arXiv:1210.1329v1 [math-ph] 4 Oct 2012

Asymptotics of the ground state energy of heavy molecules and related topics

Victor Ivrii

October 5, 2012

Abstract

We consider asymptotics of the ground state energy of heavy atoms and molecules in the strong external magnetic field and derive it including Schwinger and Dirac corrections (if magnetic field is not too strong). In the next instalment we extend this paper to consider also related topics: an excessive negative charge, ionization energy and excessive negative charge when atoms can still bind into molecules.

Contents

Contents **1**

0 Introduction **3**
0.1 Framework 3
0.2 Problems to consider 4
0.3 Magnetic Thomas-Fermi theory 5
0.4 Main results sketched and plan of the chapter 7

1 Magnetic Thomas-Fermi theory **8**
1.1 Framework and existence 8
1.2 Properties 11

2 Applying semiclassical methods: $M = 1$ 15
2.1 Heuristics 15
2.2 Smooth approximation 22
2.3 Rough approximation 27

3 Applying semiclassical methods: $M \geq 2$ 41
3.1 Analysis in zone $\mathcal{X}_2$ 41

4 Analysis in the boundary strip 59
4.1 Properties of W_B^{TF} as $N = Z$ 59
4.2 Semiclassical analysis in the strip $\mathcal{Y}$ for $N \geq Z$. 64
4.3 Analysis in the boundary strip $\mathcal{Y}$. Construction of the potential for $N < Z$ 67
4.4 Conclusion 72

5 Ground state energy 73
5.1 Lower estimates 73
5.2 Upper estimate: general scheme 75
5.3 Upper estimate as $M = 1$ 75
5.4 Upper estimate as $M \geq 2$ 78

6 Negatively charged systems (coming) 89

7 Positively charged systems (coming) 89

8 Appendices 89
8.A Electrostatic inequalities 89
8.B Very strong magnetic field case 94
8.C Riemann sums and integrals 95

0 Introduction

In this paper we repeat analysis of the previous paper [Ivr10]$^{1)}$ but in the case of the constant external magnetic field$^{2)}$.

0.1 Framework

Let us consider the following operator (quantum Hamiltonian)

$$\mathsf{H}=\mathsf{H}_N:=\sum_{1\le j\le N} H_{A,V,x_j}+\sum_{1\le j<k\le N}|x_j-x_k|^{-1} \tag{0.1.1}$$

on

$$\mathfrak{H}=\bigwedge_{1\le n\le N}\mathscr{H},\qquad \mathscr{H}=\mathscr{L}^2(\mathbb{R}^d,\mathbb{C}^q) \tag{0.1.2}$$

with

$$H_{V,A}=\bigl((i\nabla-A)\cdot\boldsymbol{\sigma}\bigr)^2-V(x) \tag{0.1.3}$$

describing N same type particles in the external field with the scalar potential $-V$ and vector potential $A(x)$, and repulsing one another according to the Coulomb law.

Here $x_j\in\mathbb{R}^d$ and $(x_1,\dots,x_N)\in\mathbb{R}^{dN}$, potentials $V(x)$ and $A(x)$ are assumed to be real-valued. Except when specifically mentioned we assume that

$$V(x)=\sum_{1\le k\le M}\frac{Z_m}{|x-\mathsf{y}_m|} \tag{0.1.4}$$

where $Z_k>0$ and y_k are charges and locations of nuclei. Here $\boldsymbol{\sigma}=(\sigma_1,\sigma_2,\sigma_d)$, σ_k are $q\times q$-Pauli matrices.

So far in comparison with the previous paper [Ivr10] we only changed (24.1.3) of [Ivr11] to (0.1.3) introducing magnetic field. Now spin enters not only in the definition of the space but also into operator through matrices

---

[1] Coinciding with Chapter 24 of of [Ivr11].

[2] Actually we need a magnetic field either sufficiently weak or close to a constant on the very small scale.

σ_k. Since we need $d=3$ Pauli matrices it is sufficient to consider $q=2$ but we will consider more general case as well (but q should be even).

Remark 0.1.1. In the case of the the constant magnetic field $\nabla\times A$

$$H_{A,V}=\bigl(-i\nabla-A(x)\bigr)^2+\boldsymbol{\sigma}\cdot\nabla\times A-V(x) \tag{0.1.5}$$

In the case $d=2$ this operator downgrades to

$$H_{A,V}=\bigl(-i\nabla-A(x)\bigr)^2+\sigma_3(\nabla\times A)-V(x) \tag{0.1.6}$$

Again, let us assume that

(0.1.7) Operator H is self-adjoint on $\mathfrak{H}$.

As usual we will never discuss this assumption.

0.2 Problems to consider

As in the previous Chapter we are interested in the *ground state energy* $\mathsf{E}=\mathsf{E}_N$ of our system i.e. in the lowest eigenvalue of the operator $\mathsf{H}=\mathsf{H}_N$ on $\mathfrak{H}$:

$$\mathsf{E}:=\inf\operatorname{Spec}\mathsf{H}\qquad\text{on}\ \ \mathfrak{H}; \tag{0.2.1}$$

more precisely we are interested in the asymptotics of $\mathsf{E}_N=\mathsf{E}(\underline{\mathsf{y}};\underline{Z};N)$ as V is defined by (0.1.4) and $N\asymp Z:=Z_1+Z_2+\ldots+Z_M\to\infty$ and we are going to prove that[3] E is equal to *Magnetic Thomas-Fermi energy* $\mathcal{E}_B^{\mathsf{TF}}$, possibly with Scott and Dirac-Schwinger corrections and with appropriate error.

We are also interested in the asymptotics for the *ionization energy*

$$\mathsf{I}_N:=-\mathsf{E}_N+\mathsf{E}_{N-1} \tag{0.2.2}$$

and we also would like to estimate *maximal excessive negative charge*

$$\max_{N:\ \mathsf{I}_N>0} N-Z. \tag{0.2.3}$$

---

[3] Under reasonable assumption to the minimal distance between nuclei.

All these questions so far were considered in the framework of the fixed positions $\mathsf{y}_1, \dots, \mathsf{y}_M$ but we can also consider

$$\widehat{\mathsf{E}} = \widehat{\mathsf{E}}_N = \widehat{\mathsf{E}}(\underline{\mathsf{y}}; \underline{Z}; N) = \mathsf{E} + U(\underline{\mathsf{y}}; \underline{Z}) \tag{0.2.4}$$

with

$$\sum_{1\le m<m'\le M} \frac{Z_m Z_{m'}}{|\mathsf{y}_m - \mathsf{y}_{m'}|} \tag{0.2.5}$$

and

$$\widehat{\mathsf{E}}(\underline{Z}; N) = \inf_{\mathsf{y}_1,\dots,\mathsf{y}_M} \widehat{\mathsf{E}}(\underline{\mathsf{y}}; \underline{Z}; N) \tag{0.2.6}$$

and replace I_N by $\widehat{I}_N = -\widehat{\mathsf{E}}_N + \widehat{\mathsf{E}}_{N-1}$ and modify all our questions accordingly. We call these frameworks *fixed nuclei model* and *free nuclei model* respectively.

In the free nuclei model we can consider two other problems:

- Estimate from below *minimal distance between nuclei* i.e.

$$\min_{1\le m<m'\le M} |\mathsf{y}_m - \mathsf{y}_{m'}|$$

for which such minimum is achieved;

- Estimate *maximal excessive positive charge*

$$\max_N \{Z - N:\ \widehat{\mathsf{E}} < \min_{\substack{N_1,\dots,N_M:\\ N_1+\dots N_M=N}} \sum_{1\le m\le M} \mathsf{E}(Z_m; N_m)\} \tag{0.2.7}$$

for which molecule does not disintegrates into atoms.

0.3 Magnetic Thomas-Fermi theory

As in the previous Chapter the first approximation is the Hartree-Fock (or Thomas-Fermi) theory. Let us introduce the *spacial density* of the particle with the state $\Psi \in \mathfrak{H}$:

$$\rho(x) = \rho_\Psi(x) = N \int |\Psi(x, x_2, \dots, x_N)|^2 \, dx_2 \cdots dx_N. \tag{0.3.1}$$

Let us write the Hamiltonian, describing the corresponding "quantum liquid":

$$\mathcal{E}_B(\rho) = \int \tau_B(\rho(x))\,dx - \int V(x)\rho(x)\,dx + \frac{1}{2}\mathsf{D}(\rho,\rho), \tag{0.3.2}$$

with

$$\mathsf{D}(\rho,\rho) = \iint |x-y|^{-1}\rho(x)\rho(y)\,dxdy \tag{0.3.3}$$

where τ_B is the energy density of a gas of noninteracting electrons:

$$\tau_B(\rho) = \sup_{w\geq 0}\bigl(\rho w - P(w)\bigr) \tag{0.3.4}$$

is the Legendre transform of the *pressure* $P_B(w)$ given by the formula

$$P_B(w) = \varkappa_1 B\Bigl(\frac{1}{2}w_+^{\frac{d}{2}} + \sum_{j\geq 1}(w-2jB)_+^{\frac{d}{2}}\Bigr) \tag{0.3.5}$$

with $\varkappa_1 = (2\pi)^{-1}q, (3\pi^2)^{-1}q$ for $d=2,3$ respectively.

The classical sense of the second and the third terms in the right-hand expression of (0.3.2) is clear and the density of the kinetic energy is given by $\tau_B(\rho)$ in the semiclassical approximation (see remark 0.3.1). So, the problem is

(0.3.6) Minimize functional $\mathcal{E}_B(\rho)$ defined by (0.3.2) under restrictions:

$$\rho \geq 0, \qquad \int \rho\,dx \leq N. \tag*{(0.3.7)$_{1,2}$}$$

The solution if exists is unique because functional $\mathcal{E}_B(\rho)$ is strictly convex (see below). The existence and the property of this solution denoted further by ρ_B^{TF} is known in the series of physically important cases.

Remark 0.3.1. If w is the negative potential then

$$\operatorname{tr} e(x,x,0) \approx P'_B(w) \tag{0.3.8}$$

defines the density of all non-interacting particles with negative energies at point x and

$$\int_{-\infty}^{0} \tau\, d_\tau \operatorname{tr} e(x,x,\tau)dx \approx -\int P_B(w)\,dx \tag{0.3.9}$$

is the total energy of these particles; here $\approx$ means "in the semiclassical approximation".

We consider in the case of $d=3$ a large (heavy) molecule with potential (24.1.4) of [Ivr11]. It is well-known$^{4)}$ that

Proposition 0.3.2. *(i) For $V(x)$ given by* (0.1.4) *minimization problem* (0.3.6) *has a unique solution $\rho=\rho_B^{\mathsf{TF}}$; then denote $\mathcal{E}_B^{\mathsf{TF}}:=\mathcal{E}_B(\rho_B^{\mathsf{TF}})$;*

(ii) Equality in $(0.3.7)_2$ *holds if and only if $N\le Z:=\sum_m Z_m$;*

(iii) Further, ρ^{TF} does not depend on N as $N\ge Z$;

(iv) Thus

$$\int \rho_B^{\mathsf{TF}}\,dx=\min(N,Z),\qquad Z:=\sum_{1\le m\le M} Z_m. \tag{0.3.10}$$

0.4 Main results sketched and plan of the chapter

In the first half of this Chapter we derive asymptotics for ground state energy and justify Thomas-Fermi theory. As construction of Section 24.2 of [Ivr11] works with minimal modifications (see Section 5) in the magnetic case as well we start immediately from magnetic Thomas-Fermi theory in Section 1. We discover that there are three different cases: a weak magnetic field case $B\ll Z^{\frac{4}{3}}$ when $\mathcal{E}_B^{\mathsf{TF}}\asymp Z^{\frac{5}{3}}$ and $\mathcal{E}_B^{\mathsf{TF}}=\mathcal{E}_0^{\mathsf{TF}}(1+o(1))$, a strong magnetic field case $B\gg Z^{\frac{4}{3}}$ when $\mathcal{E}_B^{\mathsf{TF}}\asymp B^{\frac{2}{5}}Z^{\frac{9}{5}}$ and $\mathcal{E}_B^{\mathsf{TF}}=\bar{\mathcal{E}}_B^{\mathsf{TF}}(1+o(1))$ where $\bar{\mathcal{E}}_B^{\mathsf{TF}}$ is Thomas-Fermi potential derived as $P_B(w)=\frac{1}{2}\varkappa_1 w^{\frac{d}{2}}$ (cf. (0.3.5)), and an intermediate case $B\sim Z^{\frac{4}{3}}$.

Then we apply semiclassical methods (like in Section 24.4) of [Ivr11] albeit now analysis is way more complicated due to two factors: semiclassical theory of magnetic Schrödinger operator is more difficult than the corresponding theory for non-magnetic Schrödinger operator and also Thomas-Fermi potential W^{TF} is not very smooth in the magnetic case, so we need to approximate it by a smooth one (on a microscale). We discover that both semiclassical methods and Thomas-fermi theory are relevant only as $B\ll Z^3$. Case of the superstrong magnetic field $B\gg Z^3$ was considered in E. H. Lieb, J. P. Solovej and J. Yngvarsson [LSY1] and we hope to cover it in the Chapter 27^1

^{1} Pending

[4)] Section IV of E. H. Lieb, J. P. Solovej and J. Yngvarsson [LSY2].

First of all, in Section 2 we consider the case $M=1$; then Thomas-Fermi potential W_B^{TF} is non-degenerate and in this case we derive sharp spectral asymptotics.

Next, in Section 3 we consider the case $M\ge 2$ but only in the zone $\{W_B^{\mathsf{TF}}+\nu\ge B\}$ where ν is a chemical potential and B is an intensity of the magnetic field. A certain weaker non-degeneracy condition is satisfied due to Thomas-Fermi equation and we derive almost sharp spectral asymptotics.

Then in Section 4 we consider the case $M\ge 2$ in the zone $\{W^{\mathsf{TF}}+\nu\le B\}$ containing the boundary of $\operatorname{supp}\rho_B^{\mathsf{TF}}$; this is the most difficult case to analyze and our remainder estimates are not sharp unless $N\ge Z-CZ^{\frac{2}{3}}$.

Finally, in Section 5 we derive asymptotics of the ground state energy. Their precision (or lack of it) follows from the precision of the corresponding semiclassical results; so our results in the case $M=1$ are sharp, but results in the case $M\ge 2$ (especially if $N\le Z-CZ^{\frac{2}{3}}$) are not.

[2] Pending

In the second half[2] of this Chapter we consider related problems. In Section 6 (cf. Section 24.5) of [Ivr11] we consider negatively charged systems ($N\ge Z$) and estimate both ionization energy I_N and excessive negative charge $(N-Z)_+$ (5).

In Section 7 (cf. Section 24.6) of [Ivr11] we consider positively charged systems ($N\le Z$) and estimate the remainder $|\mathsf{I}_N+\nu|$ in the formula $\mathsf{I}_N\approx -\nu$; as $M\ge 2$ we also estimate excessive positive charge $(Z-N)_+$ when atoms can be bound into molecule(5).

We will assume that

$$c^{-1}N\le Z_m\le cN\qquad \forall m=1,\dots,M. \tag{0.4.1}$$

1 Magnetic Thomas-Fermi theory

1.1 Framework and existence

The Thomas-Fermi theory is well developed in the magnetic case as well albeit in the lesser degree than in the non-magnetic one. The most important source now is Section IV of E. H. Lieb, J. P. Solovej and J. Yngvarsson [LSY2].

Again as in the previous Paper [Ivr10] to get the best lower estimate for the ground state energy (neglecting semiclassical errors) one needs to maximize functional $\Phi_{B,*}(W+\nu)$ defined by (24.3.1) of [Ivr11] albeit with the pressure $P_B(w)$ given for $d=2,3$ by (0.3.5). Formulae (24.3.2) and (24.3.3) of [Ivr11] also remain valid.

---

(5) In (magnetic) Thomas-Fermi theory both answers are 0.

Further, to get the best upper estimate (neglecting semiclassical errors) one needs to minimize functional $\Phi^*_B(\rho',\nu)$ defined by (24.3.4) of [Ivr11] where (24.3.4) of [Ivr11] remains valid with P replaced by P_B and respectively $\tau(\rho')$ replaced by $\tau_B(\rho')$ which is Legendre transformation of P_B (see (0.3.4)).

Since P_B is given by much more complicated expression (0.3.5) rather than $(24.3.6)_1$ of [Ivr11], and respectively

$$P'_B(w)=\frac{d}{2}\varkappa_1 B\Bigl(\frac{1}{2}w_+^{\frac{d}{2}-1}+\sum_{j\ge 1}(w-2jB)_+^{\frac{d}{2}-1}\Bigr) \tag{1.1.1}$$

(cf. $(24.3.6)_2$) of [Ivr11] there is no explicit expression for τ_B similar to (24.3.7) of [Ivr11].

Remark 1.1.1. (i) $B(x)=|\nabla\times A(x)|$;

(ii) From now on we will assume that $d=3$;

(iii) P_B is a strictly convex function and therefore τ_B is also a strictly convex function[6];

(iv) $P_B(w)\to P_0(w)$, $P'_B(w)\to P'_0(w)$ and $\tau_B(\rho)\to\tau_0(\rho)$ as $B\to 0$ where (without subscript "0") the limit functions have been defined by (24.3.6) and (24.3.7) of [Ivr11] respectively.

Remark 1.1.2. (i) Alternatively we minimize $\mathcal{E}_B(\rho)=\Phi^*_B(\rho,0)$ under assumptions

$$\rho\ge 0,\qquad \int \rho\,dx\le N; \tag*{$(1.1.2)_{1,2}$}$$

(ii) So far in comparison with the previous paper [Ivr10] we changed only definition of $P_B(w)$ and $\tau_B(\rho)$ respectively. Note that $P_B(w)$ belongs to $\mathscr{C}^{\frac{d}{2}}$ (as $d=2,3$) as function of w; this statement will be quantified later;

---

[6] As $d=2$, P_B is a convex piecewise linear function and therefore τ_B is also a convex function.

(iii) While not affecting existence (with equality in $(1.1.2)_1$ iff $N \le Z$) and uniqueness of solution, it affects other properties, especially as $B \ge Z^{\frac{4}{3}}$.

Proposition 1.1.3. *In our assumptions for any fixed $\nu \le 0$ statements (i)–(vii) of proposition 24.3.1 of [Ivr11] hold.*

Proof. The proof is the same as of proposition 24.3.1 of [Ivr11]. The proof that threshold $\nu = 0$ matches to $N = Z$ are theorems 4.9 and 4.10 of Section IV of E. H. Lieb, J. P. Solovej and J. Yngvarsson [LSY2]. ☐

Note that (24.3.8)–(24.3.9) and (24.3.10) of [Ivr11] become

$$\rho = \frac{1}{4\pi}\Delta(W - V) = P'_B(W + \nu), \tag{1.1.3}$$

$$W = o(1) \qquad \text{as}\quad |x| \to \infty \tag{1.1.4}$$

and

$$\mathcal{N}(\nu) = \int P'_B(W + \nu)\, dx \tag{1.1.5}$$

respectively.

Similarly, proposition 24.3.2 of [Ivr11] remains true:

Proposition 1.1.4. *For arbitrary W the following estimates hold with absolute constants $\epsilon_0 > 0$ and C_0:*

$$\epsilon_0 \mathsf{D}(\rho - \rho^{\mathsf{TF}}, \rho - \rho^{\mathsf{TF}}) \le \Phi_{B,*}(W^{\mathsf{TF}} + \nu) - \Phi_{B,*}(W + \nu) \le C_0 \mathsf{D}(\rho - \rho', \rho - \rho') \tag{1.1.6}$$

and

$$\epsilon_0 \mathsf{D}(\rho' - \rho^{\mathsf{TF}}, \rho' - \rho^{\mathsf{TF}}) \le \Phi^*_B(\rho, \nu) - \Phi^*_B(\rho^{\mathsf{TF}}, \nu) \le C_0 \mathsf{D}(\rho - \rho', \rho - \rho') \tag{1.1.7}$$

with $\rho = \frac{1}{4\pi}\Delta(W - V)$, $\rho' = P'_B(W + \nu)$.

Proof. This proof is rather obvious as well. ☐

1.2 Properties

Proposition 1.2.1. *The solution of the magnetic Thomas-Fermi problem has the following scaling properties*

$$W^{\mathsf{TF}}(x;\, \underline{Z};\, \underline{y};\, B;\, N;\, q) = q^{\frac{2}{3}} N^{\frac{4}{3}} W^{\mathsf{TF}}(q^{\frac{2}{3}} N^{\frac{1}{3}} x;\, N^{-1}\underline{Z};\, q^{\frac{2}{3}} N^{\frac{1}{3}} \underline{\mathsf{y}};\, q^{-\frac{2}{3}} N^{-\frac{4}{3}} B;\, 1; 1), \tag{1.2.1}$$

$$\rho^{\mathsf{TF}}(x;\, \underline{Z};\, \underline{y};\, B;\, N;\, q) = q^{2} N^{2} \rho^{\mathsf{TF}}(q^{\frac{2}{3}} N^{\frac{1}{3}} x;\, N^{-1}\underline{Z};\, q^{\frac{2}{3}} N^{\frac{1}{3}} \underline{\mathsf{y}};\, q^{-\frac{2}{3}} N^{-\frac{4}{3}} B;\, 1; 1), \tag{1.2.2}$$

$$\mathcal{E}^{\mathsf{TF}}(\underline{Z};\, \underline{y};\, B;\, N;\, q) = q^{\frac{2}{3}} N^{\frac{7}{3}} \mathcal{E}^{\mathsf{TF}}(N^{-1}\underline{Z};\, q^{\frac{2}{3}} N^{\frac{1}{3}} \underline{\mathsf{y}};\, q^{-\frac{2}{3}} N^{-\frac{4}{3}} B;\, 1;\, 1), \tag{1.2.3}$$

$$\nu^{\mathsf{TF}}(\underline{Z};\, \underline{y};\, B;\, N;\, q) = q^{\frac{2}{3}} N^{\frac{4}{3}} \nu^{\mathsf{TF}}(N^{-1}\underline{Z};\, q^{\frac{2}{3}} N^{\frac{1}{3}} \underline{\mathsf{y}};\, q^{-\frac{2}{3}} N^{-\frac{4}{3}} B;\, 1;\, 1) \tag{1.2.4}$$

where $\nu^{\mathsf{TF}} = \nu$ *is the chemical potential; recall that* $\underline{Z} = (Z_1, \dots, Z_M)$ *and* $\underline{\mathsf{y}} = (\mathsf{y}_1, \dots, \mathsf{y}_M)$ *are arrays and parameter* q *also enters into Thomas-Fermi theory.*

In particular, ν^{TF} *and* B *scale the same way.*

Proof. Proof is trivial by scaling. $\square$

Now one can guess that there are two cases $B \ll Z^{\frac{4}{3}}$ and $B \gg Z^{\frac{4}{3}}$ (recall that $N \asymp Z$) in which magnetic Thomas-Fermi theory looks very different (and also an intermediate case $B \sim Z^{\frac{4}{3}}$). To explain this difference let us consider one atom case:



First of all recall that if $B = 0$ and $N = Z$ theory (as $M = 1$) has just one parameter and we can get rid off it by rescaling; $W^{\mathsf{TF}} \asymp Z\ell^{-1}$ as $\ell \lesssim Z^{-\frac{1}{3}}$ and $W^{\mathsf{TF}} \asymp \ell^{-4}$ as $\ell \gtrsim Z^{-\frac{1}{3}}$. Then

$$\mathsf{W}^{\mathsf{TF}\,\frac{3}{2}} \ell^{3} \asymp Z^{\frac{3}{2}} \ell^{\frac{3}{2}}, \qquad \mathsf{W}^{\mathsf{TF}\,\frac{5}{2}} \ell^{3} \asymp Z^{\frac{5}{2}} \ell^{\frac{1}{2}}$$

and

$$\mathsf{W}^{\mathsf{TF}\,\frac{3}{2}} \ell^{3} \asymp \ell^{-3}, \qquad \mathsf{W}^{\mathsf{TF}\,\frac{5}{2}} \ell^{3} \asymp \ell^{-7}$$

respectively where the first factors are spacial densities of the charge and (negative) Thomas-Fermi energy respectively and therefore zone $\ell \asymp Z^{-\frac{1}{3}}$ provides the main contributions into both.

Therefore, if in this *main zone* $B \lll W^{\mathsf{TF}} \asymp Z^{\frac{4}{3}}$ we guess that the magnetic theory is similar to non-magnetic one, and actually it is true.

However, let us study an atomic case rigorously. Let $M = 1$, $\mathsf{y}_m = 0$ and $N \le Z$. Then

(1.2.5) W_B^{TF} is a spherically symmetric, and it is monotone non-increasing function of $|x|$; $W_B^{\mathsf{TF}} \to +0$ as $|x| \to \infty$;

$$W_B^{\mathsf{TF}}(x) \le -\nu \implies W_B^{\mathsf{TF}} = |x|^{-1}(Z - N). \tag{1.2.6}$$

Really, (1.2.5) is obvious and (1.2.6) follows from it and Newton screening theorem.

Two propositions below treat cases $B \lesssim Z^{\frac{4}{3}}$ and $B \gtrsim Z^{\frac{4}{3}}$ respectively; in the former case there is another fork: $B \lesssim (Z - N)_+^{\frac{4}{3}}$ and $B \gtrsim (Z - N)_+^{\frac{4}{3}}$.

Proposition 1.2.2. *Let* $M = 1$, $\mathsf{y}_m = 0$, $N \asymp Z_m$ *and* $B \le Z^{\frac{4}{3}}$.

(i) Then

$$W_B^{\mathsf{TF}} \le \min(Z|x|^{-1}, C|x|^{-4}) \tag{1.2.7}$$

and

$$\rho_B^{\mathsf{TF}} \le C\min(Z^{\frac{3}{2}}|x|^{-\frac{3}{2}} + BZ^{\frac{1}{2}}|x|^{-\frac{1}{2}}, |x|^{-6} + B|x|^{-2}); \tag{1.2.8}$$

(ii) There exists

$$r_m \asymp \min\bigl(B^{-\frac{1}{4}}, (Z - N)_+^{-\frac{1}{3}}\bigr) \tag{1.2.9}$$

such that $W_B^{\mathsf{TF}} \gtreqless -\nu$ *as* $|x| \lesseqgtr r_m$ *and then* $\rho_B^{\mathsf{TF}} = 0$ *iff* $x \ge r_m$*;*

(iii) (1.2.7) *and* (1.2.8) *become equivalencies* ($\asymp$) *as* $|x| \le (1 - \epsilon)r_m$*;*

(iv) $B \le (Z - N)_+^{\frac{4}{3}}$ *implies* $r_m \asymp (Z - N)_+^{-\frac{1}{3}}$, $\nu \asymp (Z - N)_+^{\frac{4}{3}}$ *and*

$$\begin{aligned} &W^{\mathsf{TF}} + \nu \asymp (Z - N)_+^{\frac{5}{3}}(r_m - |x|),\\ &-\partial_{|x|}W^{\mathsf{TF}} \asymp (Z - N)_+^{\frac{5}{3}} \qquad \text{as } (1 - \epsilon)r_m \le |x| \le r_m; \end{aligned} \tag{1.2.10}$$

(v) $B \ge (Z-N)_+^{\frac{4}{3}}$ *implies* $r_m \asymp B^{-\frac{1}{4}}$, $\nu \asymp (Z-N)_+ B^{\frac{1}{4}} \lesssim B$ *and*

$$\begin{aligned} &W^{\mathsf{TF}} + \nu \asymp B^2 (r_m - |x|)^4 + B^{\frac{1}{2}} (Z-N)_+ (r_m - |x|) \\ &-\partial_{|x|} W^{\mathsf{TF}} \asymp B^2 (r_m - |x|)^3 + B^{\frac{1}{2}} (Z-N)_+ \\ &\qquad\qquad\qquad\qquad\qquad\qquad \text{as} \quad (1-\epsilon) r_m \le |x| \le r_m. \end{aligned} \tag{1.2.11}$$

Proposition 1.2.3. *Let* $M=1$, $\mathsf{y}_m = 0$, $N \asymp Z_m$ *and* $B \ge Z^{\frac{4}{3}}$.

(i) Then

$$W_B^{\mathsf{TF}} \le Z|x|^{-1} \tag{1.2.12}$$

and

$$\rho_B^{\mathsf{TF}} \le CZ^{\frac{3}{2}} |x|^{-\frac{3}{2}} + CBZ^{\frac{1}{2}} |x|^{-\frac{1}{2}}; \tag{1.2.13}$$

(ii) There exist r_m *and* r'_m,

$$r_m \asymp B^{-\frac{2}{5}} Z^{\frac{1}{5}}, \qquad r'_m \asymp B^{-1} Z_m, \tag{1.2.14}$$

such that $W_B^{\mathsf{TF}} \gtrless B$ *as* $|x| \lesseqgtr r_m$, $W_B^{\mathsf{TF}} \gtrless -\nu$ *as* $|x| \lesseqgtr r'_m$ *and then* $\rho_B^{\mathsf{TF}} = 0$ *iff* $x \ge r_m$*;*

(iii) (1.2.12)–(1.2.13) *become equivalencies* ($\asymp$) *as* $|x| \le (1-\epsilon) r_m$*;*

(iv) $\nu \asymp (Z-N)_+ B^{\frac{2}{5}} Z^{-\frac{1}{5}} \lesssim B$ *and*

$$\begin{aligned} &W^{\mathsf{TF}} + \nu \asymp B^2 (r_m - |x|)^4 + r_m^{-2} (Z-N)_+ (r_m - |x|) \\ &-\partial_{|x|} W^{\mathsf{TF}} \asymp B^2 (r_m - |x|)^3 + r_m^{-2} (Z-N)_+ \\ &\qquad\qquad\qquad\qquad\qquad\qquad \text{as} \quad (1-\epsilon) r_m \le |x| \le r_m. \end{aligned} \tag{1.2.15}$$

Proofs of propositions 1.2.2 and 1.2.3. Proofs easily follow from equation and "boundary conditions" satisfied by $w(r)$ where $r = |x|$:

$$w'' + 2r^{-1} w = P'_B(w+\nu), \tag{1.2.16}$$

$$w = r^{-1} Z_m + O(1) \qquad \text{as} \quad r \to 0, \tag{1.2.17}$$

$$w(r_m) = -\nu, \qquad w'(r) = \nu r_m^{-1} \tag{1.2.18}$$

where $\nu = -(Z_m - N)_+ \bar{r}_m^{-1}$. □

Corollary 1.2.4. *Let* $M=1$, $\mathsf{y}_m=0$ *and* $N\asymp Z_m$. *Then*

(i) $W_B^{\mathsf{TF}}\lesssim B$ *if* $|x|\ge r'_m$ *where* $r'_m\asymp B^{-1}Z_m$ *as* $B\ge Z_m^{\frac{4}{3}}$ *and* $r'_m\asymp B^{-\frac{1}{4}}$ *as* $B\le cZ_m^{\frac{4}{3}}$*;*

(ii) As $B\lesssim Z_m^{\frac{4}{3}}$ *the main contribution to both the charge and the Thomas-Fermi energy is delivered by zone* $\{x: |x|\asymp r_m^*\}$ *with* $r_m^*=Z_m^{-\frac{1}{3}}$*; in particular, then* $\mathcal{E}_B^{\mathsf{TF}}\asymp\mathcal{E}^{\mathsf{TF}}\asymp Z_m^{\frac{7}{3}}$*; further, in this case* $W_B^{\mathsf{TF}}\asymp W^{\mathsf{TF}}$ *in the zone* $\{x:\ |x|\lesssim\epsilon r_m\}$*;*

(iii) Further, $\mathcal{E}_B^{\mathsf{TF}}\sim\mathcal{E}^{\mathsf{TF}}$ *as* $B\ll Z_m^{\frac{4}{3}}$*; furthermore, in this case* $W_B^{\mathsf{TF}}\sim W^{\mathsf{TF}}$ *in the zone* $\{x:\ |x|\ll r_m\}$*;*

(iv) On the other hand, as $B\ge Z^{\frac{4}{3}}$*, the main contributions to the total charge and energy are delivered by* $\{x:\ |x|\asymp r_m\}$ *and in particular* $Z_m\asymp BZ_m^{\frac{1}{2}}r_m^{\frac{5}{2}}$ *and*

$$\mathcal{E}_B^{\mathsf{TF}}\asymp BZ_m^{\frac{3}{2}}r_m^{\frac{3}{2}}\asymp B^{\frac{2}{5}}Z_m^{\frac{9}{5}}; \tag{1.2.19}$$

Recall that $r_m\asymp B^{-\frac{1}{4}}$ as $B\le Z_m^{\frac{4}{3}}$ and $r_m\asymp B^{-\frac{2}{5}}Z_M^{\frac{1}{5}}$ as $B\ge Z_M^{\frac{4}{3}}$. Note that proposition 24.3.5 of [Ivr11] (comparing W^{TF} for molecule with the sum of those for single atoms) still holds. Therefore we conclude that

Corollary 1.2.5. *(i) Assume that* $Z_m\asymp Z$ *for all* $m=1,\dots,M$*. Then all statements of corollary 1.2.4 remain true for* $M\ge 2$ *with* $|x|$ *and* Z_m *replaced by* $\ell(x)$ *and* Z *and* r_m, r'_m, r^*_m *by* r, r', r^* *respectively;*

(ii) In the general case global statements remain true, pointwise statements remain true without modification only as $\ell(x)=\ell_m(x):=|x-\mathsf{y}_m|$ *with* $Z_m\asymp Z$*.*

Also holds proposition 24.3.12 of [Ivr11] as it uses only super-additivity of $\tau(\rho)$ and $\tau_B(\rho)$ is also super-additive (this follows from convexity of $\tau_B(\rho)$ and equality $\tau_B(0)=0$).

However there is a significant difference: if there is no magnetic field atoms really repulse one another on any distances and we can attribute it to either excessive positive charge as $N<Z$ or their infinite spatial size as $N=Z$. However with magnetic field atoms have a finite size even as $N=Z$ and they do not repulse one another on the large distances. In particular,

Proposition 1.2.6. *Let* $N = Z$ *and*

$$|\mathsf{y}_m - \mathsf{y}_{m'}| \ge \bar{r}_m + \bar{r}_{m'} \qquad \forall m:\ 1 \le m < m' \le M. \tag{1.2.20}$$

Then

$$\mathcal{E}_B^{\mathsf{TF}}(\underline{Z}, \underline{\mathsf{y}}, B, Z) = \sum_{1\le m\le M} \mathcal{E}_B^{\mathsf{TF}}(Z_m, \mathsf{y}_m, B, Z_m) \tag{1.2.21}$$

and

$$\rho_B^{\mathsf{TF}}(x, \underline{Z}, \underline{\mathsf{y}}, B, Z) = \sum_{1\le m\le M} \rho_B^{\mathsf{TF}}(x, Z_m, \mathsf{y}_m, B, Z_m). \tag{1.2.22}$$

As far as we know Theorem 1 of R. Benguria [Be] (see Theorem 24.3.8 of [Ivr11]) has not been proven in the case of magnetic field; while it is not important for our arguments we still formulate

Problem 1.2.7. *Prove (or disprove) Theorem 24.3.8 of [Ivr11] in the case of the constant magnetic field.*

2 Applying semiclassical methods: $M = 1$

2.1 Heuristics

Let us consider first a mock proof of our main results; we deal here *as if* W_B^{TF} *was very smooth* which it is not the case; however later we will show that its smoothness is sufficient to employ arguments of Chapter 18 rather than those of Chapter 13 of [Ivr11]. We also will deal as if non-degeneracy conditions were satisfied leaving them also to more rigorous arguments below.

2.1.1 Total charge.

Consider

$$\int e(x,x,\nu)\psi(x)\,dx \tag{2.1.1}$$

first with γ-admissible $\psi(x)$, where $\gamma \le \epsilon\ell$.

General arguments.

The main part of semiclassical expression for (2.1.1) is of magnitude $\hbar'^{-3} + \mu'\hbar'^{-2} \asymp \zeta^3\gamma^3 + B\zeta\gamma^3$ with $\hbar' = 1/(\zeta\gamma)$ and $\mu' = B\gamma/\zeta$. Really, let us rescale $x \mapsto x/\gamma$ and $\tau \mapsto \tau/\zeta^2$ which leads to $h = 1 \mapsto \hbar'$ and $B \mapsto \mu'$. In particular, as $\gamma \asymp \ell$ we get

$$\zeta^3\ell^3 + B\zeta\ell^3. \tag{2.1.2}$$

Meanwhile the remainder in the semiclassical expression for (2.1.1) does not exceed $C\hbar'^{-2} + C\mu'\hbar'^{-2} \asymp \zeta^2\gamma^2 + B\gamma^2$ (gaining factor $\hbar'$ in the comparison to the main part; here we need smoothness and as $\mu' \ge \hbar'^{\delta-1}$ we need non-degeneracy); as $\gamma \asymp \ell$ we get

$$\zeta^2\ell^2 + B\ell^2. \tag{2.1.3}$$

Sure, we ignored the fact that $\hbar' \le 1$ does not necessarily hold even if $\gamma \asymp \ell$ but we believe that the contributions to the main part and the remainder of these zones will be less than of zone where this inequality holds, provided $B \ll Z^3$.

Finally let us sum with respect to ℓ-partition.

Moderate magnetic field.

Consider $B \le Z^{\frac{4}{3}}$ first. Then for $\ell \le Z^{-\frac{1}{3}}$ we plug $\zeta = Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}$ into (2.1.2) and (2.1.3) resulting in

$$Z^{\frac{3}{2}}\ell^{\frac{3}{2}} + BZ^{\frac{1}{2}}\ell^{\frac{5}{2}} \qquad \text{and} \qquad Z\ell + B\ell^2 \tag*{(2.1.4)$_{0,1}$}$$

in the main part and in the remainder respectively and the summation over zone $\{\ell \le Z^{-\frac{4}{3}}\}$ results in the same expressions as $\ell = Z^{-\frac{1}{3}}$ i. e. $Z + BZ^{-\frac{1}{3}} \asymp Z$ and $Z^{\frac{2}{3}} + BZ^{-\frac{2}{3}} \asymp Z^{\frac{2}{3}}$ respectively.

Further, for $\ell \ge Z^{-\frac{1}{3}}$ we plug $\zeta = \ell^{-2}$ into (2.1.2) and (2.1.3) resulting in

$$\ell^{-3} + B\ell \qquad \text{and} \qquad \ell^{-2} + B\ell^2; \tag*{(2.1.5)$_{0,1}$}$$

then summation over zone $\{Z^{-\frac{1}{3}} \le \ell \le \bar{r} = B^{-\frac{1}{4}}\}$ results in $Z + B^{\frac{3}{4}} \asymp Z$ and $Z^{\frac{2}{3}} + B^{\frac{1}{2}} \asymp Z^{\frac{2}{3}}$ respectively.

Strong magnetic field.

Consider $B \ge Z^{\frac{4}{3}}$ now. Then threshold $Z^{-\frac{1}{3}}$ disappears and we sum expressions $(2.1.4)_{0,1}$ over $\ell \le \bar{r} := Z^{\frac{1}{5}}B^{-\frac{2}{5}}$ resulting in $Z^{\frac{9}{5}}B^{-\frac{3}{5}} + Z \asymp Z$ and $Z^{\frac{6}{5}}B^{-\frac{2}{5}} + Z^{\frac{2}{5}}B^{\frac{1}{5}} \asymp Z^{\frac{2}{5}}B^{\frac{1}{5}}$.

Therefore,

(2.1.6) The total charge is $\min(N, Z)$ (due to the choice of ν) with the remainder estimate $O\bigl(\max(Z^{\frac{2}{3}}, Z^{\frac{2}{5}}B^{\frac{1}{5}})\bigr)$ which is $O(Z^{\frac{2}{3}})$ as $B \le Z^{\frac{4}{3}}$ and $O(Z^{\frac{2}{5}}B^{\frac{1}{5}})$ as $Z^{\frac{4}{3}} \le B \le Z^{3}$.

Remark 2.1.1. Remainder is less than the main part as $Z^{\frac{2}{5}}B^{\frac{1}{5}} \lesssim Z$ i.e. $B \le Z^{3}$. It means exactly that $\zeta\ell \ge 1$ as $\ell = \bar{r}$ (as $B \ge Z^{\frac{4}{3}}$). The same is true for all other semiclassical asymptotics below.

2.1.2 Semiclassical D-term.

Consider now the semiclassical D-term

$$CD\bigl(e(x,x,\nu) - \rho_B^{\mathsf{TF}}(x),\ e(x,x,\nu) - \rho_B^{\mathsf{TF}}(x)\bigr). \tag{2.1.7}$$

General arguments.

We do not have appropriate asymptotics for $e(x,x,\nu)$ in the case of magnetic field[7] but we apply Fefferman–de Llave decomposition (16.3.1 of [Ivr11]):

$$|x-y|_{\gamma}^{-1}(x,y) := |x-y|\varphi(\gamma^{-1}|x-y|) = \gamma^{-4}\int \psi_{1,\gamma}(x,z)\psi_{2,\gamma}(y,z)\,dz \tag{2.1.8}$$

where $\varphi \in \mathscr{C}^\infty([1,2])$.

Therefore contribution of $B(z,\gamma)\times B(z',\gamma)$ with $3\gamma \le |z-z'| \le 4\gamma$, $\gamma \le \epsilon\ell(z)$ to such term does not exceed $\bigl(\zeta^2\gamma^2 + B\gamma^2\bigr)^2\gamma^{-1}$. There are $\asymp \ell^3\gamma^{-3}$ of such pairs with $\ell(x) \asymp \ell$ and their total contribution does not exceed $C\bigl(\zeta^2 + B\bigr)^2\ell^3$.

Now we need to sum over $\gamma^{-1}\,d\gamma$ which does not look good as leads to the logarithmic divergency but there is a simple remedy: we treat this way only pairs $t\ell \le |z-z'| \le \ell$ and apply for pairs with $|z-z'| \le t\ell$ pointwise asymptotics; then we get

---

[7] Unless we really assume that W is smooth and apply results sections 16.5–16.8 of [Ivr11].

$$C(\zeta^2+B)^2\ell^3\bigl(1+(\log B\ell/\zeta)_+\bigr); \tag{2.1.9}$$

to get rid off this logarithmic factor we apply more delicate arguments similar to those of Subsection 16.9.3 of [Ivr11].

So, ignoring this logarithmic factor we conclude that the contribution of all pairs (z,z') with $\ell(z)\asymp \ell(z')\asymp \ell$ does not exceed $C(\zeta^2+B)^2\ell^3$ while contribution of all pairs (z,z') with $\ell(z)\asymp \ell_1 \not\asymp \ell(z')\asymp \ell_2$ does not exceed $C(\zeta_1^2+B)(\zeta_2^2+B)\ell_1^2\ell_2^2(\ell_1+\ell_2)^{-1}$.

Finally let us sum.

Moderate magnetic field.

Consider $B\le Z^{\frac{4}{3}}$. Then summation over zone $\{\ell_1\le Z^{-\frac{1}{3}}, \ell_2\le Z^{-\frac{1}{3}}\}$ results in $CZ^{\frac{5}{3}}$ and the same is true for summation over zone $\{Z^{-\frac{1}{3}}\le \ell_1\le B^{-\frac{1}{4}}, Z^{-\frac{1}{3}}\le \ell_2\le B^{-\frac{1}{4}}\}$.

Obviously, in such estimates, if there is a fixed number of zones, we do not need to sum over "mixed" pairs .

Strong magnetic field.

Consider $B\ge Z^{\frac{4}{3}}$. Then summation over zone $\{\ell_1\le Z^{\frac{1}{5}}B^{-\frac{2}{5}}, \ell_2\le Z^{\frac{1}{5}}B^{-\frac{2}{5}}\}$ results in $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$.

Therefore

(2.1.10) Term (2.1.7) does not exceed $C\max(Z^{\frac{5}{3}}, Z^{\frac{3}{5}}B^{\frac{4}{5}})$ which is $CZ^{\frac{5}{3}}$ as $B\le Z^{\frac{4}{3}}$ and $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$ as $Z^{\frac{4}{3}}\le B\le Z^3$.

2.1.3 $|\lambda_N-\nu|$ and another D-term.

Consider two other non-trace terms in the upper estimate.

Moderate magnetic field.

As $B\le Z^{\frac{4}{3}}$ we established the remainder in the total charge $O(Z^{\frac{2}{3}})$. Then using our standard arguments we conclude easily that $|\lambda_N-\nu|=O(Z)$ and then

$$|\lambda_N-\nu|\cdot|\mathsf{N}(\nu)-N|\le CZ^{\frac{5}{3}} \tag{2.1.11}$$

and

$$\begin{multline}\mathsf{D}\bigl(P'_B(W_B^{\mathsf{TF}}(x)+\lambda_N)-P'_B(W_B^{\mathsf{TF}}(x)+\nu),\\ P'_B(W_B^{\mathsf{TF}}(x)+\lambda_N)-P'_B(W_B^{\mathsf{TF}}(x)+\nu)\bigr)\le CZ^{\frac{5}{3}};\end{multline} \tag{2.1.12}$$

combining with the estimate of the previous subsubsection we conclude that

$$\mathsf{D}(\rho_\Psi-\rho_B^{\mathsf{TF}},\rho_\Psi-\rho_B^{\mathsf{TF}})\le CQ=O(Z^{\frac{5}{3}}) \tag{2.1.13}$$

as in (24.4.55) of [Ivr11].

Strong magnetic field.

Let now $Z^{\frac{4}{3}}\le B\le Z^3$. Then we established the remainder in the total charge $O(Z^{\frac{2}{5}}B^{\frac{1}{5}})$ and for the semiclassical D-term we established estimate $O(Z^{\frac{3}{5}}B^{\frac{4}{5}})$ and therefore to estimate

$$|\lambda_N-\nu|\cdot|\mathsf{N}(\nu)-N|\le CZ^{\frac{3}{5}}B^{\frac{4}{5}} \tag{2.1.14}$$

as well we want to prove that

$$|\lambda_N-\nu|=O(Z^{\frac{1}{5}}B^{\frac{3}{5}}). \tag{2.1.15}$$

Note that $|\nu|\lesssim Z\bar{r}^{-1}\asymp Z^{\frac{4}{5}}B^{\frac{2}{5}}\le CB$. Therefore if $|\lambda_N-\nu|\le\frac{1}{2}|\nu|$ we conclude that

$$\begin{multline}|\int\bigl(P'_B(W_B^{\mathsf{TF}}(x)+\lambda_N)-P'_B(W_B^{\mathsf{TF}}(x)+\nu)\bigr)\,dx|\ge\\ \epsilon|\lambda_N-\nu|B\int(W+\nu)_+^{-\frac{1}{2}}\,dx\end{multline} \tag{2.1.16}$$

with integral taken over zone $\{x: W(x)+\nu\ge|\lambda_N-\nu|\}$.

One can see easily that as $|\lambda_N-\nu|\le\epsilon|\nu|$ the right-hand expression of (2.1.16) is larger than $\epsilon|\lambda_N-\nu|\cdot Z^{\frac{1}{5}}B^{-\frac{2}{5}}$ and it must be less than $CZ^{\frac{2}{5}}B^{\frac{1}{5}}$: $|\lambda_N-\nu|Z^{\frac{1}{5}}B^{-\frac{2}{5}}\le CZ^{\frac{2}{5}}B^{\frac{1}{5}}$ which implies (2.1.15).

Let us estimate the left-hand expression of (2.1.12). For this however estimate (2.1.15) is insufficient. We consider here only atomic case. Then using (1.2.15) one can prove easily that the right-hand expression of (2.1.16) is of magnitude

$$|\lambda_N-\nu|\cdot B\bar{r}^2\times\underbrace{(|\nu|\bar{r}^{-1})^{-\frac{1}{3}}B^{-\frac{1}{3}}}_{}\asymp|\lambda_N-\nu|\cdot|\nu|^{-\frac{1}{3}}Z^{\frac{7}{15}}B^{-\frac{4}{15}}$$

provided $|\lambda-N|\le\epsilon\nu$, where the selected factor is just integral of $(B^2z^4+|\nu|\bar{r}^{-1})_+^{-\frac{1}{2}}\,dz$ (appearing due to (1.2.15)). Comparing with $Z^{\frac{2}{5}}B^{\frac{1}{5}}$ we conclude that

(a) If $|\nu|\ge C_1Z^{-\frac{1}{10}}B^{\frac{7}{10}}$ $(=Z^{\frac{2}{5}}B^{\frac{3}{5}}\times Z^{-\frac{3}{10}}B^{\frac{1}{10}})$ then

$$|\lambda_N-\nu|\le C|\nu|^{\frac{1}{3}}Z^{-\frac{1}{15}}B^{\frac{7}{15}}\tag{2.1.17}$$

which is less than $\epsilon|\nu|$ and coincides with (2.1.15) as $(Z-N)_+\asymp Z$;

(b) If $|\nu|\ge C_1Z^{-\frac{1}{10}}B^{\frac{7}{10}}$ then $|\lambda_N-\nu|\le C_2Z^{-\frac{1}{10}}B^{\frac{7}{10}}$.

In the former case one can prove easily that the left-hand expression of (2.1.12) does not exceed $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$.

In the latter case (as in Subsection 24.4.2) of [Ivr11] we consider Thomas-Fermi theory with $\nu=0$ i.e. $N=Z$ and also prove that that

(2.1.18) The left-hand expression of (2.1.15) does not exceed $Q=CZ^{\frac{3}{5}}B^{\frac{4}{5}}$.

In particular we slightly improve estimate (2.1.14) to $|\nu|^{\frac{1}{3}}Z^{\frac{3}{5}}B^{\frac{2}{3}}$ as well (if $(Z-N)\ll Z$).

Therefore in our framework we estimated all non-trace terms in the upper estimate by $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$ and therefore "proved" estimate

$$\mathsf{D}(\rho_\Psi-\rho_B^{\mathsf{TF}},\rho_\Psi-\rho_B^{\mathsf{TF}})\le CQ=O(Z^{\frac{3}{5}}B^{\frac{4}{5}}).\tag{2.1.19}$$

2.1.4 Trace

Consider now $\mathsf{Tr}^-(H_{A,W}-\nu)$. This term is of magnitude $\int(\zeta^5+B\zeta^3)\,dx$ and one can see easily that it is $\asymp Z^{\frac{7}{3}}$ as $B\le Z^{\frac{4}{3}}$ and $\asymp B^{\frac{2}{5}}Z^{\frac{9}{5}}$ as $Z^{\frac{4}{3}}\le B\le Z^3$.

Meanwhile consider the remainder. Again for simplicity consider only atomic case. As $B\le Z$ contribution of zone $\{x:\,\ell(x)\le Z^{-\frac{1}{3}}\}$ is $O(Z^{\frac{5}{3}})$ (we need to include Scott correction term in the main part) while contribution of zone $\{x:\ \ell(x)\ge Z^{-\frac{1}{3}}\}$ does not exceed

$$C\int (\zeta^3+B\zeta)\ell^{-2}\,dx \tag{2.1.20}$$

taken over this zone and it is $\asymp Z^{\frac{5}{3}}$ as well.

As $Z\le B\le B^2$ contribution of zone $\{x:\ \ell(x)\le b:=B^{-\frac{2}{3}}Z^{\frac{1}{3}}\}$ is $O(b^{-\frac{1}{2}}Z^{\frac{3}{2}})=O(Z^{\frac{4}{3}}B^{\frac{1}{3}})$ and we need to include Scott correction term. Meanwhile contribution of zone $\{x:\ \ell(x)\ge b\}$ does not exceed integral (2.1.20) taken over this zone which is $\asymp Z^{\frac{4}{3}}B^{\frac{1}{3}}+Z^{\frac{3}{5}}B^{\frac{4}{5}}$ where the last term coincides with estimate for (2.1.7) as $B\ge Z^{\frac{4}{3}}$ and does not exceed $CZ^{\frac{5}{3}}$ as $B\le Z^{\frac{4}{3}}$.

Finally, as $Z^2\le B\le B^3$ we need to reset $b=Z^{-1}$ as $\hbar=1/(\zeta\ell)$ becomes ≤ 1 inside; then we do not need Scott correction term and contribution of zone $\{x:\ \ell(x)\le b\}$ to both main part and remainder do not exceed $C\int(\zeta^5+B\zeta^3)\,dx\asymp Z^2+B\asymp B$ and contribution of zone $\{x:\ \ell(x)\ge b\}$ to the remainder does not exceed integral (2.1.20) taken over this zone which results in $CB+CZ^{\frac{3}{5}}B^{\frac{4}{5}}$ and the second term dominates due to assumption $B\ll Z^3$. Thus we arrive to

(2.1.21) The main therm in $\mathsf{Tr}^-(H_{A,W}-\nu)$ is of magnitude $Z^{\frac{7}{3}}$ as $B\le Z^{\frac{4}{3}}$ and $B^{\frac{2}{5}}Z^{\frac{9}{5}}$ as $Z^{\frac{4}{3}}\le B\le Z^3$ while remainder estimate is $O(Z^{\frac{5}{3}})$ as $B\le Z$, $O(Z^{\frac{4}{3}}B^{\frac{1}{3}})$ as $Z\le B\le Z^{\frac{4}{3}}$, $O(Z^{\frac{4}{3}}B^{\frac{1}{3}}+Z^{\frac{3}{5}}B^{\frac{4}{5}})$ as $Z^{\frac{4}{3}}\le B\le Z^3$.

As $B\le Z^{\frac{7}{4}}$ we need to include into main part Scott correction term.

2.1.5 Discussion

Now let us formulate our expectations

Remark 2.1.2. We expect

(i) Estimate (2.1.13) as $B\le Z^{\frac{4}{3}}$ and (2.1.19) as $Z^{\frac{4}{3}}\le B\le Z^3$;

(ii) Furthermore as for $B\le Z^{\frac{4}{3}}$ the main contribution to all terms needed to derive this estimate is delivered by zone $\{x:\ \ell(x)\approx Z^{-\frac{1}{3}}\}$ and effective magnetic field is $\mu=B\ell/\zeta\approx BZ^{-1}$ we expect improved to "o" (or better) estimate (2.1.13) as $B\ll Z$ and $a\gg Z^{-\frac{1}{3}}$ [8];

---

[8] Recall that $a=\min_{1\le m<m'\le M}|\mathsf{y}_m-\mathsf{y}_{m'}|$ is the minimal distance between nuclei.

(iii) Statement (ii) should be also true for the trace term; however then we need to include Schwinger term;

(iv) The remainder estimate for the ground state energy is maximum of the remainder estimate for non-trace and trace terms; therefore we expect the same remainder estimate as in (2.1.21); statement (ii) should be also correct for the ground state energy; however then we need to include both Schwinger and Dirac terms;

(v) We expect the described remainder estimate of the trace term and the ground state energy if a is large enough; otherwise it should contain term $O(a^{-\frac{1}{2}}Z^{\frac{3}{2}})$ as $B \le Z^{\frac{7}{4}}$ and $a \ge Z^{-1}$ (and in this case we include Scott correction term).

Remark 2.1.3. The other difference between cases $B \le Z^{\frac{4}{3}}$ and $B \ge Z^{\frac{4}{3}}$ is that $\mu\hbar = B\zeta^{-2} \lesssim 1$ in the former case as $\ell(x) \le \bar{r}$; however in the latter case it happens only as $\ell(x) \le B^{-1}Z$ but in the zone $\{B^{-1}Z \le \ell(x) \le B^{-\frac{2}{5}}Z^{\frac{1}{5}}\}$ an opposite inequality holds.

2.2 Smooth approximation

An approach described in Subsection 2.1 hits two obstacles: non-smoothness of W_B^{TF} and its possible degeneration i.e. ∇W_B^{TF} is not disjoint from 0. However non-smoothness of W_B^{TF} is due to the non-smoothness of P_B. So we want to consider first the zone where we can just replace $P_B(W+\nu)$ by $P(W+\nu)$ and therefore W_B^{TF} by some smooth function W which does not necessary coincides with W^{TF}.

2.2.1 Trivial arguments

Obviously we can do this as an effective magnetic field $\mu = B\ell/\zeta \lesssim 1$. In this case we do not need assumption $W+\nu \asymp \zeta^2$ and therefore we can take $\zeta = \ell^{-4}$ as $B \lesssim Z^{\frac{4}{3}}$ and $\ell \gtrsim Z^{-\frac{1}{3}}$ and $\zeta = Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}$ in all other cases. Therefore zone in question is

$$\mathcal{X}_1 := \{x:\, \ell(x) \le \bar{r}_1\} \tag{2.2.1}$$

$$\text{with }\ \bar{r}_1 = \begin{cases} B^{-\frac{1}{3}} & \text{as }\ 1 \le B \lesssim Z,\\ B^{-\frac{2}{3}}Z^{\frac{1}{3}} & \text{as }\ Z \lesssim B \lesssim Z^2.\end{cases}$$

In this zone $\mathcal{X}_1$ for such modified W we can unleash the full power of the same smooth theory as in section 24.4 of [Ivr11] and prove the following

Proposition 2.2.1. *Let* $1\le B\le Z^2$. *Then*

(i) Contribution of zone $\mathcal{X}_1$ *defined by* (2.2.1) *to*

$$\int \Bigl(e(x,x,\nu)-P'(W(x)+\nu)\Bigr)\,dx \tag{2.2.2}$$

does not exceed $CZ^{\frac{2}{3}}$, *its contribution to*

$$\mathsf{D}\Bigl(e(x,x,\nu)-P'(W(x)+\nu),\,e(x,x,\nu)-P'(W(x)+\nu)\Bigr) \tag{2.2.3}$$

does not exceed $CZ^{\frac{5}{3}}$ *and its contribution to*

$$\int \Bigl(e_1(x,x,\nu)+P(W(x)+\nu)\Bigr)\,dx-\mathsf{Scott} \tag{2.2.4}$$

does not exceed $CZ^{\frac{5}{3}}+Ca^{-\frac{1}{2}}Z^{\frac{3}{2}}+CB^{\frac{1}{3}}Z^{\frac{3}{2}}$ [8], [9]

(ii) Further, as $B\ll Z$ *and* $a\gg Z^{-\frac{1}{3}}$ *we can recover for these contributions estimates* $CZ^{\frac{2}{3}}$, $CZ^{\frac{5}{3}}\upsilon$ *and* $CZ^{\frac{5}{3}}\upsilon$ *with*

$$\upsilon:=Z^{-\delta}+(aZ^{\frac{1}{3}})^{-\delta}+(BZ^{-1})^{\delta} \tag{2.2.5}$$

respectively where expression (2.2.4) *should be modified to*

$$\int \bigl(e_1(x,x,\nu)-P(W(x)+\nu)\bigr)\,dx-\mathsf{Scott}-\mathsf{Schwinger}; \tag{2.2.6}$$

Furthermore in this case contribution of $\mathcal{X}_1$ *to*

$$\frac{1}{2}\int \operatorname{tr} e^{\dagger}(x,y,\nu)e(x,y,\nu)\,dxdy-\mathsf{Dirac} \tag{2.2.7}$$

does not exceed $CZ^{\frac{5}{3}-\delta}$.

---

[9] As $a\lesssim Z^{-1}$ we skip Scott and reset $a=Z^{-1}$ in the remainder estimate which become CZ^2.

Remark 2.2.2. (i) So far we should use $P(.)$ instead of $P_B(.)$ but we will prove that the same results would hold for P_B as well;

(ii) In the next Subsubsections we expand this zone to one defined by $\mu \le \hbar^{-\frac{1}{3}}$ [10] but for trace term we still need a separate analysis as $\mu \lesssim 1$;

(iii) The same estimates hold as we replace in all expressions (2.2.2)–(2.2.6) P by P_B;

(iv) We assumed that $B \le Z^2$ as otherwise $\hbar \gtrsim 1$ not only in $\mathcal{X}_1$ but even in $\{x:\ W^{\mathsf{TF}} \ge B\}$;

(v) Note that as $\bar{r}_1 \gtrsim (Z-N)_+^{-\frac{1}{3}}$ this zone (and the whole analysis) could be cut short as outside zone in question $W+\nu \ge 0$. From paper [Ivr10] we already know how to deal with such irregularities;

(vi) We need to assume that $a \ge Z^{-\frac{1}{3}}$ and to include the second term $(aZ^{\frac{1}{3}})^{-\delta}$ in the definition of υ only as we estimate the trace term (2.2.4).

2.2.2 Formal expansion

Now we want to expand zone $\mathcal{X}_1$. Note first that

$$P'_B(W+\nu) - P'(W+\nu) = O(B^{\frac{3}{2}}) \tag{2.2.8}$$

and

$$P_B(W+\nu) - P(W+\nu) - \kappa_1 B^2 (W+\nu)^{\frac{1}{2}} = O(B^{\frac{5}{2}}). \tag{2.2.9}$$

Really, one can consider $P'_B(w)$ and $P_B(w)$ as Riemann sums for integrals $P'(w)$ and $P(w)$ respectively; see Appendix 8.C for details.

However under non-degeneracy assumption $|\nabla W| \asymp \zeta^2\ell^{-1}$ we can do better with the integrated expressions.

Proposition 2.2.3. *Assume that in* $B(z,\gamma)$

$$|\nabla^\alpha W| \le C_\alpha \zeta^2 \gamma^{-|\alpha|} \qquad \forall \alpha : |\alpha| \le K, \tag{2.2.10}$$

$$|\nabla W| \ge \epsilon \zeta^2 \gamma^{-1}, \tag{2.2.11}$$

---

[10] Or even to $\mu \le \hbar^{-\frac{3}{5}}$ under non-degeneracy assumption (2.2.11) with $\gamma = \ell$, in particular, in the atomic case.

and

$$B\le \zeta^2. \tag{2.2.12}$$

Then

$$\int \phi(x)\Bigl(P'_B(W(x)+\nu)-\tilde{P}'_B(W(x)+\nu)\Bigr)\,dx = O(B^2\zeta^{-1}\gamma^3), \tag{2.2.13}$$

$$\int \phi(x)\Bigl(P_B(W(x)+\nu)-\tilde{P}'_B(W(x)+\nu)\Bigr)\,dx = O(B^5\zeta^{-5}\gamma^3) \tag{2.2.14}$$

and

$$\begin{aligned}\mathsf{D}\Bigl(\phi(x)\bigl(P'_B(W(x)+\nu)-\tilde{P}'_B(W(x)+\nu)\bigr),&\\ \phi(x)\bigl(P'_B(W(x)+\nu)-\tilde{P}'_B(W(x)+\nu)\bigr)\Bigr)&=O(B^5\zeta^{-4}\gamma^5)\end{aligned} \tag{2.2.15}$$

with

$$\tilde{P}_B(w):=P(w)+\bigl(\kappa_1P''(w)B^2+\kappa_2P^{IV}B^4\bigr)\cdot\bigl(1-\varphi(w/B)\bigr) \tag{2.2.16}$$

where $\varphi\in\mathscr{C}^\infty([-2,2])$, $\varphi=1$ *on* $[-1,1]$.

Proof. Rescaling $x\mapsto x\gamma^{-1}$, $w\mapsto w\zeta^{-2}$ and therefore $B\mapsto\beta=B\zeta^{-2}$ one can reduce the case to $\gamma=\zeta=1$, $\beta\le 1$. Then estimates (2.2.13) and (2.2.14) are trivially proven by (multiple) integration by parts which integrates P_β on each step increasing its smoothness[11)]

To prove estimate (2.2.15) we apply decomposition (2.1.8). Integration by parts shows that (2.2.13) with t-admissible function ϕ is $O(\beta^3t^{\frac{3}{2}})$ as $t\ge\beta$ and therefore contribution of zone $\{(x,y):\,|x-y|\asymp t\}$ is $O(\beta^6t^3\times t^{-4})$ and therefore the total contribution of zone $\{(x,y)\,:\,|x-y|\ge\beta\}$ is $O(\beta^5)$. Meanwhile a total contribution of the zone $\{(x,y):\,|x-y|\le\beta\}$ is $O(\beta^3\times\beta^2)$. □

Therefore we expect that the zone $\mathcal{X}_1$ defined by $\mu\lesssim 1$ could be expanded to the zone $\mathcal{X}'_1$ defined by $\mu\lesssim h^{-\frac{1}{3}}$ [10)] or even larger[12)]; furthermore, under assumption $|\nabla W|\asymp\zeta^2\ell^{-1}$ we can define $\mathcal{X}'_1$ by $\mu\lesssim h^{-\frac{3}{5}}$ or even larger[12)].

[11)] In fact one can prove then estimates $O(\beta^s)$ but adding correction terms $\sum\kappa_k\beta^{2k}$. However this improvement is not carried on to (2.2.15) in full.

[12)] We do not need for each ℓ have a sharp remainder estimates but need only them to sum to a sharp estimate.

2.2.3 Expansion: justification

Now however we need to deal with $e(x,x,\nu)$ rather than $P'_B(W(x)+\nu)$ (etc).

Proposition 2.2.4. *Assume that in* $B(z,\gamma)$ *conditions* (2.2.10), $\zeta\gamma\ge 1$ *and*

$$B\le c\zeta^2(\zeta\gamma)^{-\delta} \tag{2.2.17}$$

are fulfilled. Then for γ*-admissible* ϕ

$$\int \phi(x)\Bigl(e(x,x,\nu)-\tilde{P}'_B(W(x)+\nu)\Bigr)\,dx=O(\zeta^2\gamma^2), \tag{2.2.18}$$

$$\int \phi(x)\Bigl(e_1(x,x,\tau)-\tilde{P}_B(W(x)+\nu)\Bigr)\,dx=O(\zeta^3\gamma) \tag{2.2.19}$$

and

$$\begin{multline}\mathsf{D}\Bigl(\phi(x)\bigl(e(x,x,\nu)-\tilde{P}'_B(W(x)+\nu)\bigr),\\ \phi(x)\bigl(e(x,x,\nu)-\tilde{P}'_B(W(x)+\nu)\bigr)\Bigr)=O(\zeta^4\gamma^3).\end{multline} \tag{2.2.20}$$

Proof. Estimates (2.2.18) and (2.2.19) are due to Chapter 13 of [Ivr11]. Really, rescale $x\mapsto x\gamma^{-1}$, $\tau\mapsto\tau\zeta^{-2}$ and $h=1\mapsto\hbar=\gamma^{-1}\zeta^{-1}$, $B\mapsto\mu=B\gamma\zeta^{-1}$.

To prove (2.2.20) let us apply decomposition (2.1.8); then according to (2.2.18) $J(t)$ (which is expression (2.2.18) with $t\gamma$-admissible ϕ_t) does not exceed $C\zeta^2\gamma^2t^2$ as long as $t\zeta\gamma\ge 1$; therefore contribution of zone $\{(x,y):\,|x-y|\asymp t\}$ into the left-hand expression of (2.2.20) does not exceed $C(\zeta^2\gamma^2t^2)^2\times t^{-4}\gamma^{-1}\asymp C\zeta^4\gamma^3$.

Then summation over $t\ge\mu^{-1}=B^{-1}\gamma^{-1}\zeta$ returns $C\zeta^4\gamma^3\int t^{-1}\,dt\asymp C\zeta^4\gamma^3\log\mu$ (we assume that $\mu\ge 2$; case $\mu\lesssim 2$ has covered). So, the total contribution of zone $\{(x,y):\,|x-y|\ge\mu^{-1}\}$ does not exceed $C\zeta^4\gamma^3\log\mu$.

Let us get rid off logarithmic factor. Returning back to $B(z,t)$ stretched to $B(0,1)$ one can see easily that conditions of proposition 13.6.25 of [Ivr11] are fulfilled as well with $T=\min\bigl(t^{-\delta},\hbar^{-\delta}t^{\delta}\bigr)$ and thus $|J(t)|\le C(\hbar t^{-1})^{-2}T^{-1}\le C\hbar^{-2}t^2\bigl(t^{\delta}+\hbar^{\delta}t^{-\delta}\bigr)$. Plugging into (2.1.8) we get

$$C\hbar^{-4}\gamma^{-1}\int_{\mu^{-1}}^{1}t^{-1}\bigl(t^{\delta}+h^{\delta}t^{-\delta}\bigr)\,dt\asymp C\hbar^{-4}\gamma^{-1}=C\zeta^4\gamma^3.$$

On the other hand, in zone $t \le \mu^{-1}$ we use the trivial estimate

$$e(x,x,\nu) - P'(W(x)+\nu) = O(\mu\zeta^2\gamma^2)$$

(due to simple rescaling $x \mapsto \mu x$) and its contribution to the left-hand expression of (2.2.20) does not exceed $C(\mu\zeta^2\gamma^2)^2 \times \mu^{-2}\gamma^{-1} \asymp C\zeta^4\gamma^3$. $\square$

Combining with estimates (2.2.8) and (2.2.9) we arrive to the statement (i) below; combining with Proposition 2.1.7 to the statement (ii):

Corollary 2.2.5. *Assume that in $B(z,\gamma)$ conditions* (2.2.10) *and $\zeta\gamma \ge 1$ are fulfilled. Let ϕ be γ-admissible function.*

(i) Let

$$B \le c\zeta^{\frac{4}{3}}\gamma^{-\frac{2}{3}}; \tag{2.2.21}$$

then

$$\int \phi(x)\Bigl(e(x,x,\nu) - \tilde{P}'_B(W(x)+\nu)\Bigr)\,dx = O(\zeta^2\gamma^2), \tag{2.2.22}$$

$$\int \phi(x)\Bigl(e_1(x,x,\tau) - \tilde{P}_B(W(x)+\nu)\Bigr)\,dx = O(\zeta^3\gamma) \tag{2.2.23}$$

and

$$\begin{multline} \mathsf{D}\Bigl(\phi(x)\bigl(e(x,x,\nu) - \tilde{P}'_B(W(x)+\nu)\bigr),\\ \phi(x)\bigl(e(x,x,\nu) - \tilde{P}'_B(W(x)+\nu)\bigr)\Bigr) = O(\zeta^4\gamma^3); \end{multline} \tag{2.2.24}$$

(ii) Let assumption (2.2.11) *be fulfilled and*

$$B \le c\zeta^{\frac{8}{5}}\gamma^{-\frac{2}{5}}. \tag{2.2.25}$$

Then (2.2.22)–(2.2.24) *hold.*

2.3 Rough approximation

Unless our analysis has been cut short with $\bar{r}_1 \gtrsim (Z-N)_+^{-\frac{1}{3}}$, we need to consider zone $\{x:\, \ell(x) \ge \bar{r}_1\}$ with redefined $\bar{r}_1$, so that this zone is described by $\mu \gtrsim \hbar^{-\frac{1}{3}}$ or $\mu \gtrsim \hbar^{-\frac{3}{5}}$ in the general or non-degenerate (i.e. satisfying assumption (2.2.11)) cases respectively.

In this zone we consider $\varepsilon\ell$-mollification with $\varepsilon \ll 1$. In contrast to potentials considered in Chapter 18 of [Ivr11] function W^{TF}_B is more regular.

2.3.1 Properties of mollification

First, recall regularity properties of W_B^{TF}:

Proposition 2.3.1. *W_B^{TF} have the following properties:*

$$|\nabla^\alpha W_B^{\mathsf{TF}}(x)| \le c_\alpha \zeta(x)^2 \ell(x)^{-|\alpha|} \qquad \forall \alpha : |\alpha| \le 2, \tag{2.3.1}$$

$$\begin{aligned}|\nabla^\alpha \bigl(W_B^{\mathsf{TF}}(x) - W_B^{\mathsf{TF}}(y)\bigr)| \le \\ c_0 B\ell(x)^{-\frac{5}{2}}|x-y|^{\frac{1}{2}} + c_0\zeta(x)^2\ell(x)^{-3}|x-y| \\ \forall |\alpha| = 2 \quad \forall x, y : |x-y| \le \epsilon\ell(x)\end{aligned} \tag{2.3.2}$$

where we recall

$$\zeta(x) = \min\bigl(Z^{\frac{1}{2}}\ell(x)^{-\frac{1}{2}}, \ell(x)^{-2}\bigr) \qquad \textit{as}\ \ B \le Z^{\frac{4}{3}}, \tag{2.3.3}$$

$$\zeta(x) = Z^{\frac{1}{2}}\ell(x)^{-\frac{1}{2}} \qquad \textit{as}\ \ B \ge Z^{\frac{4}{3}}; \tag{2.3.4}$$

Proof. This proof is rather obvious corollary of the Thomas-Fermi equation (1.1.3). See also arguments below. ☐

Let us consider $B(z, \ell(z))$ with $\zeta^2 \gtrsim B$ and rescale $x \mapsto x\ell^{-1}$, $W \mapsto w = \zeta^{-2}(W + \nu)$ (where we included ν for a convenience). After such rescaling $w \in \mathscr{C}^{\frac{5}{2}}$ uniformly, but there is more: Thomas-Fermi equation (1.1.3) translates into

$$\Delta w = \ell^2 P'_\beta(w) = \ell^2\zeta P'(w) + \ell^2\zeta\bigl(P'_\beta(w) - P'(w)\bigr) \tag{2.3.5}$$

with $\beta = B\zeta^{-2}$; we used that $P'_B(W)$ is positively homogeneous of degree 3 with respect to (W, B).

Note that parameter $\eta := \zeta\ell^2 \lesssim 1$ and $\eta \asymp 1$ if and only if $B \lesssim Z^{\frac{4}{3}}$ and $\ell \gtrsim Z^{-\frac{4}{3}}$ (in which case $\zeta \asymp \ell^{-2}$).

Also note that the first term and the second terms in the right-hand expression of (2.3.5) belong to $\mathscr{C}^{\frac{5}{2}}$ and $\beta\eta\mathscr{C}^{\frac{1}{2}}$ respectively with the uniform classes and

$$\beta\eta = \beta\zeta\ell^2 = B\zeta^{-1}\ell^2, \qquad \eta := \zeta\ell^2 \qquad \text{as}\ \ \beta \lesssim 1 \tag{2.3.6}$$

Because of this $w \in \mathscr{C}^{\frac{9}{2}} \oplus \beta\eta\mathscr{C}^{\frac{5}{2}}$ again with uniform classes. Iterating we conclude that $w \in \mathscr{C}^K \oplus \beta\eta\mathscr{C}^{\frac{5}{2}}$ with arbitrarily large exponent K.

On the other hand, if $B \gtrsim \zeta^2$ (i.e. $\beta \gtrsim 1$) without invoking P'_B one can prove easily that $w \in \eta\mathscr{C}^{\frac{5}{2}}$ with

$$\eta := \beta\zeta\ell^2 = B\zeta^{-1}\ell^2 \qquad \text{as } \ \beta \gtrsim 1 \tag{2.3.6$'$}$$

Therefore

(2.3.7) $w \in \mathscr{C}^K \oplus \beta\eta\mathscr{C}^{\frac{5}{2}}$ with arbitrarily large exponent K as $\beta \lesssim 1$ and $w \in \eta\mathscr{C}^{\frac{5}{2}}$ as $\beta \gtrsim 1$

and one can see easily that

(2.3.8) Parameter $\eta = B\zeta^{-1}\ell^2$ is $O(1)$ and $\eta \asymp 1$ iff either $B \le Z^{\frac{4}{3}}$ and $\ell \asymp B^{-\frac{1}{4}}$ or $B \ge Z^{\frac{4}{3}}$ and $\ell \asymp B^{-\frac{2}{5}}Z^{\frac{1}{5}}$ (i.e. near border of $\operatorname{supp}\rho_B^{\mathsf{TF}}$, uncut by ν).

Remark 2.3.2. It may seem strange to define η differently as $\beta \lesssim 1$ and $\beta \gtrsim 1$ but there is a good reason for this as we consider $M \ge 2$. Anyway, η is the magnitude of the right-hand expression of (2.3.5).

Proposition 2.3.3. *(i) Let w_ε be a ε-mollification of w with $\varepsilon \lesssim \min(\beta, \hbar^\delta)$. Then as $\beta \lesssim 1$*

$$|\nabla^\alpha(w - w_\varepsilon)| \le c_\alpha \beta\eta\varepsilon^{\frac{5}{2}-|\alpha|} \qquad \forall\alpha : |\alpha| \le 2, \tag{2.3.9}$$

$$|P_\beta(w) - P_\beta(w_\varepsilon)| \le c\beta\eta\varepsilon^{\frac{5}{2}} \tag{2.3.10}$$

and

$$|P'_\beta(w) - P'_\beta(w_\varepsilon)| \le c\beta^{\frac{3}{2}}\eta^{\frac{1}{2}}\varepsilon^{\frac{5}{4}} + c\beta\eta\varepsilon^{\frac{5}{2}}; \tag{2.3.11}$$

(ii) On the other hand as $\beta \gtrsim 1$ the right-hand expressions of (2.3.9)–(2.3.11) *should be replaced by similar expressions albeit without β:*

$$|\nabla^\alpha(w - w_\varepsilon)| \le c_\alpha \eta\varepsilon^{\frac{5}{2}-|\alpha|} \qquad \forall\alpha : |\alpha| \le 2, \tag{2.3.9$'$}$$

$$|P_\beta(w) - P_\beta(w_\varepsilon)| \le c\eta\varepsilon^{\frac{5}{2}} \qquad \forall\alpha : |\alpha| \le 2, \tag{2.3.10$'$}$$

and

$$|P'_\beta(w) - P'_\beta(w_\varepsilon)| \le c\eta^{\frac{1}{2}}\varepsilon^{\frac{5}{4}}; \qquad \forall\alpha : |\alpha| \le 2. \tag{2.3.11$'$}$$

(iii) Further, under assumption $|\nabla w|\asymp 1$ *in both cases*

$$|\int \phi(x)\bigl(P_\beta(w)-P_\beta(w_\varepsilon)\bigr)\,dx|\le c\eta\varepsilon^{\frac{7}{2}}, \tag{2.3.12}$$

$$|\int \phi(x)\bigl(P'_\beta(w)-P'_\beta(w_\varepsilon)\bigr)\,dx|\le c\eta\varepsilon^{\frac{9}{4}} \tag{2.3.13}$$

and

$$\mathsf{D}\bigl(\phi(P'_\beta(w)-P'_\beta(w_\varepsilon)),\,\phi(P'_\beta(w)-P'_\beta(w_\varepsilon))\bigr)\le c\eta^2\varepsilon^{\frac{9}{2}}. \tag{2.3.14}$$

Proof. Proof of (i) is trivial; in particular we note that $\eta\varepsilon^{\frac{5}{2}}\lesssim\beta$.

Proof of (iii) is also easy as then w_ε is different from w on the set of measure $\asymp\beta^{-1}\varepsilon$ as $\beta\le C_0$ and on the set of measure $\asymp\varepsilon$ as $\beta\ge C_0$. Actually w is uniformly smooth as $\beta\gtrsim 1$ and $\ell(x)\le\epsilon\bar{r}$ and we do not need any mollification here.

One definitely can improve estimates (2.3.12)–(2.3.14) but we do not need it. □

Consider now the analytical expression and estimate the semiclassical error. From now on until the end of the subsection we assume that $M=1$ to avoid possible degenerations.

Remark 2.3.4. Note that we can reduce operator to a canonical form as $\varepsilon\ge C(\mu^{-1}\hbar)^{\frac{1}{2}}|\log\mu|$ (see Section 18.7) of [Ivr11] but here we will have a much better estimate as we will take $\varepsilon\ge\hbar^{\frac{2}{3}-\delta}$.

2.3.2 Charge term

Let us consider the *charge term* i.e. expression $\int e(x,x,\nu)\,dx=(\operatorname{Tr}\theta(\nu-H))$.

Regular zone.

Then Section 18.9 of [Ivr11] implies that as

$$W+\nu\asymp\zeta^2 \tag{2.3.15}$$

and

$$|\nabla W|\asymp\zeta^2\ell^{-1} \tag{2.3.16}$$

contribution of the ball $B(x,\ell(x))$ to expression (2.2.2) does not exceed $C(1+\mu\hbar)\hbar^{-2}\asymp C\zeta^2\ell^2+CB\ell^2$ exactly as in the mock proof.

Then summation with respect to ℓ-partition in this zone results in $CB^{\frac{2}{3}}$ as $B\le Z$, $CZ^{\frac{2}{3}}$ as $Z\le B\le Z^{\frac{4}{3}}$ and $CB^{\frac{1}{5}}Z^{\frac{2}{5}}$ as $Z^{\frac{4}{3}}\le B\le Z^3$.

Remark 2.3.5. (i) Condition (2.3.15) is fulfilled as $\ell(x)\le \epsilon\bar{r}$.

(ii) Further, as $M=1$ both conditions (2.3.15) and (2.3.16) are fulfilled as $|x|\le (1-\epsilon)\bar{r}_m$ (we pick up $\mathsf{y}_m=0$).

Border strip.

Now we need to consider the contribution of the border strip $\mathcal{Y}:=\{x:\ \gamma(x)\le\epsilon\}$ with $\gamma(x)=\epsilon(\bar{r}-|x|)\bar{r}^{-1}$ and $\bar{r}:=\bar{r}_m$. Here $\ell\asymp\bar{r}$, $\zeta\asymp\bar{\zeta}$ with

$$\bar{r}\asymp\begin{cases}(Z-N)_+^{-\frac{1}{3}} & B\le (Z-N)_+^{\frac{4}{3}},\\ B^{-\frac{1}{4}} & (Z-N)_+^{\frac{4}{3}}\le B\le Z_+^{\frac{4}{3}},\\ Z^{\frac{1}{5}}B^{-\frac{2}{5}} & Z_+^{\frac{4}{3}}\le B\le CZ^3\end{cases}\tag{2.3.17}$$

and

$$\bar{\zeta}\asymp\begin{cases}(Z-N)_+^{\frac{2}{3}} & B\le (Z-N)_+^{\frac{4}{3}},\\ B^{\frac{1}{2}} & (Z-N)_+^{\frac{4}{3}}\le B\le Z_+^{\frac{4}{3}},\\ Z^{\frac{2}{5}}B^{\frac{1}{5}} & Z_+^{\frac{4}{3}}\le B\le CZ^3\end{cases}\tag{2.3.18}$$

and scaling we get $\mu=B\bar{r}\bar{\zeta}^{-1}$ and $\hbar=\bar{\zeta}^{-1}\bar{r}^{-1}$ here.

Let us consider first the case $\nu=0$. Then both conditions (2.3.15) and (2.3.16) are fulfilled albeit with $\ell_1=\gamma(x)\bar{r}$ and $\varsigma(x)=\bar{\zeta}\gamma(x)^2$ instead of ℓ and ζ.

Thus as $\varsigma\gamma\bar{r}\ge 1$ (i.e. $\gamma\ge\bar{\gamma}:=\hbar^{\frac{1}{3}}$) the contribution of the ball $B(x,\gamma(x)\bar{r})$ to the remainder does not exceed $C\mu\hbar^{-1}\gamma^2$ and therefore the total contribution of zone $\mathcal{Y}_1:=\{x:\ \bar{\gamma}\le\gamma(x)\le\epsilon\bar{r}\}$ to the remainder does not exceed

$$C\mu\hbar^{-1}\int\gamma(x)^{-1}\,dx\asymp C\mu\hbar^{-1}|\log\hbar| = CB\bar{r}^2|\log\hbar|\tag{2.3.19}$$

which is $O(Z^{\frac{2}{3}})$ as long as $B\le Z^{\frac{4}{3}}(\log Z)^{-2}$.

Further, the same approach works as $|\nu|\lesssim\bar{\zeta}^2\bar{\gamma}^3\asymp\bar{\zeta}^2\hbar\asymp\bar{\zeta}\bar{r}^{-1}$ which is equivalent to $(Z-N)_+\le\bar{\zeta}$ (then $|\nabla W|\asymp\varsigma^2\ell_1^{-1}$ as $\gamma(x)\ge\bar{\gamma}$) and also if this condition is violated but $|\nu|\le \bar\zeta^2$; in the latter case we need to pick up $\bar\gamma=\bar\gamma_1:=|\nu|^{\frac{1}{3}}\bar\zeta^{-\frac{2}{3}}$.

To get rid off the logarithmic factor let us consider propagation. Recall that it goes along magnetic lines i.e. as (x_1,x_2) remain constant. Let us consider propagation in the direction in which $|x_3|$ increases (i.e. $\gamma(x)$ decays); we do not need to consider zone $\mathcal{Y}_1\cap\{|x_3|\le Z^{-\delta}\bar r\}$ as contribution of this zone (2.3.19) is $o(B\bar r^2)$.

One can see easily that we can follow dynamics which does not return for a time $T_1^*(x):=T_1(x)(\gamma(x)/\bar\gamma)^\delta$ where $T(x)\asymp \ell_1^{-1}\varsigma^{-1}\asymp \bar r\bar\zeta^{-1}\gamma^{-1}$ is a time required for this dynamics to pass though $B(x,\ell_1(x))$. Therefore one can replace (2.3.19) by

$$C\mu\hbar^{-1}\int_{\{x:\,\gamma(x)\ge\bar\gamma\}}\bar\gamma^\delta\gamma(x)^{-1-\delta}\,dx\asymp C\mu\hbar^{-1}=CB\bar r^2. \tag{2.3.20}$$

Further, as $|\nu|\ge B\bar r$ we need also to consider zone $\mathcal{Y}_0:=\{x:\,\gamma(x)\le\bar\gamma\}$. In this zone we take $\ell_1=\bar\ell_1=\bar\gamma\bar r$ and $\varsigma=\bar\varsigma=(|\nu|\bar\gamma)^{\frac{1}{2}}$ with $\ell_1\varsigma\ge 1$ and since $|\nabla W|\asymp\varsigma^2\ell_1^{-1}$, contribution of $B(x,\ell_1(x))$ to the remainder does not exceed $CB\bar\ell_1^2$ and the total contribution of $\mathcal{Y}_2$ does not exceed $CB\bar r^2$ which what exactly we got for zone $\mathcal{Y}_1$ after we got rid off logarithm. We take mollification parameter $\varepsilon=\bar\varsigma^{-1}Z^{\delta}$[13].

Furthermore, zone $\mathcal{Y}_3=\{x:\,|x|\ge\bar r+\bar\ell_1\}$ is classically forbidden. So we can take here

$$\ell_1(x)=\epsilon(|x|-\bar r),\qquad \varsigma(x)=\min\bigl(\bar\varsigma\bar\ell_1^{-\frac{1}{2}}\ell_1(x)^{\frac{1}{2}},|\nu|^{\frac{1}{2}}\bigr) \tag{2.3.21}$$

and prove easily that its contribution also does not exceed $CB\bar r^2$.

Returning to the case $|\nu|\lesssim\bar\zeta$ we see that the contribution of zone $\mathcal{Y}_2$ to the remainder does not exceed $CB\bar r^2$ because effective semiclassical parameter here is $\hbar_*=1$ and non-degeneracy condition is of no concern for us. We take mollification parameter $\varepsilon=\bar\varsigma^{-1}Z^{\delta}$[13].

Moreover, we can modify W in $\mathcal{Y}_2$ (make it negative there) so that this zone would be classically forbidden with ℓ_1, ς defined by (2.3.21) with $|\nu|$ replaced by $\bar\zeta$.

Finally in the case $B\le|\nu|$ (i. e. $B\le C(Z-N)_+^{\frac{4}{3}}$ we can apply the above arguments with $\bar\gamma=1$ and arrive to the same result. Therefore we proved in all cases

---

[13] One can see easily that the resulting errors in the expressions(2.2.2) and (2.2.3)–(2.2.4) will not violate our claims.

(2.3.22) The total contribution of the border strip $\mathcal{Y}$ to the remainder in is $O(B\bar{r}^2)$ which does not exceed $CB^{\frac{1}{2}}$ as $B \le Z^{\frac{4}{3}}$ and $CB^{\frac{1}{5}}Z^{\frac{2}{5}}$ as $Z^{\frac{4}{3}} \le B \le Z^3$.

Conclusion.

As $Z^2 \le B \le Z^3$ we need to estimate also contribution of the *inner core* $\mathcal{X}_0 := \{x :\ \ell(x) \le CZ^{-1}\}$. By means of variational methods we will prove (see 8.B.2)

(2.3.23) As $Z^2 \le B \le Z^3$ the contributions of $\mathcal{X}_0$ to both $\int e(x,x,\nu)\,dx$ and $\int P'_B(W(x)+\nu)\,dx$ do not exceed CBZ^{-2}.

Then we arrive to the following

Proposition 2.3.6. *Let $M=1$. Then*

(i) For constructed above potential W expression (2.2.2) *does not exceed $CZ^{\frac{2}{3}} + CB^{\frac{1}{5}}Z^{\frac{2}{5}}$;*

(ii) As $B \le Z$ expression (2.2.2) *does not exceed $C(B+1)^{\delta}Z^{\frac{2}{3}-\delta}$.*

2.3.3 Trace term

Let us consider the *trace term* i.e. expression $\int e_1(x,x,\nu)\,dx = \operatorname{Tr}((H-\nu)^-)$.

Regular zone.

Here again let us consider first zone where $|x| \le (1-\epsilon)\bar{r}$. Then the contribution of $B(x,\ell(x))$ to the Tauberian remainder[14] does not exceed $C\zeta^2(\hbar^{-1}+\mu) \asymp C\zeta^3\ell + CB\zeta\ell$ as in the mock proof and the summation over zone results in $CZ^{\frac{5}{3}} + CB^{\frac{1}{3}}Z^{\frac{4}{3}} + CZ^{\frac{3}{5}}B^{\frac{4}{5}}$.

Border strip.

Again in zone $\mathcal{Y}_1$ contribution of $B(x,\gamma(x))$ does not exceed $CB\varsigma\ell_1$ and the summation over this zone returns

$$CB\int \varsigma\ell_1^{-2}\,dx \tag{2.3.24}$$

---

[14] We will consider a bit later transition from the Tauberian to magnetic Weyl expression.

and plugging $\ell_1=\bar{r}\gamma$ and $\varsigma=\bar{\zeta}\gamma^2$ results in $CB^{\frac{5}{4}}$ as $B\lesssim Z^{\frac{4}{3}}$ and $CB^{\frac{4}{5}}Z^{\frac{3}{5}}$ otherwise. The analysis of zone $\mathcal{Y}_0$ if there $\mathcal{Y}_2=\emptyset$ is also easy.

Consider zones $\mathcal{Y}_2$ and $\mathcal{Y}_0$. The same arguments as before imply that their contributions to the remainder do not exceed $CB\bar{r}^2\bar{\varsigma}\bar{\ell}_1^{-1}$ which is what we got before.

Justification: from Tauberian to magnetic Weyl expression.

Case $\mu\hbar\le C_0$. We need to prove that with the announced error we can replace the Tauberian expression by magnetic Weyl one. Note that the canonical form of $\zeta^{-2}H_{A,W}$ as described in Sections 13.2 and 18.7 of [Ivr11] is

$$\begin{aligned}\mathcal{H}=\mathcal{H}_0+&\\ &\mu^{-2}\omega_1(x_1,\mu^{-1}\hbar D_1,x_3)+\mu^{-2}\omega_2(x_1,\mu^{-1}\hbar D_1,x_3)(x_2^2+\mu^2\hbar^2D_2^2)\\ &\quad+\mu^{-1}\hbar\omega_3(x_1,\mu^{-1}\hbar D_1,x_3)+O\bigl(\mu^{-3}\hbar(\gamma+\mu^{-1})^{-\frac{3}{2}}+\mu^{-4}\bigr)\end{aligned}\tag{2.3.25}$$

with

$$\mathcal{H}_0=\hbar^2D_3^2-(x_2^2+\mu^2\hbar^2D_2^2\pm\mu\hbar)+w(x_1,\mu^{-1}\hbar D_1,x_3)\tag{2.3.26}$$

and

$$\gamma=\epsilon\min_j|w-2j\mu\hbar|\tag{2.3.27}$$

where we used the fact that $w\in\mu\hbar\mathscr{C}^{\frac{5}{2}}+\mathscr{C}^K$, $\mu^{-3}\hbar=\mu^{-4}\cdot\mu\hbar$. Here we have signs "+" and "−" on $q/2$ of the diagonal elements equally.

Then the Tauberian expression is

$$\begin{aligned}\mathsf{const}\cdot\mu\hbar^{-2}\int\sum_{j\ge0}\bigl(w-2j\mu\hbar-\mu^{-2}\omega_1-2j\mu^{-1}\hbar\omega_2\bigr)_+^{\frac{3}{2}}\times&\\ \bigl(\psi+\mu^{-2}\psi_1+2j\mu^{-1}\hbar\psi_2\bigr)\,dx&\end{aligned}\tag{2.3.28}$$

where term with $j=0$ enters with the weight $\frac{1}{2}$ and an error does not exceed

$$C\hbar^{-3}\Bigl(\mu^{-4}+\mu^{-3}\hbar\int(\gamma+\mu^{-1})^{-\frac{3}{2}}\,dx\Bigr)\asymp C\mu^{-\frac{7}{2}}\hbar^{-3}$$

because an integral does not exceed $C\mu^{\frac{1}{2}}(\mu\hbar)^{-1}$; since $\mu\ge\hbar^{-\frac{3}{5}}$ this error does not exceed $C\hbar^{-\frac{9}{10}}$ which is better than $O(\hbar^{-1})$.

On the other hand, if we consider the difference between (2.3.28) and the same expression with $\omega_1=\omega_2=\psi_1=\psi_2=0$ and consider it as a Riemannian sum and replace it by an integral we get $G\mu^{-2}\hbar^{-3}$ with an error not exceeding $C\mu^{-4}(\mu\hbar)^{\frac{1}{2}}\hbar^{-3}$ which is even less. Therefore (2.3.28) becomes

$$\int P_{\mu\hbar}(w)\psi\, dx + G\mu^{-2}\hbar^{-3}$$

and comparing with the result as $\mu\asymp\hbar^{-\frac{3}{5}}$ when we get the same answer albeit with $G=0$ we conclude that G *must* be 0. This concludes the justification in $\mathcal{X}_2$.

Case $\mu\hbar\ge C_0$. In this case we need a simplified version of (2.3.25) $\mathcal{H}=\mathcal{H}_0+O(\mu^{-1}\hbar)$ and we need to consider only $j=0$ and replacing $\mathcal{H}$ by $\mathcal{H}_0$ brings and error $C\mu\hbar^{-2}\times\mu^{-1}\hbar=O(\hbar^{-1})$. This takes care of $\mathcal{X}_2$ and after scaling of $\mathcal{Y}$.

Conclusion.

As $Z^2\le B\le Z^3$ we need to estimate also contribution of $\mathcal{X}_0=\{x:\,\ell(x)\le CZ^{-1}\}$. By means of variational methods we will prove (see 8.B.2)

(2.3.29) As $Z^2\le B\le Z^3$ the contributions of $\mathcal{X}_0$ to both $\int e_1(x,x,\nu)\,dx$ and $\int P'_B(W(x)+\nu)\,dx$ do not exceed CB.

Then we arrive to the following

Proposition 2.3.7. *Let $M=1$. Then*

(i) For constructed above potential W expression (2.2.4) *does not exceed $CZ^{\frac{5}{3}}+CB^{\frac{1}{3}}Z^{\frac{4}{3}}+CZ^{\frac{3}{5}}B^{\frac{4}{5}}$;*

(ii) As $B\le Z$ expression (2.2.4) *does not exceed $C(B+1)^{\delta}Z^{\frac{5}{3}-\delta}$ (but one should subtract a Schwinger term from the trace).*

2.3.4 Semiclassical D-term: local theory

Unfortunately, we do not have non-smooth theory (cf. Section 16.7) of [Ivr11] here but actually we almost do not need it as singularities are rather rare. Let us introduce a scaling function (2.3.27) and consider

$$J_\lambda(z)=\int \phi_{z,\lambda}(x)\bigl(e(x,x,\tau)-P_\beta(w(x)+\tau)\bigr)\,dx \tag{2.3.30}$$

with $\phi_{z,\lambda}(x)=\phi(\lambda^{-1}(x-z))$ and $\lambda\le\gamma(z)$. Scaling $x\mapsto\lambda^{-1}(x-z)$ we have $\mu\mapsto\mu'=\lambda\mu$ and $\hbar\mapsto\hbar'=\lambda^{-1}\hbar$.

Then, according to Section 13.4 of [Ivr11]

$$|J_\lambda(z)|\le C\hbar'^{-2}(1+\mu'\hbar')\asymp C\lambda^2\hbar^{-2}(1+\mu\hbar) \tag{2.3.31}$$

as long as $\lambda\ge\hbar$.

Really, transition from the Tauberian decomposition to magnetic Weyl one in this case is easy: skipping all perturbation terms $O(\mu^{-2})$ in (2.3.25) and also setting $\psi_1=\psi_2=0$ results in error $O\bigl(\mu^{-1}\cdot\mu\hbar^{-3}+\mu^{-2}\hbar^{-3}\bigr)$ in (2.3.28)-like expression albeit with the power $\frac{1}{2}$ rather than $\frac{3}{2}$ and without integration:

$$\mathsf{const}\cdot\mu\hbar^{-2}\sum_{j\ge 0}\bigl(w-2j\mu\hbar-\mu^{-2}\omega_1-2j\mu^{-1}\hbar\omega_2\bigr)_+^{\frac{1}{2}}\times \\ \bigl(\psi+\mu^{-2}\psi_1+2j\mu^{-1}\hbar\psi_2\bigr) \tag{2.3.32}$$

Let us apply this estimate (2.3.31) to the Fefferman–de Llave decomposition.

Case $\mu\le C_0\hbar^{-1}$.

(i) Consider first a pair (z,z') such that $|z'-z''|\le\epsilon_0\gamma(z')$; then also $|z'-z''|\le\epsilon_0\gamma(z'')$ and we take $\lambda=\epsilon|z'-z''|$.

Then in the virtue of (2.3.31) the total contribution to D-term of all such pairs belonging to $B(z,\gamma(z))$, and with $|z'-z''|\asymp\lambda$ does not exceed

$$C\gamma^3\lambda^{-3}\times\lambda^{-1}\times\lambda^2\hbar^{-4}(1+\mu\hbar)^2\asymp C\gamma^3\hbar^{-4}(1+\mu\hbar)^2 \tag{2.3.33}$$

where $C\gamma^3\lambda^{-3}$ estimate the number of such pairs, λ^{-1} the inverse distance between them, and $C\lambda^2\hbar^{-2}(1+\mu\hbar)$ is the right-hand expression of (2.3.31).

Then summation over $\lambda\in(\mu^{-1},\gamma)$ results in $C\gamma^3(1+\mu\hbar)^2\hbar^{-4}|\log(\mu\gamma)|$.

Further, summation over all balls $B(z,\gamma)\subset B(0,1)$ with $\gamma(z)\asymp\gamma$ results in $C(\mu\hbar)^{-1}\gamma\hbar^{-4}|\log(\mu\gamma)|$ as there are $\asymp(\mu\hbar)^{-1}\gamma^{-2}$ such balls due to non-degeneracy assumption $|\nabla w|\asymp 1$. Summation over $\gamma\in(\mu^{-1},\mu\hbar)$ results in $C\hbar^{-4}|\log(\mu^2\hbar)|$.

As $\lambda\le\mu^{-1}$ we can apply standard non-magnetic methods without Fefferman–de Llave decomposition. Coefficients are smooth after scaling as long as $\varepsilon\ge\mu^{-1}$.

(ii) Consider *disjoint* pairs (z',z'') with $|z'-z''|\ge\max\bigl(\gamma(z'),\gamma(z'')\bigr)$. Here estimate (2.3.31) is not sufficient and it should be replaced by

$$|J_\gamma(z)|\le C\lambda^3\hbar^{-2}(1+\mu\hbar) \tag{2.3.34}$$

as

$$\gamma\ge\hbar^{\frac{2}{3}-\delta}. \tag{2.3.35}$$

Really, the shift for time T with respect to ξ_3 is $\asymp T$ provided $|\nabla_{x_3}w|\asymp 1$ and this shift is observable if $T\times\gamma\gtrsim\hbar^{1-\delta}$. Similarly, in the canonical form the shift for time T with respect to $\mu^{-1}\xi_i$ is $\asymp\mu^{-1}T$ provided $|\nabla_{x_i}w|\asymp 1$ and this shift is observable if $\mu^{-1}T\times\gamma\gtrsim\mu^{-1}\hbar^{1-\delta}$. In both cases shift with $T\in(\gamma^{\frac{1}{2}},\epsilon_0)$ is observable under assumption (2.3.35) and therefore we can extend $T\asymp\gamma^{\frac{1}{2}}$ to $T\asymp 1$.

Note that for $\varepsilon\ge\hbar^{\frac{2}{3}-\delta}$ assumption (2.3.35) is fulfilled automatically. Then contribution of each disjoint pair to D-term does not exceed

$$C\hbar^{-4}(1+\mu\hbar)^2\gamma(z')^3\gamma(z'')^3|z'-z''|^{-1}$$

and the total contribution does not exceed

$$C\hbar^{-4}(1+\mu\hbar)^2\iint|z'-z''|^{-1}\,dz'dz''\asymp C\hbar^{-4}(1+\mu\hbar)^2.$$

(iii) To shed off logarithm in (i) we need a slightly better estimate than (2.3.31). The same arguments as in (ii) result in

$$|J_\lambda(z)|\le C\lambda^2\hbar^{-2}(1+\mu\hbar)\cdot(1+\lambda\gamma/\hbar)^{-\delta}. \tag{2.3.36}$$

Really, we just advance from time $T\asymp\lambda$ to $T\asymp\lambda(1+\lambda\gamma/\hbar)^{\delta}$.

Then the same factor is acquired by the right-hand expression of (2.3.33) and the summation with respect to $\lambda \in (\hbar\gamma^{-1},\gamma)$ results in $C\gamma^3(1+\mu\hbar)^2\hbar^{-4}$ but summation with respect to $\lambda \in (\mu^{-1},\hbar\gamma^{-1})$ results

$$C\gamma^3(1+\mu\hbar)^2\hbar^{-4}\bigl(1+|\log(\mu\hbar\gamma^{-1})|\bigr).$$

Further, summation over all balls $B(z,\gamma)\subset B(0,1)$ with $\gamma(z)\asymp\gamma$ results in $C(\mu\hbar)^{-1}\gamma(1+\mu\hbar)^2\hbar^{-4}\bigl(1+|\log(\mu\hbar\gamma^{-1})|\bigr)$ and, finally, summation over $\gamma\lesssim\mu\hbar$ results in $C\hbar^{-4}(1+\mu\hbar)^2$.

Note that in all cases perturbation terms in (2.3.25) and (2.3.32) result in the error not exceeding the announced one.

Case $\mu \ge C_0\hbar^{-1}$. So far factor $(1+\mu\hbar)$ was for compatibility only. Now it is important.

Exactly the same arguments work as $\mu\ge C_0\hbar^{-1}$ with a minor modifications:

(a) $\gamma(x)$ now is defined by (2.3.27) with $j=0$ and its upper bound is 1 rather than $\mu\hbar$;

(b) Also the number of γ-balls is $\asymp\gamma^{-2}$ rather than $\asymp\gamma^{-2}$;

(c) λ now runs from $\hbar$ to γ in (i), (iii).

(d) We need to estimate contribution of pairs (z',z'') with $|z'-z''|\le\hbar$. One can see easily that $e(x,x,\tau)\le\mu\hbar^{-2}$ and therefore the total contribution of these pairs does not exceed $C\mu^2\hbar^{-4}\iint|z'-z''|^{-1}\,dz'dz''\asymp C\mu^2\hbar^{-4}\times\hbar^2\asymp C\mu^2\hbar^{-2}$.

Therefore we have the following

Proposition 2.3.8. *As* $|\nabla w|\asymp 1$ *and* $\varepsilon\ge\hbar^{\frac{2}{3}-\delta}$ *in* $B(0,1)$ *and* $\phi\in\mathscr{C}^\infty(B(0,\frac{1}{2})$

$$\mathsf{D}\Bigl(\phi\bigl(e(x,x,\tau)-P'_\beta(w(x)+\tau)\bigr),\phi\bigl(e(x,x,\nu)-P'_\beta(w(x)+\tau)\bigr)\Bigr)\le C(1+\mu\hbar)^2\hbar^{-4}. \tag{2.3.37}$$

Remark 2.3.9. One can see easily that one can select $\varepsilon\ge\hbar^{\frac{2}{3}-\delta}$ such that expressions (2.3.12), (2.3.13) and (2.3.13) will be respectively $O(\hbar^{2+\delta})$, $O(\hbar^{1+\delta})$ and $O(\hbar^{2+\delta})$.

2.3.5 Semiclassical D-term: global theory

Regular zone.

The above results allow us to consider a total contribution of zone $\mathcal{X}_2$ into semiclassical D-term. As before let us consider ℓ-admissible partition of unity there and apply it to Fefferman–de Llave decomposition. Then the total contribution of the elements which are not disjoint does not exceed

$$\sum_n \ell_n^{-1}\bigl(1+B\zeta_n^{-2}\bigr)^2\ell_n^4\zeta_n^4 \asymp \int \underbrace{\bigl(\zeta^4+B^2\bigr)\ell^3}\ \ell^{-1}d\ell \tag{2.3.38}$$

where $\bigl(1+B\zeta_n^{-2}\bigr)^2$ and $\ell_n^4\zeta_n^4$ are $(1+\mu\hbar)$ and $\hbar^{-4}$ respectively and ℓ_n^{-1} is a scaling factor.

Then as $\zeta^2=Z\ell^{-1}$ an integral equals to the value of the selected expression as ℓ reaches its maximum, i.e. as $\ell=Z^{-\frac{1}{3}}$ for $B\le Z^{\frac{4}{3}}$ and $\ell=Z^{\frac{1}{5}}B^{-\frac{2}{5}}$ for $Z^{\frac{4}{3}}\le B\le Z^3$ and we arrive to $CZ^{\frac{5}{3}}$ and $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$ respectively.

On the other hand, as $\zeta^2=\ell^{-4}$ an integral equals to the value of the selected expression as ℓ reaches its minimum, i.e. $\ell=Z^{-\frac{1}{3}}$ and only in the case $B\le Z^{\frac{4}{3}}$ and we arrive to $CZ^{\frac{5}{3}}$ again.

Furthermore, the total contribution of the disjoint elements does not exceed

$$\sum_{n,p}|z_n-z_p|^{-1}\bigl(1+B\zeta_n^{-2}\bigr)\bigl(1+B\zeta_p^{-2}\bigr)\ell_n^2\zeta_n^2\ell_p^2\zeta_p^2 \asymp \\ \iint \underbrace{(\ell+\ell')^{-1}\bigl(\zeta^2+B\bigr)\ell^2\bigl(\zeta'^2+B\bigr)\ell'^2}\ \ell^{-1}d\ell\,\ell'^{-1}d\ell'. \tag{2.3.39}$$

Then as $\zeta^2=Z\ell^{-1}$ and $\zeta'^2=Z\ell'^{-1}$ an integral equals to the value of the selected expression as both ℓ and ℓ' reach their maxima, and we arrive to $CZ^{\frac{5}{3}}$ and $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$ respectively.

On the other hand, as $\zeta^2=\ell^{-4}$ and $\zeta'^2=\ell'^{-4}$ (we do not need to consider mixed pair) an integral equals to the value of the selected expression as both ℓ and ℓ' reach their minima, and we arrive to $CZ^{\frac{5}{3}}$.

Therefore we conclude (combining with proposition 8.B.2 as $Z^2\le B\le Z^3$)

Proposition 2.3.10. *Let* $M=1$*. Then* *(i) The total contribution of the zone* $\{x:\, \ell(x)\le (1-\epsilon)\bar{r}\}$ *to the semiclassical* D*-term does not exceed* $CZ^{\frac{5}{3}}$ *and* $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$ *as* $B\le Z^{\frac{4}{3}}$ *and* $Z^{\frac{4}{3}}\le B\le Z^3$ *respectively;*

(ii) As $B\le Z$ *this contribution does not exceed* $C(B+1)^{\delta}Z^{\frac{5}{3}-\delta}$*.*

Border strip.

Border strip $\mathcal{Y}=\{x:\, (1-\epsilon)\bar{r}\le \ell(x)\le (1+\epsilon)\bar{r}\}$ is more subtle. Here we need to use the same $\ell_1(x)=\epsilon_0(\bar{r}-|x|)$ partition as before.

Remark 2.3.11. $\mathcal{Y}$ is already covered by our arguments if $\bar{r}\asymp (Z-N)_+^{-\frac{1}{3}}$.

Close elements. Consider first contribution of elements which are not disjoint. It is given by the left-hand expression of (2.3.38) with ℓ, ζ replaced by $\ell_1(x)=\bar{r}\gamma(x)$ and $\varsigma(x)=\bar{\zeta}\gamma(x)^2$ respectively. However since the layer $\{x:\, \gamma(x)\asymp\gamma\}$ contains $\asymp\gamma^{-2}$ elements the right-hand expression should be replaced by

$$\int \underbracket{B^2\bar{r}^3\gamma}\,\gamma^{-1}d\gamma \asymp B^2\bar{r}^3$$

since $\varsigma^2\le B$; so we arrive to $O\bigl(\max(B^{\frac{5}{4}}, Z^{\frac{3}{5}}B^{\frac{4}{5}}\bigr)$.

Meanwhile for $\mathcal{Y}_2$ we have $\gamma(x)=\bar{\gamma}\le 1$ and $\varsigma(x)=\bar{\varsigma}=\bar{\zeta}\bar{\gamma}^2$ with and its contribution does not exceed what we got for $\mathcal{Y}_1$.

Disjoint elements. Consider contribution of the disjoint elements. It is given by the left-hand expression of (2.3.39) with ℓ, ζ replaced by γ and ς respectively. Note that $\sum_{n,p}|z_n-z_p|^{-1}\asymp \bar{r}^{-1}\gamma^{-2}\gamma'^{-2}$ where we sum with respect to all pairs with $\gamma_n\asymp\gamma$ and $\gamma_p\asymp\gamma'$. Therefore the right-hand expression should be replaced by

$$\int \underbracket{\bar{r}^3B^2}\,\gamma^{-1}d\gamma\,\gamma'^{-1}d\gamma' \tag{2.3.40}$$

which leads to $C\bar{r}^3B^2|\log(\bar{r}^{-1}\bar{\gamma})|^2$ which differs from what we got before by a logarithmic factor. To get rid off it we will use exactly the same trick as in paragraph "Border strip" proving proposition 2.3.6 because considering disjoint pairs we consider the same objects as there. Then instead of (2.3.40) we arrive to

$$\int \underbracket{\bar{r}^3B^2\gamma^{-\delta}\gamma'^{-\delta}\bar{\gamma}^{2\delta}}\,\gamma^{-1}d\gamma\,\gamma'^{-1}d\gamma'$$

which results in $C\bar{r}^3B^2$.

Meanwhile for $\mathcal{Y}_2$ we have $\gamma(x)=\bar{\gamma}\le\bar{r}$ and $\varsigma(x)=\bar{\varsigma}=\bar{\zeta}\bar{\gamma}^2$ with and its contribution does not exceed what we got for $\mathcal{Y}_1$.

Conclusion. Finally, analysis in the outer zone is trivial. Therefore we arrive

Proposition 2.3.12. *Let $M=1$. Then for constructed above potential W*

(i) Expression (2.2.3) *does not exceed $CZ^{\frac{5}{3}}+CZ^{\frac{3}{5}}B^{\frac{4}{5}}$;*

(ii) As $B\le Z$ expression (2.2.3) *does not exceed $C(B+1)^{\delta}Z^{\frac{5}{3}-\delta}$.*

3 Applying semiclassical methods: $M\ge 2$

Let us consider now the molecular case ($M\ge 2$). The major problem is that non-degeneracy condition $|\nabla w|\asymp 1$ is not necessarily fulfilled. Therefore we need to find an alternative approach to the zone where $\mu\ge h^{-\frac{1}{3}}$. Recall that it consists of three zones: zone $\mathcal{X}_2:=\{\mu h\le C_0,\ W_B^{\mathsf{TF}}\ge\epsilon_0\zeta^2\}$[15], zone $\mathcal{X}_3:=\{\mu h\ge C_0,\ W_B^{\mathsf{TF}}\ge\epsilon_0\zeta^2\}$[16], and the (most difficult) *boundary strip* $\mathcal{Y}=\{W_B^{\mathsf{TF}}\le\epsilon_0\zeta^2\}$.

3.1 Analysis in zone $\mathcal{X}_2$

3.1.1 Scaling functions in zone $\mathcal{X}_2$

Step 1.

We will use the scaling method here; the good news is that W_B^{TF} is sufficiently regular for a proper rescaling. Recall that after we rescale in $x\mapsto\bar{\ell}^{-1}(x-\bar{x})$, $\tau\mapsto\bar{\zeta}^{-2}\tau$ in $B(\bar{x},\frac{1}{2}\bar{\ell})$ with $\bar{\ell}=\ell(\bar{x})$, $\bar{\zeta}=Z^{\frac{1}{2}}\bar{\ell}^{-1}$, the rescaled potential $w=\bar{\zeta}^{-2}W_B^{\mathsf{TF}}$ satisfies in $B(0,2)$ equation

$$\Delta w=\eta P'_{\beta}(w)\qquad\text{with }\ \eta=\bar{\zeta}\bar{\ell}^2\le 1,\quad \beta=\mu\hbar=B\bar{\zeta}^{-2}\le 1 \tag{3.1.1}$$

---

[15] Only as $B\le C_1Z^2$; this zone disappears for $C_1Z^2\le B\lesssim Z^3$.

[16] Only as $Z^{\frac{4}{3}}\lesssim B\lesssim Z^3$; this zone disappears for $B\lesssim Z^{\frac{4}{3}}$.

and therefore in $B(0,1)$

$$w=-\frac{1}{4\pi}\int |x-z|^{-1}\eta P'_\beta(w(z))\phi(z)\,dz+w' \tag{3.1.2}$$

with $\phi\in\mathscr{C}_0^\infty B(0,\frac{5}{6})$ and $w'\in\mathscr{C}_0^\infty B(0,\frac{3}{4})$.

Let us introduce a function

$$\gamma_0(x)=\Bigl(\min_j |w-2j\beta|^3+|\nabla w|^4+|\nabla^2 w|^6+|\nabla^3 w'|^{12}\Bigr)^{\frac{1}{12}}. \tag{3.1.3}$$

Note that we cannot replace w' by w in the last term because $w\in\mathscr{C}^{\frac{5}{2}}$ only.

Proposition 3.1.1. $\gamma_0(x)$ *is a scaling function i.e.* $|x-y|\le\epsilon\gamma_0(x)\implies\gamma_0(y)\asymp\gamma_0(x)$*.*

Proof. (i) If w belonged to $\mathscr{C}^4$ (and we would put w instead of w' in the last term of (3.1.3)) then we would just prove that $|\nabla\gamma_0|\le c$. Here we should be more subtle. We need to prove that if

$$(3.1.4)_{1,2}\qquad \min_j|w-2j\beta|\le\gamma_0^4,\qquad |\nabla w|\le\gamma_0^3,$$

$$(3.1.4)_{3,4}\qquad |\nabla^2 w|\le\gamma_0^2,\qquad |\nabla^3 w'|\le\gamma_0$$

at point x, then at point y the same inequalities hold with γ_0 replaced by $c\gamma_0$ [17]. Definitely this is true for $(3.1.4)_4$ since w' is smooth.

(ii) Consider $|\nabla^2 w|$. Consider $|\Lambda_\alpha w(y)-\nabla^\alpha w(x)|$ with $\Lambda_\alpha=\nabla^\alpha-\frac{1}{3}\delta_{ij}\Delta$, $\alpha=(i,j)$. Then due to (3.1.2)

$$\begin{aligned}(3.1.5)\quad &|\Lambda_{\alpha,y}w(y)-\Lambda_{\alpha,x}w(x)|=\\ &\qquad|\eta\int\bigl(\Lambda_{\alpha,x}|x-z|^{-1}-\Lambda_{\alpha,y}|y-z|^{-1}\bigr)P'_\beta(w(z))\phi(z)\,dz|+\epsilon_1\gamma_0^2\end{aligned}$$

where the last term estimates $|\Lambda_{\alpha,y}w'(y)-\Lambda_{\alpha,x}w'(x)|$ and we used $(3.1.4)_4$. Integrals are understood in the sense of the principal value (vrai) and $\epsilon_1=\epsilon_1(\epsilon)\to+0$ as $\epsilon\to+0$.

---

[17] We need to prove a bit of reverse as well; see (iii).

Note that the integral in expression (3.1.5) does not change if we add to $P'_\beta(w)$ any constant with respect to z.

Let us consider first this integral over $\{z : |x-z| \geq 2\gamma_0\}$, as $\gamma_0 \geq |x-y|$, and note that this integral does not exceed

$$\eta|\int |x-z|^{-4}|x-y|\times \\ \bigl(|\nabla w(x)|\cdot|x-z| + |\nabla^2 w(x)|\cdot|x-z|^2 + |x-z|^{\frac{5}{2}}\bigr)^{\frac{1}{2}}\,dz| \leq C\epsilon_1\eta.$$

Consider now integral over $\{z : |x-z| \leq 2\gamma_0\}$

$$|\eta\int \Lambda_{\alpha,x}|x-z|^{-1}\cdot\bigl(P'_\beta(w(z)) - P'_\beta(w(x))\bigr)\,dz|$$

and note that it also does not exceed $\epsilon_1\eta$. The same arguments work for this integral with x replaced by y but still integrated over $\{z : |x-z| \leq 2\gamma_0\}$ (one needs to remember that $|\nabla w(x)-\nabla w(y)| = O(\gamma_0)$ and $|\nabla^2 w(x)-\nabla^2 w(y)| = O(\gamma_0^{\frac{1}{2}})$).

Further, the same arguments work also for these expressions integrated over $\{z : |y-z| \leq 2\gamma_0\}$ and we are left with

$$|\eta\int \omega(x,y,z)\bigl(P'_\beta(w(x)) - P'_\beta(w(y))\bigr)\,dz|$$

integrated over $\{z :\ |x-z| \asymp \gamma_0\}$ and $\omega \asymp \gamma_0^{-3}$ and one can estimate it by $\epsilon_1\eta$ easily in the same way. Therefore since $|\Delta w(x) - \Delta w(y)|$ does not exceed $\epsilon_1\eta$ we conclude that $|\nabla^2 w(x) - \nabla^2 w(y)|$ does not exceed $\epsilon_1(\eta+\gamma_0^2)$.

However (3.1.2) implies that $\eta \leq c\gamma_0^2$ since $P'_\beta(w) \asymp 1$ in $\mathcal{X}_2$ and therefore $|\nabla^2 w(x) - \nabla^2 w(y)| \leq \epsilon_1\gamma_0^2$ in $B(x,\gamma_0)$.

Combining with $(3.1.4)_2$ we conclude that $|\nabla w(x) - \nabla w(y)| \leq \epsilon_1\gamma_0^3$ in $B(x,\gamma_0)$; next, combining with $(3.1.4)_1$ we conclude that $|w(x)-w(y)| \leq \epsilon_1\gamma_0^4$ in $B(x,\gamma_0)$.

(iii) Therefore $(3.1.4)_{1-4}$ are fulfilled in $y \in B(x,\epsilon\gamma_0)$ with γ_0 replaced by $\gamma_0(1+C\epsilon_1)$. Further, if we redefine γ_0 as the minimal scale such that inequalities $(3.1.4)_{1-4}$ are fulfilled in x, then $(3.1.4)_{1-4}$ fail in $y \in B(x,\epsilon\gamma_0)$ with γ_0 replaced by $\gamma_0(1-C\epsilon_1)$. Therefore with appropriate $\epsilon > 0$ we conclude that $\frac{1}{2} \leq \gamma_0(x)/\gamma_0(y) \leq 2$.

Obviously $\gamma_0 \asymp \gamma_{0\mathsf{old}}$ where $\gamma_{0\mathsf{old}}$ was defined by (3.1.3) and therefore the same conclusion also holds for $\gamma_{0\mathsf{old}}$. $\square$

Now let us reintroduce the scaling function

$$(3.1.3)^* \quad \gamma_0(x) = \epsilon\Bigl(\min_j |w - 2j\beta|^3 + |\nabla w|^4 + |\nabla^2 w|^6 + |\nabla^3 w'|^{12}\Bigr)^{\frac{1}{12}} + C_0\hbar^{\frac{1}{3}} + C_0\eta^{\frac{1}{2}}.$$

Then

(3.1.6) $|x-y| \le 2\gamma_0(x) \implies \gamma_0(y) \asymp \gamma_0(x)$ and $(3.1.4)_{1-4}$ hold (with some constant factor in the right-hand expression).

Consider $B(\bar{x}, \bar{\gamma}_0)$, $\bar{\gamma}_0 = \gamma_0(\bar{x})$, and scale again $x \mapsto \bar{\gamma}_0^{-1}(x-\bar{x})$, $\tau \mapsto \bar{\gamma}_0^{-4}\tau$ and respectively $w \mapsto w_1 = \bar{\gamma}_0^{-4}(w - 2\bar{j}\beta))$, $\hbar \mapsto \hbar_1 = \hbar\bar{\gamma}_0^{-3}$.

If $\bar{\gamma}_0 \asymp \hbar^{\frac{1}{3}}$ then $\hbar_1 \asymp 1$ and we are done. If $\bar{\gamma}_0 \asymp \eta^{\frac{1}{2}}$ then go to *Step 3.*

Step 2.

So, let $\eta_1 \ll 1$. Let us introduce scaling function in $B(0,1)$ obtained after the previous rescaling

$$(3.1.7) \qquad \gamma_1(x) = \epsilon\Bigl(\min_j |w_1 + 2(\bar{j} - j)\beta\bar{\gamma}_0^{-4}|^2 + |\nabla w_1|^3 + |\nabla^2 w_1|^6\Bigr)^{\frac{1}{6}}.$$

Then

$$(3.1.8)_{1-2} \qquad \min_j |w_1 + 2(\bar{j} - j)\beta\bar{\gamma}_0^{-4}| \le C_0\gamma_1^3, \qquad |\nabla w_1| \le C_0\gamma_1^2,$$

$$(3.1.8)_3 \qquad |\nabla^2 w_1| \le C_0\gamma_1.$$

Proposition 3.1.2. *(i)* $\gamma_1(x)$ *is a scaling function:* $|x-y| \le 2\gamma_1(x) \implies \gamma_1(y) \asymp \gamma_1(x)$*;*

(ii) If $\eta_1 \le \epsilon_0$ *then*

$$(3.1.9) \qquad |\nabla^2 w_1(x) - \nabla^2 w_1'(x)| \le \epsilon_2\gamma_1, \qquad w_1' = (w' - \bar{j}\beta)\gamma_0^{-4}.$$

Proof. Proof is similar but simpler than one of Proposition 3.1.1; it is based on the rescaled version of (3.1.1)–(3.1.2):

$$\Delta w_1 = \eta_1 P_\beta (w_1 \bar{\gamma}_0^4 + 2\bar{j}\beta), \qquad \eta_1 = \eta \bar{\gamma}_0^{-2} \le 1 \tag{3.1.10}$$

and therefore in $B(0,1)$

$$w_1 = -\frac{1}{4\pi}\int |x-z|^{-1}\eta_1 P'_\beta (w_1(z)\bar{\gamma}_0^4 + 2\bar{j}\beta)\phi(z)\,dz + w'_1. \tag{3.1.11}$$

$\square$

Now let us reintroduce scaling function

$$\gamma_1(x) = \epsilon\Bigl(\min_j |w_1 + 2(\bar{j}-j)\beta\bar{\gamma}_0^{-4}|^2 + |\nabla w_1|^3 + |\nabla^2 w_1|^6\Bigr)^{\frac{1}{6}} + C_0\hbar_1^{\frac{2}{5}} \tag*{(3.1.7)*}$$

Then

(3.1.12) $|x-y| \le 2\gamma_0(x) \implies \gamma_0(y) \asymp \gamma_0(x)$ and $(3.1.8)_{1-3}$ hold (with some constant factor in the right-hand expression).

Let us consider $\bar{x} \in B(0,1)$ (it is a new point), $B(\bar{x}, \bar{\gamma}_1)$, $\bar{\gamma}_1 = \gamma_1(\bar{x})$, and scale again $x \mapsto \bar{\gamma}_1^{-1}(x-\bar{x})$, $\tau \mapsto \bar{\gamma}_1^{-3}\tau$ and respectively $w_1 \mapsto w_2 = \bar{\gamma}_1^{-3} w_1$, $\hbar_1 \mapsto \hbar_2 = \hbar_1 \bar{\gamma}_1^{-\frac{5}{2}}$.

If $\bar{\gamma}_1 \asymp \hbar_1^{\frac{2}{5}}$ then $\hbar_2 \asymp 1$ and we are done. If $|\nabla w_2| \asymp 1$ we are done as well.

Step 3.

So, consider the remaining case $|\nabla^2 w_2| \asymp 1$. Then we introduce scaling function

$$\gamma_2(x) = \epsilon\Bigl(\min_j |w_2 + 2(\bar{j}-j)\beta\bar{\gamma}_0^{-4}\bar{\gamma}_1^{-3}| + |\nabla w_2|^2\Bigr)^{\frac{1}{2}} + C_0\hbar_2^{\frac{1}{2}}. \tag{3.1.13}$$

Let us consider $\bar{x} \in B(0,1)$ (it is a new point), $B(\bar{x}, \bar{\gamma}_2)$, $\bar{\gamma}_2 = \gamma_2(\bar{x})$, and scale again $x \mapsto \bar{\gamma}_2^{-1}(x-\bar{x})$, $\tau \mapsto \bar{\gamma}_2^{-2}\tau$ and respectively $w_2 \mapsto w_3 = \bar{\gamma}_2^{-2} w_2$, $\hbar_2 \mapsto \hbar_3 = \hbar_2 \bar{\gamma}_2^{-2}$.

If $\bar{\gamma}_2 \asymp \hbar_1^{\frac{1}{2}}$ then $\hbar_3 \asymp 1$ and we are done. If $|\nabla w_3| \asymp 1$ we are done as well.

Step 4.

Finally, introduce

$$\gamma_3(x) = \epsilon \min_j |w_3 + 2(\bar{j} - j)\beta\bar{\gamma}_0^{-4}\bar{\gamma}_1^{-3}\bar{\gamma}_2^{-2}| + C\hbar_3^{\frac{2}{3}}. \tag{3.1.14}$$

3.1.2 Scaling arguments in zone $\mathcal{X}_2$: Semiclassical N-term.

Now we apply scaling arguments using scaling functions constructed above.

We revert our steps. While we call γ_{1-3} *relative scaling functions* let us introduce *absolute scaling functions* $\alpha_0(x) = \gamma_0(x)$, $\alpha_1(x) = \gamma_0(x)\gamma_1(x)$, $\alpha_2(x) = \gamma_0(x)\gamma_1(x)\gamma_2(x)$, and $\alpha_3(x) = \gamma_0(x)\gamma_1(x)\gamma_2(x)\gamma_3(x)$ [18].

We need first

Proposition 3.1.3. *Consider* $B(0,1)$ *and assume that in it*

$$|\nabla w| \asymp \theta, \tag{3.1.15}$$

and

$$|\nabla^2 w| \le c\theta. \tag{3.1.16}$$

Let

$$\gamma(x) := \epsilon \min_j |w - 2\mu\hbar j|\theta^{-1} + \hbar_*^{\frac{2}{3}} \tag{3.1.17}$$

and

$$\varepsilon \ge (\hbar_*)^{\frac{2}{3}-\delta}, \qquad \hbar_* = \hbar\theta^{-\frac{1}{2}}. \tag{3.1.18}$$

Let $\varphi \in \mathscr{C}_0^\infty([-\epsilon_0, \epsilon_0])$. *Then for* $\alpha \le \bar{\gamma} := \gamma(\bar{x})$,

$$\begin{gathered}|\int \phi_\alpha(x)\Bigl(e_\varphi(x,x,\tau) - P'_{\mu\hbar,\varphi}\bigl(w(x)+\tau\bigr)\Bigr)\,dx| \le \\ C\mu\hbar^{-1}\alpha^3 + C\mu\hbar^{-1}\alpha^3\bar{\gamma}^{-1}(\hbar_*\bar{\gamma}^{-\frac{1}{2}}\alpha^{-1})^s\end{gathered} \tag{3.1.19}$$

with

$$e_\varphi(x,y,\tau) := \varphi\bigl(\hbar^2 D_{x_3}^2(\mu\hbar)^{-1}\bigr)e(x,y,\tau) \tag{3.1.20}$$

---

[18] So far we ignore the first scaling $x \mapsto (x - \bar{x})\ell^{-1}$.

and Weyl expression

$$P'_{\beta,\varphi}(w+\tau) = \mathsf{const}\sum_j \beta(w+\tau-2j\mu\hbar)^{\frac{1}{2}}\varphi(w+\tau-2j\mu\hbar) \tag{3.1.21}$$

with the standard constant where for each x *only one term is present in this sum. Here we take large* $s>0$ *as* $\hbar_*\le\bar\gamma\alpha$ *and* $s=0$ *otherwise.*

Proof. The proof is standard and based on the standard reduction to canonical form, standard estimates for $U(x,y,t)$ a Schwartz kernel of $e^{i\hbar_*^{-1}\theta^{-1}tH}$:

$$|F_{t\to\hbar_*^{-1}\tau}\int \bar\chi_T(t)\phi_\alpha(x)U_\varphi(x,x,t)\,dx|\le C\mu\hbar^{-1}\alpha^3 \tag{3.1.22}$$

as $T\asymp 1$ and

$$|F_{t\to\hbar_*^{-1}\tau}\int \phi_\alpha(x)\bigl(\bar\chi_T(t)-\bar\chi_{\bar T}(t)\bigr)U_\varphi(x,x,t)\,dx|\le \\ C\mu\hbar^{-1}\alpha^3(\hbar_*\bar\gamma^{-\frac{1}{2}}\alpha^{-1})^s \tag{3.1.23}$$

with $\bar T=\epsilon\gamma^{\frac{1}{2}}$ where U_φ is defined similarly to (3.1.20).

Here obviously we can skip in (3.1.19) all perturbation terms in the argument and in ϕ_α transformed. □

Then plugging into (3.1.19) $\alpha=\gamma$ $(=\bar\gamma)$, we have factor $(\hbar\gamma^{-\frac{3}{2}})^s$ in the second term.

There are two cases: $\theta\le\mu\hbar$ and $\theta\ge\mu\hbar$.

In the former case taking the sum over γ-partition of 1-element we estimate the same expressions with ϕ_1 instead of ϕ_γ by their right-hand expressions integrated over $\gamma^{-3}\,d\gamma$ which return $C\mu\hbar^{-1}$.

On the other hand, in the latter case let $\lambda=\mu h\theta^{-1}$. Taking the sum over γ_3-partition of λ-element ϕ_λ by the right-hand expressions which return $C\mu\hbar_2^{-1}\lambda^2$. In this case summation over λ-partition return $C\theta\hbar^{-2}$.

In both cases we arrive to

$$|\int \phi(x)\Bigl(e_\varphi(x,x,\tau)-P'_{\mu\hbar,\varphi}\bigl(w(x)+\tau\bigr)\Bigr)\,dx|\le C\theta\hbar^{-2}+C\mu\hbar^{-1}. \tag{3.1.24}$$

Applying this estimate after α_2-scaling we conclude that the left-hand expression with $\phi=\phi_{\alpha_2}$ (in the non-scaled settings) does not exceed

$C\theta\hbar^{-2}\alpha_2^2 + C\mu\hbar^{-1}\alpha_2^2$. Here the first term is $O(\hbar^{-2}\alpha_2^3)$ and the summation over 1-element returns $O(\hbar^{-2})$.

Consider the second term $C\mu\hbar^{-1}\alpha_2^2 = C\mu\hbar^{-1}\gamma_0^2\gamma_1^2\gamma_2^2$. Then summation over α_2-partition of α_1-element returns

$$C\mu\hbar^{-1}\gamma_0^2\gamma_1^2\int \gamma_2^2\times\gamma_2^{-3}\,dx \asymp C\mu\hbar^{-1}\gamma_0^2\gamma_1^2(1+|\log\hat{\gamma}_2|)$$

where $\hat{\gamma}_k$ is a minimal value of γ_k over γ_{k-1}-element. However in fact there will be no logarithmic factor because in virtue of equation (3.1.1) there is a positive eigenvalue of $\mathsf{Hess}\, w_2$ of the maximal size (cf. Section 5.1.1) of [Ivr11]. Therefore, in fact we have $C\mu\hbar^{-1}\gamma_0^2\gamma_1^2$.

Now summation summation over α_1-partition of α_0-element returns

$$C\mu\hbar^{-1}\gamma_0^2\int \gamma_1^2\times\gamma_1^{-3}\,dx \asymp C\mu\hbar^{-1}\gamma_0^2(1+|\log\hat{\gamma}_1|).$$

Finally, summation over α_0-partition of 1-element returns

$$C\mu\hbar^{-1}\int \gamma_0^2(1+|\log\hat{\gamma}_1|)\times\gamma_0^{-3}\,dx \asymp C\mu\hbar^{-1}\hat{\gamma}_0^{-1}(1+|\log\check{\gamma}_1|).$$

where $\check{\gamma}_k$ is an absolute minimum of γ_k. However $\gamma_0^2\geq\eta$ and $\gamma_1\geq\eta$ and therefore expression above does not exceed $C\mu\hbar^{-1}\eta^{-\frac{1}{2}}(1+|\log\eta|)$.

Remark 3.1.4. Recall that we estimated only cut-off expression. To calculate the full expression we need to calculate also the contribution of zone $\{\xi_3^3\geq \mu h\}$. However this is easy.

Really, instead of $\varphi(\hbar^2D_3^2/(\mu\hbar))$ consider $\varphi'(\hbar^2D_3^2/\theta)$ with $\varphi'\in\mathscr{C}^\infty([1,4])$ and $\epsilon\mu h\leq\theta\leq 1$. Without any scaling one can prove easily that such modified expression (3.1.24) does not exceed $C\theta\hbar^{-2}$. We leave easy details to the reader.

Therefore plugging $\theta=2^n\mu\hbar$ and taking a sum over $n=0,\dots,\lfloor|\log_2\mu\hbar|\rfloor$ we get required expressions. Also note that in such expressions we need to consider perturbed argument $w+\mu^{-2}\omega_1+j\mu^{-1}\hbar\omega_2$ (all other terms which are $O(\mu^{-4}+\mu^{-\frac{3}{2}}\hbar)$ could be skipped and also a perturbed function transformed.

Remark 3.1.5. (i) However we need to do get rid off these perturbations for $\theta\leq\mu\hbar^{1-\delta}$ only. Really, for $\theta\geq\mu\hbar^{1-\delta}$ we need canonical form *only* to study propagation and calculations could be performed without it. But then getting rid off perturbation is trivial provided perturbation does not exceed $C\mu\hbar^{1+\delta}$ which is the case if

$$\mu \ge \hbar^{-\frac{1}{3}-\delta}. \tag{3.1.25}$$

(ii) Note that in the smooth approximation contributions of $\mathcal{X}_1$ is always less than $CZ^{\frac{2}{3}-\delta_1}$, $C\max(Z^{\frac{5}{3}}, Z^{\frac{3}{5}}B^{\frac{4}{5}})Z^{-\delta_1}$ or $C\max(Z^{\frac{5}{3}}, Z^{\frac{3}{5}}B^{\frac{4}{5}})Z^{-\delta_1} + CB^{\frac{1}{2}}Z^{\frac{7}{6}-\delta_1}$ respectively with an exception of first two and only in the case of the threshold value between $\ge Z^{-\frac{1}{3}-\delta_2}$. However in this case $\eta \ge Z^{-\delta_3}$ and the errors of the smooth approximation approach in fact are less than $CZ^{\frac{2}{3}-\delta_4}$, $CZ^{\frac{5}{3}-\delta_4}$ as well. Therefore there are in fact no exception.

(iii) It is important to have $\varepsilon \le \mu\hbar^*$ and with $\varepsilon = \hbar_*^{\frac{2}{3}-\delta}$ we need $\mu \ge \hbar^{-\frac{1}{3}-\delta}\theta^{\frac{1}{3}-\frac{1}{2}\delta}$ which is due to (3.1.25).

Therefore we conclude that in completely non-scaled settings with $\phi = \phi_\ell(x)$

$$\begin{multline} |\int \phi(x)\Bigl(e(x,x,\tau) - P'_B\bigl(W(x)+\tau\bigr)\Bigr)| \le \\ C\zeta^2\ell^2 + CB\ell\zeta^{-\frac{1}{2}}(1+|\log \ell^2\zeta|) \end{multline} \tag{3.1.26}$$

where the first term is $C\hbar^{-2}$ and the second term is $C\mu\hbar^{-1}\eta^{-\frac{1}{2}}(1+|\log\eta|)$; recall that $\hbar^{-1} \asymp \ell\zeta$, $\mu \asymp B\ell\zeta^{-1}$ and $\eta \asymp \ell^2\zeta$. In comparison with the non-degenerate case $|\nabla W_B^{\mathsf{TF}}| \asymp \zeta^2\ell^{-1}$ we acquired the last term.

Assume first that

$$Z_m \asymp Z \qquad \forall m = 1, \dots, M. \tag{3.1.27}$$

Then

(i) As $B \le Z^{\frac{4}{3}}$, $\ell \le Z^{-\frac{1}{3}}$ we have $\zeta = Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}$ and the right-hand expression of (3.1.26) returns $CZ\ell + CB\ell^{\frac{5}{4}}Z^{-\frac{1}{4}}$ and the summation with respect to ℓ results in its value as $\ell = Z^{-\frac{1}{3}}$ i.e. $CZ^{\frac{2}{3}} + CBZ^{-\frac{2}{3}}$ with the dominating first term.

(ii) As $B \le Z^{\frac{4}{3}}$, $\ell \ge Z^{-\frac{1}{3}}$ we have $\zeta = \ell^{-2}$ and the right-hand expression of (3.1.26) returns $C\ell^{-2} + CB\ell^2$. We need to sum as long as $\mu h \le 1$ i.e. $Z^{-\frac{1}{3}} \le \ell \le B^{-\frac{1}{4}}$ and the summation returns $CZ^{\frac{2}{3}} + CB^{\frac{1}{2}}$ with the dominating first term.

(iii) As $Z^{\frac{4}{3}} \le B \le Z^2$, $\ell \le B^{-1}Z$ we have $\zeta = Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}$ and the right-hand expression of (3.1.26) returns $CZ\ell + CBZ^{-\frac{1}{4}}\ell^{\frac{5}{4}}$. Then summation results in $CZ^2B^{-1} + CZ^2B^{-\frac{1}{4}}Z \lesssim Z^{\frac{2}{3}}$.

If assumption (3.1.27) is not fulfilled we can estimate in the first term of the right-hand expression of (3.1.26) parameter ζ from above by $\mathsf{min}(Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}, \ell^{-2})$ and in the second term from below by $\zeta_m = \mathsf{min}(Z_m^{\frac{1}{2}}\ell_m^{-\frac{1}{2}}, \ell_m^{-2})$ as $\ell = \ell_m := |x - \mathsf{y}_m|$ and repeat all arguments above.

Therefore we arrive to the statement (i) below. Furthermore, as $B \le Z$ zone $\mathcal{X}_2$ is contained in $\{x : \ell(x) \ge B^{-\frac{3}{10}} \ge Z^{-\frac{3}{10}}$ (really, $\mu \ge h^{-\frac{1}{3}}$ in $\mathcal{X}_2$) and we arrive to the statement (ii) below.

Proposition 3.1.6. *(i) As $B \le Z^2$ contribution of zone $\mathcal{X}_2$ to expression*

$$\int \bigl(e(x,x,\nu) - P'_B(W(x)+\nu)\bigr)\,dx \tag{3.1.28}$$

does not exceed $CZ^{\frac{2}{3}}$;

(ii) As $B \le Z$ contribution of zone $\mathcal{X}_2$ to expression (3.1.28) *does not exceed $CZ^{\frac{2}{3}-\delta}$.*

3.1.3 Scaling arguments in zone $\mathcal{X}_2$: Semiclassical D-term

We need to estimate

$$\mathsf{D}\bigl(\phi_\alpha\bigl(e(x,x,\nu) - P'_B(W(x)+\nu)\bigr), \phi_\alpha\bigl(e(x,x,0) - P'_B(W(x))\bigr)\bigr) \tag{3.1.29}$$

where $\phi_\alpha(x)$ is an α-admissible function. Again we revert our steps.

Consider $B(x,\bar{\alpha}_3)$ and apply Fefferman–de Llave decomposition. Then in the framework of Proposition 3.1.3 contribution of pairs $B(x,\alpha)$ and $B(y,\alpha)$ with $3\alpha \le |x-y| \le 4\alpha$ does not exceed the right-hand expression of (3.1.19) squared and multiplied by α^{-4} where $C\alpha^{-3}$ estimates the number of the pairs and α^{-1} is the inverse distance. At this moment we discuss a cut-off version of (3.1.29) i.e. with $e_\varphi(.,.,.)$ and $P'_{\mu\hbar,\varphi}(.)$. So, we have

$$C\mu^2\hbar^{-2}\bigl(1 + \gamma_3^{-1}(\hbar_2\gamma_3^{-\frac{1}{2}}\alpha^{-1})^s\bigr)^2\alpha^4.$$

Then integrating this expression with respect to $\alpha^{-1}\,d\alpha$ with $\alpha\le\gamma_3$ we arrive to $C\mu^2\hbar^{-2}\bigl(\gamma_3^4+\hbar_2^2\gamma_3\bigr)$.

Therefore

(3.1.30) A cut-off version of (3.1.29) with $\alpha=\alpha_3$ does not exceed $C\mu^2\hbar^{-2}\bigl(\gamma_3^4+\hbar_2^2\gamma_3\bigr)\alpha_2^3$

The first term here $C\mu^2\hbar^{-2}\gamma_3^4\alpha_2^3$ does not exceed $C\mu^2\hbar^{-2}\alpha_3^3$ (recall that $\alpha_j=\alpha_{j-1}\gamma_j$) and the summation over α_3-partition of 1-element returns $C\mu^2\hbar^{-2}$.

Consider the second term $C\mu^2\hbar^{-2}\hbar_2^2\gamma_3\alpha_2^3$; its summation with respect to α_3-partition of α_2-element returns $C\mu^2\hbar^{-2}\alpha_2^3\hbar_2^2\int\gamma_3^{-2}\,d\gamma_3\lesssim C\mu^2\hbar^{-2}\alpha_2^3$ (really, recall that according to (3.1.14) $\gamma_3\ge\hbar_3^{\frac{2}{3}}$) and then the summation over α_2-partition of 1-element returns $C\mu^2\hbar^{-2}$.

Consider $B(x,\bar\alpha_2)$ and apply Fefferman–de Llave decomposition. There are two kinds of pairs:

(a) those with $|x-y|\ge\epsilon(\alpha_3(x)+\alpha_3(y))$ for all (x,y) and

(b) those with $|x-y|\le\min(\alpha_3(x),\alpha_3(y))$ for all (x,y).

The total contribution of the pairs of the second type (i.e. summation is taken over *all* pairs of α_3-elements in $B(0,1)$) as we already know is $O(\mu^2\hbar^{-2})$. Meanwhile according to the analysis in the previous subsubsection *Scaling arguments in zone* $\mathcal{X}_2$*: Semiclassical* N*-term* a contribution of one pair of kind (a) does not exceed

$$C\underbrace{\bigl(\hbar^{-2}+\mu\hbar^{-1}\gamma_0^{-1}\gamma_1^{-1}\gamma_2^{-1}\gamma_3^{-1-\delta}\bar\gamma_3^{\delta}\bigr)\alpha_3^3}_{\text{at }x}\times\\
\underbrace{\bigl(\hbar^{-2}+\mu\hbar^{-1}\gamma_0^{-1}\gamma_1^{-1}\gamma_2^{-1}\gamma_3^{-1-\delta}\bar\gamma_3^{\delta}\bigr)\alpha_3^3}_{\text{at }y}\times|x-y|^{-1}$$

where each of two first factors is just an estimate of the integral (3.1.24) calculated over corresponding argument. If we take the first term in the first factor and sum over α_3-partition of 1-element we get only the second factor multiplied by $\mu\hbar^{-1}$ and then summation was done in the previous subsubsection. Similarly we can deal with the first term in the second factor.

On the other hand, if we take only second factors and sum over pairs of α_3-subelements of the same α_2-element we get

$$C\mu^2\hbar^{-2}\gamma_0^{-2}\gamma_1^{-2}\gamma_2^{-2}\alpha_2^5 \asymp C\mu^2\hbar^{-2}\alpha_2^3.$$

Then summation with respect to α_2-partition of 1-element returns $C\mu^2\hbar^{-2}$.

Consider now $B(\bar{x},\bar{\alpha}_1)$ and apply here Fefferman-de Llave decomposition. There are two kinds of pairs:

(a) those with $|x-y|\ge \epsilon(\alpha_2(x)+\alpha_2(y))$ for all (x,y) and

(b) those with $|x-y|\le \min(\alpha_2(x),\alpha_2(y))$ for all (x,y).

According to the above analysis the total contribution of the pairs of the second type (i.e. the summation is taken over *all* pairs of α_2-elements in $B(0,1)$) as we already know is $O(\mu^2\hbar^{-2})$. Meanwhile according to the analysis in the previous subsubsection *Scaling arguments in zone $\mathcal{X}_2$: Semiclassical* N-*term* a contribution of one pair of kind (a) does not exceed

$$C\underbrace{\bigl(\hbar^{-2}+\mu\hbar^{-1}\gamma_0^{-1}\gamma_1^{-1}\gamma_2^{-1}\bigr)\alpha_3^3}_{\text{at } x}\times\underbrace{\bigl(\hbar^{-2}+\mu\hbar^{-1}\gamma_0^{-1}\gamma_1^{-1}\gamma_2^{-1}\bigr)\alpha_3^3}_{\text{at } y}\times|x-y|^{-1}$$

and here again we can "forget" about the first terms in each factor. Then the summation with respect to pairs of α_2-subelements of the same α_1-element results in $C\mu^2\hbar^{-2}\gamma_0^{-2}\gamma_1^{-2}\alpha_1^5\asymp C\mu^2\hbar^{-2}\alpha_1^3$ where we avoid logarithmic factor in virtue of the same positive eigenvalue of $\operatorname{Hess} w$. Summation with respect to α_1-admissible partition of 1-element returns $C\mu^2\hbar^{-2}$.

Consider now $B(\bar{x},\bar{\alpha}_0)$ and apply here Fefferman–de Llave decomposition. Again there are two kinds of pairs and the total contributions of the pairs of the second kind we already calculated and contribution of the pairs of α_1-subelements of the same 1-element does not exceed

$$C\mu^2\hbar^{-2}\gamma_0^{-2}(1+|\log\hat{\gamma}_1|)^2\alpha_0^5\lesssim C\mu^2\hbar^{-2}(1+|\log\eta|)^2\alpha_0^3$$

and the summation with respect to α_0-partition of 1-element returns $C\mu^2\hbar^{-2}(1+|\log\eta|)^2$.

Finally, consider $B(\bar{x},1)$ and apply here Fefferman-de Llave decomposition. Again there are two kinds of pairs and the total contributions of the pairs of kind (b) we already estimated while the total contribution of the pairs of kind (a) does not exceed $C\hbar^{-4}+C\mu^2\hbar^{-2}\eta^{-1}(1+|\log\eta|)^2$ where we recalled the forgotten terms.

Again, this is estimate for cut-off expression. Going to uncut expression we repeat the same trick as before but as we deal with D-term we need to consider "mixed" pairs when one "factor" comes with θ and another with θ' but then contribution of such pair does not exceed $C(\nu\hbar^{-4})^{\frac{1}{2}}(\nu'\hbar^{-4})^{\frac{1}{2}}$. Easy details are left to the reader.

Therefore returning to the original scale we conclude that the contribution of ℓ-layer to (3.1.29) does not exceed

$$C\zeta^4\ell^3 + CB^2\ell\zeta^{-1}(1+|\log \ell^2\zeta|)^2 \tag{3.1.31}$$

which is the right-hand expression of (3.1.26) squared and multiplied by ℓ^{-1} due to scaling. In comparison with the non-degenerate case $|\nabla W_B^{\mathsf{TF}}| \asymp \zeta^2\ell^{-1}$ we acquired the last term.

Assume first that condition (3.1.27) is fulfilled. Then

(i) As $B \le Z^{\frac{4}{3}}$, $\ell \le Z^{-\frac{1}{3}}$ we have $\zeta = Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}$ and expression (2.2.18) returns $CZ^2\ell + CB^2\ell^{\frac{3}{2}}Z^{-\frac{1}{2}}$ and the summation with respect to ℓ results in its value as $\ell = Z^{-\frac{1}{3}}$ i.e. $CZ^{\frac{5}{3}} + CB^2Z^{-1}$ with the dominating first term.

(ii) As $B \le Z^{\frac{4}{3}}$, $\ell \ge Z^{-\frac{1}{3}}$ we have $\zeta = \ell^{-2}$ and expression (2.2.18) returns $C\ell^{-5} + CB^2\ell^3$. We need to sum as long as $\mu h \le 1$ i.e. $Z^{-\frac{1}{3}} \le \ell \le B^{-\frac{1}{4}}$ and the summation returns $CZ^{\frac{5}{3}} + CB^{\frac{5}{4}}$ with the dominating first term.

(iii) As $Z^{\frac{4}{3}} \le B \le Z^2$, $\ell \le B^{-1}Z$ we have $\zeta = Z^{\frac{1}{2}}\ell^{-\frac{1}{2}}$ and expression (2.2.18) returns $CZ^2\ell + CB^2Z^{-\frac{1}{2}}\ell^{\frac{3}{2}}$. Then summation results in $CZ^3B^{-1} + CB^{\frac{1}{2}}Z \lesssim Z^{\frac{3}{5}}B^{\frac{4}{5}}$.

Sure, we need to consider also mixed pairs of the layers and their contributions are

$$C\bigl(\zeta^2\ell^2 + CB\ell\zeta^{-\frac{1}{2}}(1+|\log \ell^2\zeta|)\bigr)\times\bigl(\zeta'^2\ell'^2 + CB\ell'\zeta'^{-\frac{1}{2}}(1+|\log \ell'^2\zeta'|)\bigr)\times(\ell+\ell')^{-1}$$

and the summation with respect to ℓ and ℓ' returns the same expression as above.

If assumption (3.1.27) is not fulfilled we use the same trick as in the previous subsubsection. Therefore we arrive to the statement (i) below. Applying the same arguments as in the proof of Proposition 3.1.6 we arrive to the statement (ii) below.

Proposition 3.1.7. *(i) As $B \le Z^2$ contribution of zone $\mathcal{X}_2 \times \mathcal{X}_2$ to expression* (3.1.29) *does not exceed $C\max(Z^{\frac{5}{3}}, Z^{\frac{3}{5}}B^{\frac{4}{5}})$;*

(ii) As $B \le Z$ contribution of zone $\mathcal{X}_2$ to expression (3.1.29) *does not exceed* $CZ^{\frac{5}{3}-\delta}$.

3.1.4 Semiclassical T-term: zone $\mathcal{X}_1$ extended

What is $\mathcal{X}_1$ extended?

First, let us analyze the precise extension in the framework of D- and N-terms. For these terms we have errors $C(\mu\hbar)^2\eta^{-\frac{1}{2}}\hbar^{-3} = CB^2\zeta^{-\frac{3}{2}}\ell^2$ and this expression squared and multiplied by ℓ^{-1} i.e. $CB^4\zeta^{-3}\ell^3$ respectively, and both sum to their values as $\ell = r$.

(i) As $B \le Z^{\frac{4}{3}}$ and $r \ge Z^{-\frac{1}{3}}$ we have CB^2r^5 and CB^4r^9 and we want them to be less than $CZ^{\frac{2}{3}}$ and $CZ^{\frac{5}{3}}$ respectively. As $B < Z^{\frac{7}{6}}$ the first requirement is more restrictive and we set $CB^2r^5 = Z^{\frac{2}{3}}$ i.e. $r = B^{-\frac{2}{5}}Z^{\frac{2}{15}}$ and then $\mu = Br^3 = B^{-\frac{1}{5}}Z^{\frac{6}{15}} \ge Z^{\frac{1}{6}}$ and $\hbar = B^{-\frac{2}{5}}Z^{\frac{2}{15}} \ge Z^{-\frac{1}{3}}$ so $\mu \ge \hbar^{-\frac{1}{2}}$.

(ii) As $Z^{\frac{7}{6}} \le B \le Z^{\frac{4}{3}}$ the second requirement is more restrictive and we set $B^4Z^{-\frac{3}{2}}r^{\frac{9}{2}} = Z^{\frac{5}{3}}$, i.e. $r = B^{-\frac{8}{9}}Z^{\frac{19}{27}}$ and $\mu = B^{-\frac{1}{3}}Z^{\frac{5}{9}}$ and $\hbar = Z^{-\frac{1}{2}}r^{-\frac{1}{2}} = B^{\frac{4}{9}}Z^{-\frac{23}{27}}$ and $\mu \ge \hbar^{-\frac{3}{7}}$ (better than $\hbar^{-\frac{1}{3}}$).

But $\mu\hbar \le \eta$ iff $r \ge B^2Z^{-3}$ i.e. $B \le Z^{\frac{50}{39}}$ and this is less than $Z^{\frac{4}{3}}$ so we test μ and $\hbar$ in this case: $\mu = Z^{\frac{5}{39}}$ and $\hbar = Z^{\frac{11}{39}}$ and $\mu \ge \hbar^{-\frac{5}{11}}$. As $B \ge Z^{\frac{5}{11}}$ we have error $(\mu\hbar)^3\hbar^{-6}r^{-1} = B^3r^5$ and we want it to $\le Z^{\frac{5}{3}}$, so $r = Z^{\frac{1}{3}}B^{-\frac{3}{5}}$ and $\mu = B^{\frac{1}{10}}$ and $\hbar = B^{\frac{3}{10}}Z^{-\frac{2}{3}}$ and we test it as $B = Z^{\frac{4}{3}}$ when $\mu = B^{\frac{1}{10}}$ and $\hbar = B^{-\frac{1}{5}}$, so exponent $-\frac{5}{11}$ fits.

As $B \le Z^{\frac{4}{3}}$ we consider $B^3r^5 = B^{\frac{4}{5}}Z^{\frac{3}{5}}$ i.e. $r = B^{-\frac{11}{25}}Z^{\frac{3}{25}}$, $\mu = B^{\frac{17}{50}}Z^{-\frac{8}{25}}$, $\hbar = B^{\frac{11}{50}}Z^{-\frac{14}{25}}$ and exponent $-\frac{5}{11}$ fits as well.

When we can use the same method for T-term?

As far as semiclassical T-expression is concerned an error of such approach in the localized and scaled settings would be $C(\mu\hbar)^{\frac{5}{2}}\hbar^{-3}$ which is $O(\hbar^{-1})$ only as $\mu \ge \hbar^{-\frac{1}{5}}$. One can extend it to $\mu \ge \hbar^{-\frac{1}{5}-\delta}$ using the same trick as in Remark 3.1.5 but we need to do better than this.

Note first that in fact an error does not exceed $C(\mu\hbar)^3\eta^{-\frac{1}{2}}\hbar^{-3} \asymp CB^3\ell^2\zeta^{-\frac{7}{2}}$ in the localized scaled settings (we skip $C\hbar^{-1}$ as the result here is already taken into account). The simple proof is left to the reader. This is translated into $CB^3\ell^2\zeta^{-\frac{3}{2}}$ into unscaled settings. Summation with respect to $\ell \le r$ returns its value as $\ell = r$.

So we get CB^3r^5 as $B \le Z^{\frac{4}{3}}$ and $r \ge Z^{-\frac{1}{3}}$. Let $B \le Z$. Then we want $CB^3r^5 \le CZ^{\frac{5}{3}}$ i.e. $r = B^{-\frac{3}{5}}Z^{\frac{1}{3}}$ which is greater than $Z^{-\frac{1}{3}}$ as $B \le Z^{\frac{10}{9}}$. Then $\mu = Br^3 = B^{-\frac{4}{5}}Z \ge Z^{\frac{1}{5}}$ and $\hbar = r = B^{-\frac{3}{5}}Z^{\frac{1}{3}} \ge Z^{-\frac{4}{15}}$ and $\mu \ge \hbar^{-\frac{3}{4}}$.

As $Z \le B \le Z^{\frac{4}{3}}$ but $r \ge Z^{-\frac{1}{3}}$ still we want $CB^3r^5 \le CB^{\frac{1}{3}}Z^{\frac{4}{3}}$ i.e. $r = B^{-\frac{8}{15}}Z^{\frac{4}{15}}$ and we want it to be $\ge Z^{-\frac{1}{3}}$ i.e. $B \le Z^{\frac{9}{8}}$. Then $\mu = Br^3 = B^{-\frac{3}{5}}Z^{\frac{4}{5}} \ge Z^{\frac{1}{8}}$ and $\hbar = B^{-\frac{8}{15}}Z^{\frac{4}{15}} \ge B^{-\frac{1}{3}}$ and $\mu \ge \hbar^{-\frac{3}{8}}$. It is not as good as $\mu \ge \hbar^{-\frac{3}{7}}$.

Then we use the smooth canonical form. In the operator perturbation terms have factors μ^{-2}, μ^{-4} etc and we can use the standard approach to get rid off $\mu^{-4} \le \mu\hbar$, so we need to consider only μ^{-2}.

However let before scaling the second derivative of W be of magnitude θ; then after scaling it becomes of magnitude $\theta' = \gamma_0^2\gamma_1^2\theta$ and then the perturbation is of magnitude $\theta\mu^{-2}$ but contribution of the error will be (after we compare the true Riemann sum and the corresponding integral and their difference $C\hbar^{-3}\nu\mu^{-2}(\mu\hbar)^2(\theta')^{-\frac{1}{2}} \times \alpha_1^3 \le C\theta^{\frac{1}{2}}\hbar^{-1}\gamma_0^{-1}\gamma_1^{-1}\alpha_1^3 \le C\hbar^{-1}\gamma_1^{-\frac{1}{2}}\alpha_1^3$ where we used that $\theta \le C\gamma_0^2\gamma_1$. Then summation over α_1-partition of α_0 element returns $C\hbar^{-1}\alpha_0^3$ and the summation over γ_0-partition returns $C\hbar^{-1}$ as desired. Therefore we covered zone $\mathcal{X}_1$ for T-term.

3.1.5 Semiclassical T-term: zone $\mathcal{X}_2$

Tauberian estimate.

Tauberian estimate for cut-off expression is rather simple:

$$C\mu\hbar^{-1}\gamma_0^{-1}\gamma_1^{-1}\gamma_2^{-1} \times \hbar\gamma_0^{-3}\gamma_1^{-\frac{5}{2}}\gamma_1^{-2} \times \gamma_0^4\gamma_1^3\gamma_2^2\alpha_2^3 \asymp C\mu\gamma_1^{-\frac{1}{2}}\gamma_2^{-1}\alpha_2^3$$

which nicely sums to $C\mu$ without logarithm due to the same positive eigenvalue arguments as before; for θ-cut-off with $\theta \ge \mu\hbar$ we get the same albeit with γ_j defined by the same formula albeit with $(w_j - 2j\mu\hbar s_j^{-1})$ replaced by θs_j^{-1} where s_j^{-1} means the scale; and this should be multiplied by $\theta/(\mu\hbar)$. The result nicely sums to $C\hbar^{-1}$. This is what was required.

Magnetic Weyl expression.

Now we will get the same answer albeit $C\mu^{-4}$ term will be supplemented by $C\mu^{-\frac{3}{2}}\hbar$ which in cut-off sum adds $C\mu^{-\frac{3}{2}}\hbar \times \mu\hbar^{-2} \le C\hbar^{-1}$.

We can use the standard approach, with an error $C\mu^{-\frac{3}{2}}\hbar \times \theta/(\mu\hbar) \times \mu\hbar^{-2} \asymp C\theta\mu^{-\frac{3}{2}}\hbar^{-2}$ which means that we can take $\theta = \mu^{\frac{3}{2}}\hbar$ which is sufficient to deal with with $\theta \ge C\mu^{\frac{3}{2}}\hbar$; in particular, for $\mu \ge \hbar^{\frac{2}{3}}$ we are done. But for $\theta \ge \mu\hbar^{1-\delta}$ we can apply the weak magnetic field approach, which is sufficient. So we arrive to inequality

$$|\int_{-\infty}^{\tau}\int \phi(x)\Bigl(e_{\varphi}(x,x,\tau) - P'_{\beta,\varphi}\bigl(w(x)+\tau\bigr)\Bigr)\,dx\,d\tau| \le C\hbar^{-1} \tag{3.1.32}$$

and therefore

Proposition 3.1.8. *(i) As $B \le Z^2$ contribution of zone $\mathcal{X}_2$ to expression*

$$\int_{-\infty}^{\tau}\int \phi(x)\Bigl(e_{\varphi}(x,x,\tau) - P'_{B,\varphi}\bigl(W(x)+\tau\bigr)\Bigr)\,dx\,d\tau \tag{3.1.33}$$

does not exceed $C\max\bigl((Z+B)^{\frac{1}{3}}Z^{\frac{4}{3}}, Z^{\frac{3}{5}}B^{\frac{4}{5}}\bigr)$;

(ii) As $B \le Z$ contribution of zone $\mathcal{X}_2$ to expression (3.1.33) *does not exceed $CZ^{\frac{5}{3}-\delta}$.*

3.1.6 Mollification errors

We need to estimate

$$\int \phi(x)\Bigl(P'_B(W(x)+\tau)) - P'_B(W_B^{\mathsf{TF}}(x)+\tau))\Bigr)\,dx, \tag{3.1.34}$$

$$\int \phi(x)\Bigl(P_B(W(x)+\tau)) - P_B(W_B^{\mathsf{TF}}(x)+\tau))\Bigr)\,dx, \tag{3.1.35}$$

$$\begin{multline}\mathsf{D}\Bigl(\phi(x)\bigl(P'_B(W(x)+\tau)) - P'_B(W_B^{\mathsf{TF}}(x)+\tau))\bigr),\\ \phi(x)\bigl(P'_B(W(x)+\tau)) - P'_B(W_B^{\mathsf{TF}}(x)+\tau))\bigr)\Bigr)\end{multline} \tag{3.1.36}$$

and

$$\|\phi(x)\nabla\bigl(W(x) - W_B^{\mathsf{TF}}(x)\bigr)\|^2. \tag{3.1.37}$$

We start from local versions (so in fact we dealing with w and w_β^{TF}).

Obviously after all rescalings $\hbar_* = \hbar\gamma_0^{-3}\gamma_1^{-\frac{5}{2}}\gamma_2^{-2}$ and therefore $\varepsilon = \hbar_*^{\frac{3}{2}} = \hbar^{\frac{3}{2}}(\gamma_0^{-3}\gamma_1^{-\frac{5}{2}}\gamma_2^{-2})^{-\frac{3}{2}}$ where we set $\delta = 0$ but we will show that we have a reserve to set it as $\delta > 0$ if we want to estimate (3.1.34) by $\hbar$ and (3.1.35)–(3.1.37) by $\hbar^2$.

We claim that

$$|w - w_\beta^{\mathsf{TF}}| \le C\varsigma := C\beta\eta\bigl(\gamma_0^4\gamma_1^3\gamma_2^2\varepsilon\bigr)^{\frac{5}{2}}. \tag{3.1.38}$$

and

$$|\nabla(w - w_\beta^{\mathsf{TF}})| \le C\varsigma_1 := C\beta\eta\bigl(\gamma_0^4\gamma_1^3\gamma_2^2\varepsilon\bigr)^{\frac{3}{2}}. \tag{3.1.39}$$

Really it follows from equation (3.1.2).

Then the contribution of α_2-element to (3.1.35) does not exceed $C\varsigma\varepsilon\alpha_2^3$ as measure of zone of α_2-element where $w \ne w_\beta^{\mathsf{TF}}$ is $O(\varepsilon\alpha_2^3)$. One can see easily that $\varsigma\varepsilon = O(\hbar^{\frac{7}{3}})$ and therefore $C\varsigma\varepsilon\alpha_2^3 = O(\hbar^{\frac{7}{3}}\alpha_2^3)$ and the summation over α_2-partition of 1-element returns $O(\hbar^{\frac{7}{3}})$.

Modulo above calculations the contribution of α_2-element to (3.1.34) does not exceed $C\beta\varsigma^{\frac{1}{2}}\varepsilon\alpha_3^2$. One can check easily that $\varsigma\varepsilon = O(\hbar^{\frac{3}{2}}\gamma_2^{-\frac{1}{2}})$ and therefore $C\varsigma^{\frac{1}{2}}\varepsilon\alpha_2^3 = O(\hbar^{\frac{7}{3}}\alpha_1^3\gamma_2^{\frac{5}{2}})$ and the summation over α_2-partition of α_1-element returns $O(\hbar^{\frac{3}{2}}\alpha_1^3)$ and then the summation over α_1-partition of 1-element returns $O(\hbar^{\frac{3}{2}})$.

Similarly, expression (3.1.36) with $\phi = \phi_{\alpha_2}$ does not exceed $C\varsigma\varepsilon_2\alpha_2^5 \le C\hbar^3\alpha_2^4$ and the summation over α_2-partition of 1-element returns $O(\hbar^3)$. However we need to consider disjoint pairs of α_2-elements belonging to given α_1-element and their contribution does not exceed

$$C\hbar^3\int \gamma_{2x}^{-\frac{1}{2}}\gamma_{2y}^{-\frac{1}{2}}|x-y|^{-1}\,dxdy \le C\hbar^2\alpha_1^5$$

and then summation over α_1-partition of 1-element returns $O(\hbar^{\frac{3}{2}})$. We need also to consider disjoint pairs of α_1-elements belonging to given 1-element and their contribution does not exceed $C\hbar^3\int|x-y|^{-1}\,dxdy = O(\hbar^3)$.

Finally, contribution of α_2-element to (3.1.37) does not exceed $C\varsigma_1^2\varepsilon\alpha_2^3$ and one can check easily that this does not exceed $C\alpha_1^3\gamma_2^{\frac{8}{3}}\hbar^{\frac{8}{3}}$ and the summation over α_2-partition of α_1-element returns $C\alpha_1^3\hbar^{\frac{8}{3}}$; then summation over 1-partition of 1-element returns $O(\hbar^{\frac{8}{3}})$.

So, scaled versions of (3.1.34) and (3.1.35)–(3.1.37) do not exceed $C\hbar$ and $C\hbar^2$ respectively. Then original versions of (3.1.34), (3.1.35), (3.1.35), and (3.1.37) do not exceed $C\zeta^3\ell^3 \times (\zeta\ell)^{-1} = C\zeta^2\ell^2$, $C\zeta^5\ell^3 \times (\zeta\ell)^{-2} = C\zeta^3\ell$, $C\zeta^6\ell^5 \times (\zeta\ell)^{-2} = C\zeta^4\ell^3$, and $C\zeta^4\ell \times (\zeta\ell)^{-2} = C\zeta^2\ell^{-1} \le C\zeta^4\ell^3$.

Leaving easy details to the reader we arrive to

Proposition 3.1.9. *(i) Contribution of zone $\mathcal{X}_2$ to the mollification error* (3.1.34) *does not exceed $CZ^{\frac{2}{3}}$;*

(ii) Contribution of zone $\mathcal{X}_2$ to the mollification error (3.1.35) *does not exceed $C\max(Z^{\frac{5}{3}}, B^{\frac{1}{3}}Z^{\frac{4}{3}})$;*

(iii) Contributions of zone $\mathcal{X}_2$ to the mollification errors s(3.1.36) *and* (3.1.36) *do not exceed $CZ^{\frac{5}{3}}$.*

and

Proposition 3.1.10. *Let $B \le Z$. Then*

(i) Contribution of zone $\mathcal{X}_2$ to the mollification error (3.1.34) *does not exceed $CB^{\delta}Z^{\frac{2}{3}-\delta}$;*

(ii) Contributions of zone $\mathcal{X}_2$ to the mollification errors (3.1.36)–(3.1.36) *do not exceed $CB^{\delta}Z^{\frac{5}{3}-\delta}$.*

Remark 3.1.11. Obviously scaled $\varepsilon = \gamma_0\gamma_1\gamma_2\gamma(\hbar/\gamma_0^3\gamma_1^{\frac{5}{2}}\gamma_2^2)^{\frac{2}{3}-\delta} \ge \hbar^{\frac{2}{3}-\delta} \ge (\mu^{-1}\hbar)^{\frac{1}{2}-\delta_1}$ which makes reduction possible.

3.1.7 Zone $\mathcal{X}_3$

Zone $\mathcal{X}_3$ defined by $\mu\hbar \ge C_0$, $\hbar \le 1$, $\{\ell(x) \le \epsilon_0\bar{r}\}$ appears only as $B \ge Z^{\frac{4}{3}}$ here W_B^{TF} is smooth and no mollification is necessary. Further here the canonical form contains only one number $j = 0$ and $|D^\alpha W| \le C_\alpha\zeta^2\ell^{-|\alpha|}$ and $W \asymp \zeta^2$.

Therefore we have non-degeneracy condition fulfilled and applying the standard theory we conclude that in the scaled version contribution of $B(0,1)$ to the semiclassical errors in N- and T-terms and into D-term are $C\mu\hbar^{-1}$, $C\mu$ and $C\mu^2\hbar^{-2}$ respectively.

In the scaled version they become $CB\ell^2$, $CB\ell\zeta \le CBZ^{\frac{1}{2}}\ell^{\frac{1}{2}}$ and $CB^2\ell^3$ and after summation (where for D-term we need to consider mixed contribution of different layers) we arrive to the same expressions calculated as $\ell = \bar{r} = B^{-\frac{2}{5}}Z^{\frac{1}{5}}$ i.e. $CB^{\frac{1}{5}}Z^{\frac{2}{5}}$, $CB^{\frac{4}{5}}Z^{\frac{3}{5}}$ and $CB^{\frac{4}{5}}Z^{\frac{3}{5}}$ respectively. Thus we have proven

Proposition 3.1.12. *Let* $Z^{\frac{4}{3}} \le B \le Z^3$. *Then*

(i) Contribution of zone $\mathcal{X}_3$ *to the* N*-error does not exceed* $CZ^{\frac{2}{5}}B^{\frac{1}{5}}$*;*

(ii) Contributions of zone $\mathcal{X}_3$ *to the* T*-error and* D*-term do not exceed* $CZ^{\frac{3}{5}}B^{\frac{4}{5}}$.

4 Analysis in the boundary strip

4.1 Properties of W_B^{TF} as $N = Z$

To finish our analysis we need to get the same estimates as before in the strip

$$\mathcal{Y} := \{W + \nu \le \epsilon Z\bar{r}^{-1},\, \epsilon\bar{r} \le \ell(x) \le c\bar{r}\} \tag{4.1.1}$$

with $\bar{r} \asymp B^{-\frac{1}{4}}$ for $B \le Z^{\frac{4}{3}}$ [19] and $\bar{r} \asymp B^{-\frac{2}{5}}Z^{\frac{1}{5}}$ for $B \ge Z^{\frac{4}{3}}$. Let us rescale $x \mapsto x' = x\bar{r}^{-1}$. Then

(a) In the case $B \le Z^{\frac{4}{3}}$ we need to rescale then $w(x') = B^{-1}W_B^{\mathsf{TF}}(x'\bar{r})$ and take $\hbar = B^{-\frac{1}{4}} \le 1$, $\mu = B^{\frac{1}{4}} \ge 1$, $\mu\hbar = 1$. [20]. One can treat the case $B \le 1$ exactly as in paper [Ivr10].

(b) On the other hand, for $B \ge N^{\frac{4}{3}}$ one should set $w(x') = \bar{r}N^{-1}W_B^{\mathsf{TF}}(x'\bar{r})$ and $\hbar = (N\bar{r})^{-\frac{1}{2}} = (BN^{-3})^{\frac{1}{5}} \le 1$, $\mu = BN^{-\frac{1}{2}}\bar{r}^{\frac{3}{2}} = B^{\frac{2}{5}}N^{-\frac{1}{5}} \ge 1$, $\mu h = B^{\frac{3}{5}}N^{-\frac{4}{5}} \ge 1$ ($\mu\hbar \asymp 1$ iff $B \le cN^{\frac{4}{3}}$, $\hbar \asymp 1$ iff $B \asymp N^3$).

We will use now only rescaled coordinates unless the opposite is specified. Then in our zone

$$\Delta w = \kappa w_+^{\frac{1}{2}}, \qquad \kappa = 12, w \to \theta = \nu\bar{\zeta}^{-2} \quad \text{as} \quad |x| \to \infty, \tag{4.1.2}$$

with

$$\bar{\zeta}^2 = \begin{cases} B & \text{for} \quad B \le N^{\frac{4}{3}}, \\ N\bar{r}^{-1} = B^{\frac{2}{5}}N^{\frac{3}{5}} & \text{for} \quad B \ge N^{\frac{4}{3}} \end{cases} \tag{4.1.3}$$

where one can always get $\kappa = 12$ after rescaling $w \mapsto 144\kappa^{-2}w$.

---

[19] And right now we consider only case $B \ge (Z - N)^{\frac{4}{3}}$.

[20] We skip bar over w for simplicity here.

Proposition 4.1.1. *Let $Z = N$. Then in $\mathcal{Y}$ after rescaling*

$$|D^\alpha w| \le C_\alpha w \gamma^{-|\alpha|} \qquad \forall \alpha \tag{4.1.4}$$

with the scaling function $\gamma = w^{\frac{1}{4}}$ and

$$|\nabla w^{\frac{1}{4}}| \le 1 + Cw^t \tag{4.1.5}$$

with some constant C and exponent $t > 0$.

Proof. Rescaling $x \mapsto x\bar{r}^{-1}$ we get an equations $(4.1.2)_0 = (4.1.2)$ with $\theta = 0$. We know that $W = 0$ for $\ell(x) \ge c\bar{r}$; so after rescaling $w = 0$ for $\ell(x) \ge c$. On the other hand, $w \asymp 1$ as $\ell(x) \le \epsilon$ (uniformly with respect to all the parameters).

Let us consider solution of the equation

$$\Delta w_s = 12 w_+^s \tag{4.1.6}$$

in $\Omega = \{w \le \epsilon, \ell \le c\}$ with boundary conditions $w_s = w$ at $\partial\Omega$; $s > \frac{1}{2}$.

Note first that $w_s \ge 0$. Really, w_s is the solution of the variational problem to minimize $\|\nabla w\|^2 + 24(s+1)^{-1}\int w_+^{s+1}dx$ and one makes this functional only less replacing w by w_+.

Further, the standard maximum principle arguments show that $w_s \searrow$ as $s \searrow$[21]. Obviously $w_s \searrow w$ and $w_s \to w$ in $\mathscr{C}^\infty$ in $\{x:\ w(x) > 0\}$ as $s \searrow \frac{1}{2}$.

We claim that

$$w_s \in \mathscr{C}^{4s+2}. \tag{4.1.7}$$

To prove (4.1.7) note first that $w \in \mathscr{C}^{2-\delta'}$ uniformly with respect to all the parameters for any $\delta' > 0$. Then $w_s^s \in \mathscr{C}^{s-\delta}$ and then (4.1.6) yields that $w_s \in \mathscr{C}^{2+s-\delta}$ as soon as $s - \delta \notin \mathbb{Z}$. Then since $w_s \ge 0$ we get $|\nabla w_s| \le c w_s^{\frac{1}{2}}$ and so $w_s^s \in \mathscr{C}^{s-\frac{1}{2}}$. Then equation (4.1.6) again yields that $w_s \in \mathscr{C}^{s+\frac{3}{2}}$. Now we need more subtle arguments.

First, for $|y| = 1$

$$w_s(x+ty) = w_s(x) + t(\nabla w_s)_x \cdot y + \frac{1}{2}(\nabla^2 w_s)_x(y)t^2 + O(t^3). \tag{4.1.8}$$

---

[21] If $\Delta w_i = f_i(w_i)$ in Ω, $f_i(w) \nearrow$ as $w \nearrow$ and $f_1(w) \ge f_2(w)$ then $\Delta(w_1 - w_2) > 0$ as $w_1 > w_2$ and then $w_1 - w_2$ does not reach maximum inside Ω.

Then the lowest eigenvalue ς of $\nabla^2 w_s$ at x should be greater than $-Cw_s^{\frac{1}{3}}$. Really, otherwise we can take y as the corresponding eigenvector and t with $|t| = \epsilon\varsigma$ and with a sign making second term non-positive and get $w_s(x+ty) < 0$.

This lower estimate for eigenvalues of $\nabla^2 w_s$ and equation (4.1.6) yield that $|\nabla^2 w_s| \le Cw_s^{\frac{1}{3}}$. But then $|\nabla w_s| \le Cw_s^{\frac{2}{3}}$. Really, otherwise picking $y = |\nabla w_s|^{-1}\nabla w_s$ with $|t| = \epsilon|\nabla w_s|^{\frac{1}{2}}$ and an appropriate sign we would get $w_s(x+ty) < 0$.

These estimates yield that $w_s(x') \asymp w_s(x)$ in $B\bigl(x, \gamma(x)\bigr)$ with $\gamma(x) = \epsilon w_s^{\frac{1}{3}}$. Then $w_s^s \in \mathscr{C}^{\frac{3}{2}}$. In fact, let us consider $f = w_s^{s-1}\nabla w$ and $|f(x) - f(x')|$. Let us consider first $|x - x'| \ge \frac{1}{3}\bigl(\gamma(x) + \gamma(x')\bigr)$; since $|f(x)| \le \gamma(x)^{\frac{1}{2}}$ at each point we get that $|f(x) - f(x') \le |x - x'|^{\frac{1}{2}}$.

On the other hand, for $|x - x'| \le \frac{1}{3}\bigl(\gamma(x) + \gamma(x')\bigr)$ we conclude that $\gamma(x) \asymp \gamma(x')$ and $|f(x) - f(x')| \le |\nabla f| \cdot |x - x'| \le |x - x'|^s$ because $|\nabla f| \le |\nabla^2 w_s| w_s^{s-1} + |\nabla w_s|^2 w_s^{s-2} \le C\gamma^{s-1}$.

Therefore $w_s^s \in \mathscr{C}^{s+1}$ and equation (4.1.6) yields that $w_s \in \mathscr{C}^{3+s}$.

In the next round we assume that $w \in \mathscr{C}^{4+s-\delta}$ with *some* $\delta \in (0,1)$. Then

$$w_s(x+ty) \le w_s(x) + t(\nabla w_s)_x \cdot y + \frac{1}{2}(\nabla^2 w_s)_x(y)t^2 + \frac{1}{6}(\nabla^3 w_s)_x(y)t^3 + C|t|^p \tag{4.1.9}$$

with $p = \min(4, 4+s-\delta)$.

We claim now that the lowest possible eigenvalue ς of $\bigl(\nabla^2 w_s\bigr)_x$ is greater than $-Cw_s^{(p-2)/p}$. Really, otherwise let us pick up y as the corresponding eigenvector, t with $|t| = \epsilon|\varsigma|^{1/(p-2)}$ and with a sign making expression

$$t(\nabla w_s)_x \cdot y + \frac{1}{6}t^3(\nabla^3 w_s)_x(y)$$

non-positive and get $w_s(x+ty) < 0$ again. Now equation (4.1.6) yields that inequality

$$|\nabla^k w_s| \le Cw_s^{(p-k)/p} \tag*{(4.1.10)$_k$}$$

holds with $k = 2$.

Further, we claim that this inequality holds with $k = 1, 3$. Indeed, if one or both of these inequalities are violated then let us take corresponding y and t with

$$|t| = \epsilon\Big(|\nabla w_s|^{1/(p-1)} + |\nabla^3 w_s(y)|^{1/(p-3)}\Big)$$

(calculated on y); replacing ϵ by 2ϵ if necessary we get

$$|t(\nabla w_s)_x \cdot y + \frac{1}{6}t^3(\nabla^3 w_s)_x(y)\big| \ge \epsilon_0|t(\nabla w_s)_x \cdot y| + |\frac{1}{6}t^3(\nabla^3 w_s)_x(y)|$$

and choosing an appropriate sign of t we get $w(x + ty) < 0$.

Therefore inequalities $(4.1.10)_{1-3}$ *hold.* The same arguments as above with $\gamma = w_s^{1/p}$ lead us to $w^s \in \mathscr{C}^{ps}$ and then equation (4.1.6) yields that $w_s \in \mathscr{C}^{ps+2}$. So, now we came back with δ replaced by $\delta' = 2 + s - ps$ and one can see easily that if $\delta > s$ then $\delta' = s + (2 - 4s) + (\delta - s)s$ and after few repeats $\delta < s$. Then we get (4.1.7). Unfortunately, constants depend on s due to the fact that $\Delta w \in \mathscr{C}^2$ fails to yield $w \in \mathscr{C}^4$.

Now we are going to finish the proof of (4.1.4). Let us consider w_s again and let $\gamma = \gamma_{s,\delta} = w_s^{1/(4-\delta)}$. Due to the previous inequalities $\gamma \in \mathscr{C}^1$. We claim that $|\nabla\gamma|$ is bounded uniformly with respect to s, δ. Note first that $\Delta\gamma^{4-\delta} = \gamma^{(4-\delta)s}$ implies that

$$a|\nabla\gamma|^2 + b\gamma\Delta\gamma = \gamma^\sigma \tag{4.1.11}$$

with $a = \frac{1}{12}(4 - \delta)(3 - \delta)$, $b = \frac{1}{12}(4 - \delta)$, and $\sigma = 4s - 2 + (1 - s)\delta$. Let $\psi = |\nabla\gamma|^2$; obviously ψ is uniformly bounded at $\partial\Omega$. Let us consider maximum of ψ reached inside Ω. At the point of maximum

$$\sum_i \gamma_{x_i x_j}\gamma_{x_i} = 0 \tag{4.1.12}$$

and

$$\frac{1}{2}\Delta\psi = \sum_{i,j}\gamma^2_{x_i x_j} + \sum_i \gamma_{x_i}\big(\Delta\gamma\big)_{x_i} = \sum_{i,j}\gamma^2_{x_i x_j} + b^{-1}\sum_i \gamma_{x_i}\Big(\gamma^{-1}\big(\gamma^\sigma - a|\nabla\gamma|^2\big)\Big)_{x_i}$$

due to (4.1.11) and due to (4.1.12) this expression is equal to

$$\sum_{i,j}\gamma^2_{x_i x_j} - b^{-1}\gamma^{-2}|\nabla\gamma|^2\big(\gamma^\sigma - a|\nabla\gamma|^2\big) + b^{-1}\sigma\gamma^{\sigma-2}|\nabla\gamma|^2$$

and therefore at an inner point of minimum $a|\nabla\gamma|^2 \le \gamma^\sigma$. So, $|\nabla\gamma| \le C$ is proven and for $s \searrow \frac{1}{2}$, $\delta \searrow 0$ we get that $|\nabla w^{\frac{1}{4}}| \le C$.

Let us pick $\gamma(x) = \epsilon' w^{\frac{1}{4}}(x)$; then $|\nabla w| \le \frac{1}{2}$ and $w(x) \asymp w(\mathrm{x})$ in $B\bigl(x, \gamma(\mathrm{x})\bigr)$. This and equation (4.1.6) easily yield (4.1.4).

To prove inequality (4.1.5) let us consider w_s again and let us take now $\psi = |\nabla\gamma|^2 - M\gamma^{2t}$ with $t > 0$; obviously ψ is non-positive at $\partial\Omega$ for sufficiently large M. Let us consider maximum of ψ reached inside Ω. At the point of maximum

$$\sum_i \gamma_{x_i x_j}\gamma_{x_i} - Mt\gamma^{2t-2}\gamma_{x_j} = 0 \tag{4.1.12$'$}$$

and the same arguments as before (plus inequality $|\nabla\gamma| \le C_0$) show that at an inner point of maximum $a|\nabla\gamma|^2 \le \gamma^\sigma + CtM\gamma^{2t}$ where C does not depend on M and small $t > 0$. Then at this point $\psi \le 1$ for small enough $t > 0$ and as $s \to \frac{1}{2}$ and $\delta \to 0$ we get (4.1.5). $\square$

The following statement heavily uses estimate (4.1.5):

Proposition 4.1.2. *The following estimate holds*

$$\mathsf{D}(\gamma^{-1+s}, \gamma^{-1+s}) \le Cs^{-2} \tag{4.1.13}$$

with some constant C which does not depend on $s \in (0,1)$ where we set $\gamma^{-1+s} := w_+^{\frac{1}{4}(-1+s)}$ (i.e. it is 0 as $w \le 0$).

Proof. As in the notations of the proof of proposition 4.1.1 $\delta = 0$ and $s = \frac{1}{2}$ we have (4.1.11) with $a = 1$, $b = 3$ and $\sigma = 0$:

$$\frac{1}{3}\gamma\Delta\gamma + |\nabla\gamma|^2 = 1. \tag{4.1.14}$$

Then

$$\gamma^{-1+s} = \gamma^{-1+s}|\nabla\gamma|^2 + \frac{1}{3}\gamma^s\Delta\gamma = (1 - \frac{s}{3})\gamma^{-1+s}|\nabla\gamma|^2 + \frac{1}{3(1+s)}\Delta\gamma^{1+s}$$

and

$$\begin{aligned}\mathsf{D}(\gamma^{-1+s}, \gamma^{-1+s}) \le (1 - \frac{s}{3})\mathsf{D}(\gamma^{-1+s}|\nabla\gamma|^2, \gamma^{-1+s}) + C \le \\ (1 - \frac{s}{3})\mathsf{D}(\gamma^{-1+s}, \gamma^{-1+s}) + C\mathsf{D}(\gamma^{-1+t+s}, \gamma^{-1+s}) + C\end{aligned}$$

due to (4.1.5) and this yields

$$\mathsf{D}(\gamma^{-1+s},\gamma^{-1+s}) \le Cs^{-2}\mathsf{D}(\gamma^{-1+t+s},\gamma^{-1+t+s}) + Cs^{-1}.$$

Substituting $s+mt$ instead of s, $0\le m\le Ct^{-1}$ we recover (4.1.13). $\square$

4.2 Semiclassical analysis in the strip $\mathcal{Y}$ for $N\ge Z$.

It is really easy to construct the proper potential in this case: we just take

$$w_\varepsilon = w\phi_\varepsilon,\qquad \phi_\varepsilon = f(w\varepsilon^{-4}) \tag{4.2.1}$$

with $f\in\mathscr{C}_0^\infty((\frac{1}{2},\infty))$, $0\le f\le 1$, $f(t)=1$ for $t>1$. Note that due to (4.1.2)

$$\begin{aligned}
&\mathsf{D}(\gamma^{-1}\phi_\varepsilon,\gamma^{-1}\varphi_\varepsilon) \le C\varepsilon^{-2s}\mathsf{D}(\gamma^{s-1},\gamma^{s-1}) \le Cs^{-2}\varepsilon^{-2s},\\
&\mathsf{D}(1-\phi_\varepsilon,1-\phi_\varepsilon) \le C\varepsilon^{2-2s}\mathsf{D}(\gamma^{s-1},\gamma^{s-1}) \le Cs^{-2}\varepsilon^{2-2s};
\end{aligned}$$

then minimizing with respect to s ($=|\log\varepsilon|^{-1}$) the right-hand expression we conclude that

$$\mathsf{D}\bigl(\gamma^{-1}\phi_\varepsilon,\gamma^{-1}\varphi_\varepsilon\bigr)+\varepsilon^{-2}\mathsf{D}\bigl(1-\phi_\varepsilon,1-\phi_\varepsilon\bigr)\le C\bigl(1+|\log\varepsilon|\bigr)^2 \tag{4.2.2}$$

and therefore

$$\int\gamma^{-1}\phi_\varepsilon\,dx+\varepsilon^{-1}\int(1-\phi_\varepsilon)\,dx\le C\bigl(1+|\log\varepsilon|\bigr). \tag{4.2.3}$$

Remark 4.2.1. (i) Recall that all these integrals are taken over domain $\{x:\ w(x)>0\}$. To avoid possible troubles we pick $\varepsilon=\hbar^{\frac{1}{3}}$ and set in the zone $\{x:\ w(x)\le C_0\hbar^{\frac{4}{3}}\}$

$$\gamma(x)=\mathsf{dist}(x,\{w\ge 2C_0\hbar^{\frac{4}{3}}\}),$$

$$w_\varepsilon=\left\{\begin{aligned}-\gamma^4\phi'_\varepsilon\quad&\text{for}\quad\gamma\le\varepsilon,\\ -\varepsilon^4\quad&\text{for}\quad\gamma\ge\varepsilon\end{aligned}\right. \tag*{(4.2.1)$'$}$$

with $\phi'_\varepsilon=f(\gamma\varepsilon^{-1})$ and then in the complemental domain $\{x:\ w(x)\le-\varsigma^2\}$ with $\varsigma=\varepsilon^2$ and $\varsigma\gamma=\gamma^3\ge\hbar$;

(ii) Further, for $\varepsilon = \hbar^{\frac{1}{3}-\delta}$ with sufficiently small exponent $\delta > 0$ it does not break estimate for mollification error in T-term;

(iii) Furthermore, for $t > \varepsilon$

$$\operatorname{mes}(\{x:\ \gamma(x) \le t\}) \le Ct^3\varepsilon^{-3}\operatorname{mes}(\{x:\ \gamma(x) \le \epsilon\varepsilon\})$$

and therefore

$$\hbar^m \int \gamma^{-1-m}\varsigma^{-m}\,dx \le C\varepsilon^{-1}\operatorname{mes}(\{x:\ \gamma(x) \le \epsilon\varepsilon\}) \le CL := C(1+|\log\hbar|)$$

for sufficiently large m.

Using these estimates and the last remark we can prove easily

Proposition 4.2.2. *Let $N \ge Z$. Then*

(i) Contribution of $\mathcal{Y}\cup\mathcal{X}_4$ with $\mathcal{X}_4 := \{x:\ w(x)=0\}$ to mollification and approximation errors in N-term do not exceed $CT_0\varepsilon^3(1+|\log\varepsilon|)$ and $R_0(1+|\log\varepsilon|)$ respectively with

$$T_0 = B^{\frac{3}{4}},\qquad R_0 = B^{\frac{1}{2}},\qquad T = B^{\frac{7}{4}},\qquad R = B^{\frac{5}{4}}\qquad \text{as}\ \ B \le Z^{\frac{4}{3}} \tag*{(4.2.4)$_1$}$$

and

$$T_0 = Z,\qquad R_0 = B^{\frac{1}{5}}Z^{\frac{2}{5}},\qquad T = Z^{\frac{9}{5}}B^{\frac{2}{5}},\qquad R = Z^{\frac{3}{5}}B^{\frac{4}{5}}\qquad \text{as}\ \ Z^{\frac{4}{3}} \le B \le Z^3; \tag*{(4.2.4)$_2$}$$

(ii) Contribution of $\mathcal{Y}\cup\mathcal{X}_4$ to mollification and semiclassical D-terms do not exceed $CT\varepsilon^6(1+|\log\varepsilon|)$ and $R(1+|\log\varepsilon|)^2$ respectively;

(iii) Contribution of $\mathcal{Y}\cup\mathcal{X}_4$ to both mollification and approximation errors in T-term do not exceed $CT\varepsilon^6(1+|\log\varepsilon|)$ and CR respectively.

Proof. Really, estimates for mollification errors and terms immediately follow from the inequality

$$\operatorname{mes}(\{x:\ w(x) \le \varepsilon^4\}) \le C\varepsilon(1+|\log\varepsilon|) \tag{4.2.5}$$

which is due to (4.2.3).

Let us consider semiclassical errors and terms.

(i) Let us consider N-term first. The contribution of each ball $B\bigl(x,\gamma(x)\bigr)$ to the semiclassical error does not exceed $C\mu\hbar^{-1}\gamma^2 \asymp CB\bar{r}^2$ and the total contribution does not exceed $CR_0\int \gamma^{-1}\,dx \le CR_0\bigl(1+|\log\varepsilon|\bigr)$ where $R_0=B\bar{r}^2$;

(ii) Consider semiclassical D-term. The similar arguments with the analysis of disjoint balls of different types and with analysis of the intersecting balls (of the same type) lead us to the proper estimate of the contribution of $\mathcal{Y}_4\cup\mathcal{X}_4$ to semiclassical D-term: namely, it does not exceed $CR_0^2\bar{r}^{-1}\bigl(1+|\log\varepsilon|\bigr)^2$ where $R_0^2\bar{r}^{-1}\asymp R$;

(iii) Consider T-term. Let us consider all possible balls and their contributions. The contribution of each ball $B\bigl(x,\gamma(x)\bigr)$ to the semiclassical error does not exceed $C\bar{\zeta}^2\mu\varsigma^2\gamma \asymp CB\bar{\zeta}^2\bar{r}\varsigma\gamma^2$ and the total contribution does not exceed $CR\int\varsigma\gamma^{-2}\,dx\asymp CR$ where $R=B\bar{\zeta}^2\bar{r}$. ☐

Then picking appropriate $\varepsilon=\hbar^{\frac{1}{3}}$ we conclude that

Corollary 4.2.3. *Let $N\ge Z$. Then*

(i) Contributions of $\mathcal{Y}\cup\mathcal{X}_4$ to all errors in N-terms do not exceed CR_0L with $L=(1+|\log BZ^{-3}|)$;

(ii) Contribution of $\mathcal{Y}\cup\mathcal{X}_4$ to all D-terms do not exceed CRL^2;

(iii) Contribution of $\mathcal{Y}\cup\mathcal{X}_4$ to all errors in T-terms do not exceed CRL.

Remark 4.2.4. Could we get rid off logarithmic factors i.e. make $L=1$ as in the case $M=1$?

(i) With the mollification errors we need to replace (4.2.5) by

$$\operatorname{mes}(\{x:\ w(x)\le\varepsilon^4\})\le C\varepsilon; \tag{4.2.6}$$

(ii) With the semiclassical terms our arguments here are insufficient even if we established (4.2.6); we need extra propagation arguments in the direction of decaying w along magnetic lines–exactly as in the case $M=1$. Surely there could be points where such arguments do not work; f.e. consider $M=2$ and nuclei so that $|\mathsf{y}_1-\mathsf{y}_2|$ is slightly less than $\bar{r}_1+\bar{r}_2$ where $\bar{r}_{1,2}$ are precise radii of support. Then w reaches its minimum at $\mathcal{Y}$.

So, we need to prove that the measure of such points is sufficiently small (f.e. less than $C|\log BZ^{-3}|^{-1}$).

Unfortunately I do not know how to make the above remark work and I suggest

Problem 4.2.5. *Follow through the discussed plan. For* $M = 2$ *it could be easier due to rotational symmetry.*

4.3 Analysis in the boundary strip $\mathcal{Y}$. Construction of the potential for $N < Z$

Now let us consider $N < Z$.

Remark 4.3.1. (i) We can replace $\nu < 0$ by $\nu = 0$ in the definition of W_B^{TF} without increasing any of errors as long as $B\bar{r}^3\nu^{\frac{1}{2}}(\nu/\bar{\zeta}^2)^{\frac{1}{4}} \le R_0$;

(ii) As $B \le Z^{\frac{4}{3}}$ this translates into $|\nu|^{\frac{3}{4}} \le Z^{\frac{2}{3}}$ i.e. $(Z-N)_+ \le Z^{\frac{8}{9}}B^{-\frac{1}{4}}$;

(iii) On the other hand, as $Z^{\frac{4}{3}} \le B \le Z^3$ it translates into $|\nu|^{\frac{3}{4}} \le B^{1/2}$ i.e. $(Z-N)_+ \le Z^{\frac{1}{5}}B^{\frac{4}{15}}$.

Therefore let us consider a case not covered by remark 4.3.1. Let $\theta = \nu\bar{\zeta}^{-2} < 0$.

Proposition 4.3.2. *Consider dependence of* $W_B^{\mathsf{TF}} = W_{B(\nu)}^{\mathsf{TF}}(x)$ *on* ν. *Then*

(i) $W_{B(\nu)}^{\mathsf{TF}}(x) + \nu$ *is non-decreasing with respect to* ν *at each point* x*;*

(ii) $W_{B(\nu)}^{\mathsf{TF}}(x)$ *is non-increasing with respect to* ν *at each point* x*;*

(iii) In particular $W_{B(\nu)}^{\mathsf{TF}}(x) + \nu \nearrow W_{B(0)}^{\mathsf{TF}}(x)$ *and* $W_{B(\nu)}^{\mathsf{TF}}(x) \searrow W_{B(0)}^{\mathsf{TF}}(x)$ *at each point* x *as* $\nu \to -0$.

Proof. (i) Consider $W_j = W_{B(\nu_j)}^{\mathsf{TF}} + \nu_j$, $0 > \nu_1 > \nu_2$. One can prove easily that $W_B^{\mathsf{TF}} - V$ is a continuous function and since $W_1 - W_2 = \nu_1 - \nu_2 > 0$ as $\ell(x) \ge C$ we conclude that $W_1 \ge W_2$ at each point x (which is exactly our statement (i)) unless $W_1 - W_2$ achieves a negative minimum at some point x^*.

(a) Let $x^* \ne \mathsf{y}_m$; then $\Delta(W_1 - W_2)(x^*) = P_B'(W_1) - P_B'(W_2) \le 0$ because $W_1 < W_2$ at x^* and x^* cannot be such point.

(b) Let $x^* = \mathsf{y}_m$. From Thomas-Fermi equations for $W_{1,2}$ one can prove easily that

$$\begin{aligned}(W_1 - W_2)(x) = (W_1 - W_2)(\mathsf{y}_m) + L_m(x - \mathsf{y}_m) + \\ \kappa_m |x - y_m|^{\frac{3}{2}} (W_1 - W_2)(\mathsf{y}_m) + O(|x - \mathsf{y}_m|^2)\end{aligned}$$

near y_m where $L_m(x)$ is a linear function and $\kappa_m > 0$ and therefore if $(W_1 - W_2)(\mathsf{y}_m) < 0$, y_m cannot be a minimum point either.

(ii) So, $W_1 \geq W_2$ and therefore $W_1 - W_2$ is a subharmonic function and since $W_1 - W_2 = \nu_1 - \nu_2$ as $\ell(x) \geq C$ we conclude that $W_1 - W_2 \leq \nu_1 - \nu_2$ i.e. $W^{\mathsf{TF}}_{B,(\nu_1)} \leq W^{\mathsf{TF}}_{B,(\nu_2)}$ at each point.

(iii) Statement (iii) follows from (i), (ii). $\square$

Therefore in the zone $\{x :\ W^{\mathsf{TF}}_{B(\nu)} \geq (1 + \epsilon)|\nu|$ we can apply the same (γ, ς) scaling with $\varsigma = \gamma^2$ defined for $\nu = 0$. Really, we know that there $W^{\mathsf{TF}}_{B(\nu)} + \nu \asymp W^{\mathsf{TF}}_{B(0)} \asymp \varsigma^2$ and $\varsigma = \gamma^2$.

Then Thomas-Fermi equation (1.1.3) implies that

$$|\nabla^\alpha W^{\mathsf{TF}}_{B(\nu)}| \leq C_\alpha \varsigma^2 \gamma^{-|\alpha|} \qquad \forall \alpha$$

and then we arrive to the statement (i) in Proposition 4.3.3 below.

On the other hand, in the zone $\{x :\ W^{\mathsf{TF}}_{B(\nu)} \leq (1 - \epsilon)|\nu|\}$ we can apply the same arguments but this zone is classically forbidden and we arrive to to the statement (ii) below. In both cases $\varsigma\gamma \geq \hbar$ (where in the latter case γ is the distance from x to $W^{\mathsf{TF}}_{B(\nu)}$ (scaled) and $\varsigma = |\nu|^{\frac{1}{2}}$ because in both cases $\varsigma \geq |\nu|^{\frac{1}{2}} \bar{\zeta}^{-1}$ and $\gamma \geq \nu^{\frac{1}{4}} \bar{\zeta}^{-\frac{1}{2}}$ so $\varsigma\gamma \geq |\nu|^{\frac{3}{4}} \bar{\zeta}^{-\frac{3}{2}}$.

Then we arrive to the statement (ii) in Proposition 4.3.3 below.

Proposition 4.3.3. *Let* *either* $B \leq Z^{\frac{4}{3}}$ *and* $|\nu|^{\frac{3}{4}} \geq Z^{\frac{2}{3}}$ *or* $Z^{\frac{4}{3}} \leq B \leq Z^3$ *and* $|\nu|^{\frac{3}{4}} \geq B^{\frac{1}{2}}$. *Then*

(i) Contributions of zone $\{x :\ W^{\mathsf{TF}}_B(x) \geq (1 + \epsilon_0)|\nu|\}$ *to semiclassical errors in* N- *and* T-*terms and into semiclassical* D-*term do not exceed* $CR_0 L$, CR *and* CRL^2 *respectively;*

(ii) Contributions of zone $\{x :\ W^{\mathsf{TF}}_B(x) \leq (1 - \epsilon_0)|\nu|\}$ *to semiclassical errors in* N- *and* T-*terms and into semiclassical* D-*term do not exceed* $CR_0 L$, CR *and* CRL^2 *respectively.*

Remark 4.3.4. (i) Here actually $L = 1 + |\log |\nu|\bar{\zeta}^{-2}|$ with

$$|\nu|\bar{\zeta}^{-2} \asymp \begin{cases} (Z-N)_+B^{-\frac{3}{4}} & \text{as } (Z-N)_+^{\frac{4}{3}} \le B \le Z^{\frac{4}{3}}, \\ (Z-N)_+Z^{-1} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3; \end{cases} \tag{4.3.1}$$

(ii) To have a proper estimate for D-term with ν replaced by 0 we need

$$(Z-N)_+ \lesssim \begin{cases} Z^{\frac{10}{3}}B^{-\frac{25}{12}} & \text{as } B \le Z^{\frac{4}{3}}, \\ Z^{\frac{1}{5}}B^{\frac{4}{15}} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3 \end{cases} \tag{4.3.2}$$

with the right-hand expression larger than $Z^{\frac{5}{9}}$.

Therefore we need to explore zone

$$\mathcal{Y}^* := \{x :\ (1-\epsilon_0)|\nu| \le W_B^{\mathsf{TF}}(x) \le (1+\epsilon_0)|\nu|\}$$

in the framework of Proposition 4.3.3.

We assume that $\hbar_* \lesssim 1$ where $\theta = |\nu|\bar{\zeta}^{-2}$ and

$$\hbar_* = \hbar\theta^{-\frac{3}{4}} \asymp \begin{cases} (Z-N)_+^{-\frac{3}{4}}B^{\frac{5}{16}} & \text{as } B \le Z^{\frac{4}{3}}, \\ (Z-N)_+^{-\frac{3}{4}}Z^{\frac{3}{20}}B^{\frac{1}{5}} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3. \end{cases} \tag{4.3.3}$$

This assumption is a bit more relaxed that the negation of the framework of Proposition 4.3.2.

Proposition 4.3.5. *Assume that condition* (4.3.2) *is violated. Let us consider* $B(x,\eta)$ *(after rescaling* $x \mapsto x\bar{r}^{-1}$*) with* $\eta := \theta^{\frac{1}{4}}$ *and scale it* $x \mapsto x\eta^{-1}$*. Let* $w := (W_B^{\mathsf{TF}} + \nu)|\nu|^{-1}$.

Then there exist scaling functions $\alpha_0, \dots, \alpha_3$ *with* $\alpha_j = \gamma_j\alpha_{j-1}$, $\alpha_0 = \gamma_0\theta^{-\frac{1}{4}}$ *such that*

$$|\nabla^\alpha w| \le C_\alpha \varsigma_0^2 \gamma_0^{-|\alpha|} \qquad \textit{with } \varsigma_0 = \gamma_0^2, \tag{4.3.4$_0$}$$

$$\gamma_0 \asymp \max\bigl(|w|^{\frac{1}{4}},\ |\nabla w|^{\frac{1}{3}},\ |\nabla^2 w|^{\frac{1}{2}},\ |\nabla^3 w|\bigr), \tag{4.3.5$_0$}$$

and after rescaling $x \mapsto x\gamma_0^{-1}$, $w \mapsto w_0 = w\varsigma_0^{-2}$

$$|\nabla^\alpha w_0| \le C_\alpha \varsigma_1^2 \gamma_1^{-|\alpha|} \qquad \textit{with } \varsigma_1 = \gamma_1^{\frac{3}{2}}, \tag{4.3.4$_1$}$$

$$\gamma_1 \asymp \max\bigl(|w_0|^{\frac{1}{3}},\ |\nabla w_0|^{\frac{1}{2}},\ |\nabla^2 w_0|\bigr), \tag{4.3.5$_1$}$$

and after rescaling $x\mapsto x\gamma_1^{-1}$, $w_0\mapsto w_1=w_0\varsigma_1^{-2}$

$$|\nabla^\alpha w_1|\le C_\alpha \varsigma_2^2\gamma_2^{-|\alpha|}\qquad \textit{with}\ \ \varsigma_2=\gamma_2, \tag{4.3.4$_2$}$$

$$\gamma_2\asymp \max\bigl(|w|^{\frac{1}{2}},\,|\nabla w|\bigr), \tag{4.3.5$_2$}$$

and after rescaling $x\mapsto x\gamma_2^{-1}$, $w_1\mapsto w_2=w_1\varsigma_2^{-2}$

$$|\nabla^\alpha w_2|\le C_\alpha \varsigma_3^2\gamma_3^{-|\alpha|}\qquad \textit{with}\ \ \varsigma_3=\gamma_3^{-\frac{1}{2}}, \tag{4.3.4$_3$}$$

$$\gamma_3\asymp |w|. \tag{4.3.5$_3$}$$

Proof. Proof easily follows the arguments of the beginning of the Section 3; easy details we leave to the reader. □

If needed we replace γ_j by $\max(\gamma_j,\bar\gamma_j)$ with

$$\bar\gamma_1:=C\hbar_{*,0}^{\frac{2}{5}},\qquad \bar\gamma_2:=C\hbar_{*,1}^{\frac{1}{2}},\qquad \bar\gamma_3:=C\hbar_{*,2}^{\frac{2}{3}},$$
$$\hbar_{*,0}:=\hbar_*\gamma_0^{-3},\qquad \hbar_{*1}:=\hbar_{*,0}\gamma_1^{-\frac{2}{5}},\qquad \hbar_{*,2}:=\hbar_{*,1}\gamma_2^{-\frac{1}{2}}.$$

We leave to the reader to prove by our standard methods

Proposition 4.3.6. *In the framework of Proposition 4.3.5 consider* α_i*-element* ϕ_i*. Then one can apply* ε_i*-mollification to potential* w_i *with* $\varepsilon_i=\bar\gamma_i$ *as* $\gamma_i\le 2\bar\gamma_i$ *such that*

(i) Contributions of ϕ_i*-element to both semiclassical and approximation errors in* N*-term do not exceed* $CB\bar r^2\alpha_3^2$, $CB\bar r^2\alpha_2^2$, $CB\bar r^2\alpha_1^2(1+|\log\hbar_{*,1}|)$ *and* $CB\bar r^2\alpha_0^2(1+|\log\hbar_{*,1}|)^2$ *for* $i=3,2,1,0$ *respectively;*

(ii) Contribution of (ϕ_i,ϕ_i)*-pair to both semiclassical and approximation in* D*-terms do not exceed* $CB^2\bar r^3\alpha_3^3$, $CB^2\bar r^3\alpha_2^3$, $CB^2\bar r^3\alpha_1^3(1+|\log\hbar_{*,1}|)^2$, *and* $CB^2\bar r^3\alpha_0^3(1+|\log\hbar_{*,1}|)^4$ *for* $i=3,2,1,0$ *respectively;*

(iii) Contributions of ϕ_i*-element to both semiclassical and approximation errors in* T*-term do not exceed* $CB^2\bar r^3\alpha_3^2$, $CB^2\bar r^2\alpha_2^3$, $CB^2\bar r^3\alpha_1^3$, *and* $CB^2\bar r^2\alpha_0^3$ *for* $i=3,2,1,0$ *respectively.*

Then we arrive to

Proposition 4.3.7. *In the framework of Proposition 4.3.5 there exists potential* W_ε *such that*

(i) Contributions of $\mathcal{Y}^*$ *to both semiclassical and approximation errors for* N*-term do not exceed* $CB\bar{r}^2J$ *with*

$$J := \int_{\bar{r}^{-1}\mathcal{Y}^*} \alpha_0(x)^{-1}(1+|\log \hbar_*\gamma_0^{-3}|)^2\, dx \tag{4.3.6}$$

where $\bar{r}^{-1}\mathcal{Y}^*$ *is* $\mathcal{Y}^*$ *scaled by* $x\mapsto x\bar{r}^{-1}$*) and* $\alpha_0=\gamma_0\theta^{-\frac{1}{4}}$*;*

(ii) Contributions of $\mathcal{Y}^*\times\mathcal{Y}^*$ *to both semiclassical and approximation* D*-terms do not exceed* $CB^2\bar{r}^3J^2$*;*

(iii) Contributions of $\mathcal{Y}^*$ *to both semiclassical and approximation errors for* T*-term do not exceed* $CB^2\bar{r}^3$*.*

Remark 4.3.8. Obviously

$$J\le CL\max\int_{B(0,1)} \gamma_0(x)^{-1}(1+|\log \hbar_*\gamma_0^{-3}|)^2\, dx \tag{4.3.7}$$

where this integral is taken over $B(x, C_0\theta^{\frac{1}{3}})$ rescaled which in turn does not exceed $C\hbar_*^{-\frac{1}{3}}$ which translates into

$$(Z-N)_+^{\frac{1}{4}}L\times\begin{cases} B^{\frac{19}{48}} & \text{as}\ \ B\le Z^{\frac{4}{3}},\\ Z^{\frac{7}{20}}B^{\frac{2}{15}} & \text{as}\ \ Z^{\frac{4}{3}}\le B\le Z^3\end{cases} \tag{4.3.8}$$

for errors in N-term and

$$(Z-N)_+^{\frac{1}{2}}L^2\times\begin{cases} B^{\frac{25}{24}} & \text{as}\ \ B\le Z^{\frac{4}{3}},\\ Z^{\frac{1}{2}}B^{\frac{2}{3}} & \text{as}\ \ Z^{\frac{4}{3}}\le B\le Z^3\end{cases} \tag{4.3.9}$$

for D-terms.

4.4 Conclusion

Let us recall that

$$\theta = \begin{cases} 1 & \text{as } B \le (Z-N)_+^{\frac{4}{3}}, \\ (Z-N)_+ B^{-\frac{3}{4}} & \text{as } (Z-N)_+^{\frac{4}{3}} \le B \le Z^{\frac{4}{3}}, \\ (Z-N)_+ Z^{-1} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3, \end{cases} \tag{4.4.1}$$

$$\hbar = \begin{cases} (Z-N)_+^{-\frac{1}{3}} & \text{as } B \le (Z-N)_+^{\frac{4}{3}}, \\ B^{-\frac{1}{4}} & \text{as } B \le Z^{\frac{4}{3}}, \\ B^{\frac{1}{5}} Z^{-\frac{3}{5}} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3 \end{cases} \tag{4.4.2}$$

and therefore

$$\hbar_* = \hbar\theta^{-\frac{3}{4}} = \begin{cases} (Z-N)_+^{-\frac{1}{3}} & \text{as } B \le (Z-N)_+^{\frac{4}{3}}, \\ (Z-N)_+^{-\frac{3}{4}} B^{\frac{5}{16}} & \text{as } (Z-N)_+^{\frac{4}{3}} \le B \le Z^{\frac{4}{3}}, \\ (Z-N)_+^{-\frac{3}{4}} Z^{\frac{3}{20}} B^{\frac{1}{5}} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3 \end{cases} \tag{4.4.3}$$

and

$$R_0 \hbar_*^{-\frac{1}{3}} = \begin{cases} (Z-N)_+^{-\frac{5}{9}} B & \text{as } B \le (Z-N)_+^{\frac{4}{3}}, \\ (Z-N)_+^{\frac{1}{4}} B^{\frac{19}{48}} & \text{as } (Z-N)_+^{\frac{4}{3}} \le B \le Z^{\frac{4}{3}}, \\ (Z-N)_+^{\frac{1}{4}} Z^{\frac{7}{20}} B^{\frac{2}{15}} & \text{as } Z^{\frac{4}{3}} \le B \le Z^3 \end{cases} \tag{4.4.4}$$

Therefore we arrive

Proposition 4.4.1. *Let $M \ge 2$. Then for a constructed above potential W*

(i) Total semiclassical and approximation errors in N-term do not exceed

$$C \begin{cases} Z^{\frac{2}{3}} + (Z-N)_+^{-\frac{5}{9}} B & \textit{as } B \le (Z-N)_+^{\frac{4}{3}}, \\ Z^{\frac{2}{3}} + B^{\frac{1}{2}} L + (Z-N)_+^{\frac{1}{4}} B^{\frac{19}{48}} L^* & \textit{as } (Z-N)_+^{\frac{4}{3}} \le B \le Z^{\frac{4}{3}}, \\ Z^{\frac{2}{5}} B^{\frac{1}{5}} L + (Z-N)_+^{\frac{1}{4}} Z^{\frac{7}{20}} B^{\frac{2}{15}} L^* & \textit{as } Z^{\frac{4}{3}} \le B \le Z^3 \end{cases} \tag{4.4.5}$$

where here and below $L = (1 + |\log BZ^3|)$, $L^* = (1 + |\log \theta|)$;

(ii) Both semiclassical and approximation D*-terms do not exceed*

$$C\begin{cases} Z^{\frac{5}{3}} + B^2(Z-N)_+^{-\frac{7}{9}} & \text{as}\ \ B\le (Z-N)_+^{\frac{4}{3}},\\ Z^{\frac{5}{3}} + B^{\frac{5}{4}}L + (Z-N)_+^{\frac{1}{2}}B^{\frac{25}{24}}L^* & \text{as}\ \ (Z-N)_+^{\frac{4}{3}}\le B\le Z^{\frac{4}{3}},\\ Z^{\frac{3}{5}}B^{\frac{4}{5}}L + (Z-N)_+^{\frac{1}{2}}Z^{\frac{1}{2}}B^{\frac{2}{3}}L^* & \text{as}\ \ Z^{\frac{4}{3}}\le B\le Z^3; \end{cases} \tag{4.4.6}$$

(iii) Total approximation error in T*-term does not exceed*

$$C\begin{cases} Z^{\frac{5}{3}} + B^{\frac{5}{4}}L & \text{as}\ \ B\le Z^{\frac{4}{3}},\\ Z^{\frac{3}{5}}B^{\frac{4}{5}}L & \text{as}\ \ Z^{\frac{4}{3}}\le B\le Z^3; \end{cases} \tag{4.4.7}$$

(iv) Total semiclassical error in T*-term does not exceed*

$$\text{(4.4.7)} + CZ^{\frac{4}{3}}B^{\frac{1}{3}} + CZ^{\frac{3}{2}}a^{-\frac{1}{2}} \tag{4.4.8}$$

provided $a\ge Z^{-1}$*; as* $a\le Z^{-1}$ *the last term should be replaced by* CZ^2*.*

Also we arrive to

Proposition 4.4.2. *Let* $M\ge 2$*,* $B\le Z$ *and* $a\ge Z^{-\frac{1}{3}}$*. Then for a constructed above potential* W

(i) Total semiclassical and approximation errors in N*-term do not exceed* $CZ^{\frac{2}{3}}\bigl((BZ^{-1})^{\delta} + (aZ^{\frac{1}{3}})^{-\delta} + Z^{-\delta}\bigr)$*;*

(ii) Both semiclassical and approximation D*-terms do not exceed and total semiclassical and approximation errors in* T*-term do not exceed* $CZ^{\frac{5}{3}}\bigl((BZ^{-1})^{\delta} + (aZ^{\frac{1}{3}})^{-\delta} + Z^{-\delta}\bigr)$*.*

5 Ground state energy

5.1 Lower estimates

Now the lower estimates are already proven: in virtue of the analysis given in the previous paper [Ivr10] these estimates are due to estimates of the semiclassical and approximation errors for $\lambda=\nu$:

Proposition 5.1.1. *Let* $M = 1$. *Then*

(i) The following estimate holds:

$$E^{\mathsf{TF}} \geq \mathcal{E}^{\mathsf{TF}} + \Bigl(\operatorname{Tr}(H_{A,W} - \nu)^{-} + \int P_B(W^{\mathsf{TF}} + \nu)\,dx\Bigr) - CR; \tag{5.1.1}$$

(ii) The following estimate holds:

$$E^{\mathsf{TF}} \geq \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} - CR - CZ^{\frac{4}{3}}B^{\frac{1}{3}}; \tag{5.1.2}$$

(iii) As $B \leq Z$

$$E^{\mathsf{TF}} \geq \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} + \mathsf{Dirac} + \mathsf{Schwinger} - CZ^{\frac{5}{3}}\bigl(Z^{-\delta} + (BZ^{-1})^{\delta}\bigr). \tag{5.1.3}$$

Proposition 5.1.2. *Let* $M \geq 2$. *Then*

(i) The following estimate holds:

$$E^{\mathsf{TF}} \geq \mathcal{E}^{\mathsf{TF}} + \Bigl(\operatorname{Tr}(H_{A,W} - \nu)^{-} + \int P_B(W^{\mathsf{TF}} + \nu)\,dx\Bigr) - (4.4.7); \tag{5.1.4}$$

(ii) The following estimate holds:

$$E^{\mathsf{TF}} \geq \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} - (4.4.7) - CZ^{\frac{4}{3}}B^{\frac{1}{3}} - CZ^{\frac{3}{2}}a^{-\frac{1}{2}} \tag{5.1.5}$$

provided $a \geq Z^{-1}$ *and* $B \leq Z^2$; *otherwise we can skip* Scott *and replace the last term by* CZ^2;

(iii) As $B \leq Z$ *and* $a \geq Z^{-\frac{1}{3}}$

$$E^{\mathsf{TF}} \geq \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} + \mathsf{Dirac} + \mathsf{Schwinger} - \\ CZ^{\frac{5}{3}}\bigl(Z^{-\delta} + (BZ^{-1})^{\delta} + (aZ^{\frac{1}{3}})^{-\delta}\bigr). \tag{5.1.6}$$

5.2 Upper estimate: general scheme

Recall that for an upper estimate in addition to the trace we need to estimate also $|\lambda_N-\nu|$ and

$$|\lambda_N-\nu|\cdot|\mathsf{N}(H_{A,W})-N| \tag{5.2.1}$$

and also three D-terms: two of them semiclassical

$$\mathsf{D}\Bigl(e(x,x,\lambda)-P'_B(W(x)+\lambda),\, e(x,x,\lambda)-P'_B(W(x)+\lambda)\Bigr) \tag*{(5.2.2)$_{1,2}$}$$

with $\lambda=\nu$ and $\lambda=\lambda_N$ and also

$$\mathsf{D}\Bigl(P'_B(W(x)+\lambda_N)-P'_B(W(x)+\nu),\, P'_B(W(x)+\lambda_N)-P'_B(W(x)+\nu)\Bigr); \tag{5.2.3}$$

our tool will be semiclassical estimates for two semiclassical N-terms

$$\int\Bigl(e(x,x,\lambda)-P'_B(W(x)+\lambda)\Bigr)\,dx \tag*{(5.2.4)$_{1,2}$}$$

with $\lambda=\nu$ and $\lambda=\lambda_N$ and also estimate from below for the third N-term

$$|\int\Bigl(P'_B(W(x)+\lambda_N)-P'_B(W(x)+\nu)\Bigr)\,dx. \tag{5.2.5}$$

5.3 Upper estimate as $M=1$

5.3.1 Estimate for $|\lambda_N-\nu|$

As in paper [Ivr10] we have two cases: in the first case ν is small enough so we construct W^{TF} with $\nu=0$ and estimate $|\lambda_N|$ and in the second case ν we prove that $\lambda_N\asymp\nu$ and estimate $|\lambda_N-\nu|$.

Proposition 5.3.1. *Let $M=1$, $B\le Z^3$.*

(i) Let

$$(Z-N)_+\le K:=C_0\max\bigl(Z^{\frac{2}{3}},\, Z^{\frac{2}{5}}B^{\frac{1}{5}}\bigr). \tag{5.3.1}$$

and let us construct W as if $\nu=0$ i.e. $N=Z$. Then

$$|\lambda_N|\le C_1\max\bigl(Z^{\frac{8}{9}},\, B^{\frac{2}{3}}\bigr). \tag{5.3.2}$$

(ii) Let

$$(Z-N)_+ \ge K = C_0 \max\bigl(Z^{\frac{2}{3}},\, Z^{\frac{2}{5}}B^{\frac{1}{5}}\bigr) \tag{5.3.3}$$

with sufficiently large C_0. Then $\lambda_N \asymp \nu$ and

$$|\lambda_N - \nu| \le C_1 \max\bigl(Z^{\frac{2}{3}},\, B^{\frac{1}{2}}\bigr)|\nu|^{\frac{1}{4}}. \tag{5.3.4}$$

Proof. (i) One can see easily that

(5.3.5) In the framework of (i) as $B \ge (Z-N)_+^{\frac{4}{3}}$ expression (5.2.5) is

$$\asymp B|\lambda_N|^{\frac{1}{2}} \times \underbrace{\Bigl(\frac{|\lambda_N|}{B^2}\Bigr)^{\frac{1}{4}}\bar{r}^2} \asymp |\lambda_N|^{\frac{3}{4}} \min\bigl(1,\, B^{-\frac{3}{10}}Z^{\frac{2}{5}}\bigr)$$

where $\bigl(|\lambda_N|/B^2\bigr)^{\frac{1}{4}}$ is a width of the zone where $W \le -\lambda_N$ and the selected factor is the volume of this zone; this expression should be less than $C\max\bigl(Z^{\frac{2}{3}},\, Z^{\frac{2}{5}}B^{\frac{1}{5}}\bigr)$ which is exactly an error estimate in the semiclassical expression for N. Thus

$$|\lambda_N|^{\frac{3}{4}} \min\bigl(1,\, B^{-\frac{3}{10}}Z^{\frac{2}{5}}\bigr) \le C\max\bigl(Z^{\frac{2}{3}},\, Z^{\frac{2}{5}}B^{\frac{1}{5}}\bigr) \tag{5.3.6}$$

where everywhere the first and the second cases are as $B \le Z^{\frac{4}{3}}$ and $Z^{\frac{4}{3}} \le B \le Z^3$ respectively. The last inequality is equivalent to (25-6-13).

As $B \le (Z-N)_+^{\frac{4}{3}}$ inequality (5.3.6) is replaced by $|\lambda_N|^{\frac{3}{4}} \le CZ^{\frac{2}{3}}$ which coincides with (5.3.6) with B reset to $(Z-N)_+^{\frac{4}{3}}$ and also with the same inequality derived for $B=0$ in the previous Chapter; therefore (5.3.2) holds in this case as well.

(ii) One can prove easily that

(5.3.7) if (5.3.3) is fulfilled, and expression (5.2.5) does not exceed a semiclassical error then $\lambda_N \asymp \nu$ and, furthermore, expression (5.2.5) is

$$\asymp B|\lambda_N - \nu| \int P''_B(W+\nu)\,dx \asymp B|\lambda_N-\nu| \int (W+\nu)_+^{-\frac{1}{2}}\,dx \tag{5.3.8}$$

which for $B \ge (Z-N)_+^{\frac{4}{3}}$ is

$$\asymp B\bar{r}^2|\lambda_N-\nu|\cdot|\nu|^{-\frac{1}{2}}\Bigl(\frac{|\nu|}{B^2}\Bigr)^{\frac{1}{4}} \asymp |\lambda_N-\nu|\cdot|\nu|^{-\frac{1}{4}}\min\bigl(1,\, B^{-\frac{3}{10}}Z^{\frac{2}{5}}\bigr) \tag{5.3.9}$$

and this should be less than $C\max\bigl(Z^{\frac{2}{3}},\ Z^{\frac{2}{5}}B^{\frac{1}{5}}\bigr)$ which implies (5.3.4).

On the other hand, as $B\le (Z-N)_+^{\frac{4}{3}}$ the right-hand expression of (5.3.8) is $\asymp |\lambda_N-\nu|\bar{r}\asymp |\lambda_N-\nu|(Z-N)_+^{-\frac{1}{3}}$ and this should be less than $CZ^{\frac{2}{3}}$ which implies (5.3.4) in this case as well. □

Proposition 5.3.1 immediately implies

Corollary 5.3.2. *In the framework of proposition (i), (ii),*

$$|\lambda_N-\nu|\cdot \mathsf{N}([\lambda_N,\nu])\le C\max(Z^{\frac{5}{3}},Z^{\frac{3}{5}}B^{\frac{4}{5}}) \tag{5.3.10}$$

where $\mathsf{N}(\lambda_N,\nu)$ *is the number of (non-zero) eigenvalues on interval* $[\lambda_N,\nu]$ *or* $[\nu,\lambda_N]$.

5.3.2 Estimate for D-terms

Proposition 5.3.3. *In the framework of Proposition (i), (ii)*

$$\mathsf{D}\bigl(e(x,x,\lambda)-P'_B(W+\lambda),\ e(x,x,\lambda)-P'_B(W+\lambda)\bigr) \tag{5.3.11}$$

with $\lambda=\nu,\lambda_N$ *respectively and*

$$\mathsf{D}\bigl(P'_B(W+\nu)-P'_B(W+\lambda),\ P'_B(W+\nu)-P'_B(W+\lambda_N)\bigr) \tag{5.3.12}$$

with $\lambda=\lambda_N$ *do not exceed* $C\max(Z^{\frac{5}{3}},Z^{\frac{3}{5}}B^{\frac{4}{5}})$.

Proof. Term (5.3.11) with $\lambda=\nu$ has been estimated this way and the same estimate for this term with $\lambda=\lambda_N$ is proven in the same way; we leave easy details to the reader;

Term (5.3.12) is estimated using proposition 5.3.1; again we leave easy details to the reader. □

Remark 5.3.4. Let $B\le Z$. Then in (5.3.1)–(5.3.4) and therefore also in (5.3.8) and in proposition 5.3.3 one can replace C_0 and C by $C_0\varepsilon$ and $C\varepsilon$ respectively with the small parameter ε: $\max\bigl(Z^{-\delta},(BZ^{-1})^{\delta}\bigr)\le\varepsilon\le 1$.

5.3.3 Summary

Then following the scheme of subsection 24.4.4 of [Ivr11] we arrive to upper estimates in Theorem 5.3.5 below (lower estimates have been proven in Proposition 5.1.1); we also arrive to Theorem 5.3.6 below:

Theorem 5.3.5. *Let $M=1$, $B\le Z^3$. Then*

(i) The following estimate holds:

$$E^{\mathsf{TF}} \le \mathcal{E}^{\mathsf{TF}} + \Bigl(\operatorname{Tr}(H_{A,W}-\nu)^- + \int P_B(W^{\mathsf{TF}}(x)+\nu)\,dx\Bigr) + CR \tag{5.3.13}$$

with $R=\max\bigl(Z^{\frac{5}{3}}, Z^{\frac{3}{5}}B^{\frac{4}{5}}\bigr)$*;*

(ii) The following estimate holds:

$$E^{\mathsf{TF}} \le \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} + CR - CZ^{\frac{4}{3}}B^{\frac{1}{3}}; \tag{5.3.14}$$

(iii) As $B\le Z$

$$E^{\mathsf{TF}} \ge \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} + \mathsf{Dirac} + \mathsf{Schwinger} + CZ^{\frac{5}{3}}\bigl(Z^{-\delta} + (BZ^{-1})^{\delta}\bigr). \tag{5.3.15}$$

Theorem 5.3.6. *Let $M=1$, $B\le Z^3$. Then*

(i) The following estimate holds:

$$\mathsf{D}(\rho_\Psi-\rho_B^{\mathsf{TF}}, \rho_\Psi-\rho_B^{\mathsf{TF}}) \le CR; \tag{5.3.16}$$

(ii) As $B\le Z$

$$\mathsf{D}(\rho_\Psi-\rho_B^{\mathsf{TF}}, \rho_\Psi-\rho_B^{\mathsf{TF}}) \le CZ^{\frac{5}{3}}\bigl(Z^{-\delta} + (BZ^{-1})^{\delta}\bigr). \tag{5.3.17}$$

5.4 Upper estimate as $M\ge 2$

5.4.1 Estimate for $|\lambda_N-\nu|$

Again we need to consider two cases: *almost neutral molecules* (systems) when $(Z-N)_+\le C_0K$ with K slightly redefined below and we can set $\nu=0$ in the definition of Thomas-Fermi potential and establish estimate for $|\lambda_N|$ (and for optimal ν we have the same estimate for both $|\nu|$ and λ_N) and *not almost neutral molecules* (systems) when $(Z-N)_+\ge C_0K$ and we can prove that $|\lambda_N|\asymp|\nu|$ and estimate $|\lambda_N-\nu|$.

Proposition 5.4.1. *Let* $M \geq 2$, $B \leq Z^3$ *and condition* (0.4.1) *be fulfilled.*

(i) Let

$$(Z-N)_+ \leq K := C_0 \begin{cases} Z^{\frac{2}{3}} + B^{\frac{1}{2}} L & \text{as } B \leq Z^{\frac{4}{3}}, \\ Z^{\frac{1}{5}} B^{\frac{4}{15}} L & \text{as } Z^{\frac{4}{3}} \leq B \leq Z^3 \end{cases} \tag{5.4.1}$$

and let us construct W *as if* $\nu = 0$ *i.e.* $N = Z$. *Then*

$$|\lambda_N| \leq C_1 \begin{cases} Z^{\frac{8}{9}} + B^{\frac{2}{3}} L^{\frac{4}{3}} & \text{as } B \leq Z^{\frac{4}{3}}, \\ Z^{\frac{1}{5}} B^{\frac{4}{15}} L^{\frac{4}{3}} & \text{as } Z^{\frac{4}{3}} \leq B \leq Z^3; \end{cases} \tag{5.4.2}$$

recall that $L = |\log BZ^{-3}|$;

(ii) Let

$$(Z-N)_+ \geq K \tag{5.4.3}$$

with sufficiently large C_0. *Then* $\lambda_N \asymp \nu$ *and moreover*

$$|\lambda_N - \nu| \leq C \max\bigl(Z^{\frac{2}{3}}, B^{\frac{1}{2}} L_1\bigr) |\nu|^{\frac{1}{4}}; \tag{5.4.4}$$

recall that

$$L_1 = \begin{cases} 1 & \text{as } B \leq (Z-N)_+^{\frac{4}{3}}, \\ |\log((Z-N)_+/B^{\frac{3}{4}}| & \text{as } (Z-N)_+^{\frac{4}{3}} \leq B \leq Z^{\frac{4}{3}}, \\ |\log(Z-N)_+/B^{\frac{4}{5}} Z^{\frac{3}{5}}|) & \text{as } Z^{\frac{4}{3}} \leq B \leq Z^3, \end{cases} \tag{5.4.5}$$

Proof. We will apply arguments slightly more sophisticated than the obvious ones. These better arguments will allow us to derive slightly better estimates for $|\lambda_N - \nu|$ as $(Z-N)_+ \geq CK$.

Recall that estimates for $|\lambda_N - \nu|$ are derived by comparison of expression (5.2.5) and semiclassical errors for the number of eigenvalues below $\lambda = \nu$ and $\lambda = \lambda_N$: expression (5.2.5) should be less than the sum of these semiclassical errors.

Consider contribution of each ball

$$B(x, \ell(x)) \subset \mathcal{Y} = \{x : \min_m |x - \mathsf{y}_m| \geq \epsilon \bar{r}\} \tag{5.4.6}$$

to semiclassical errors as $\lambda = \nu$ and $\lambda = \lambda_N$ and compare it with its contribution to (5.2.5).

(a) Each ball contributes no more than $CB\ell^2$ to the first error (with $\lambda=\nu$) where due to our choice $\zeta\ell\ge 1$.

(b) Further, each ball with $\zeta\ge C_1|\lambda_N-\nu|^{\frac{1}{2}}$ contributes no more than $CB\ell^2$, but each ball with $\zeta\le C_1|\lambda_N-\nu|^{\frac{1}{2}}$ contributes no more than $CB\ell^{3-\delta}|\lambda_N-\nu|^{\delta/2}$ to the second error (with $\lambda=\lambda_N$); here $\delta=\frac{1}{3}$ is due to rescaling.

(c) Meanwhile each ball with $\zeta\ge C_1|\lambda_N-\nu|^{\frac{1}{2}}$ contributes no less than $\epsilon_0 B|\lambda_N-\nu|\zeta^{-1}\ell^3$, and each ball with $\zeta\ge C_1|\lambda_N-\nu|^{\frac{1}{2}}$ contributes no less than $\epsilon_0|\lambda_N-\nu|^{\frac{1}{2}}\ell^3$ to expression (5.2.5) and it is larger than its contributions to each semiclassical error (multiplied by C) as long as

$$(5.4.7)_{1,2}\qquad \zeta^2\ge|\lambda_N-\nu|\ge C_2\zeta\ell^{-1},\qquad |\lambda_N-\nu|\ge C_2\ell^{-2}.$$

Obviously in (i), (ii) we can assume that

(5.4.8) (5.4.2) and (5.4.4) respectively (with C replaced by arbitrarily large C_3) are violated.

(i) Assume first that $B\ge(Z-N)_+^{\frac{4}{3}}$.

Then in the framework of (i) $\zeta=B^2\ell^4$ with minimal $\ell=B^{-\frac{1}{3}}$ and therefore $(5.4.7)_{1,2}$ are fulfilled for $\ell\le C_2B^{-1}|\lambda_N|$. Therefore we need to account for semiclassical errors contributed by an inner shell (not exceeding $C\max(Z^{\frac{2}{3}},B^{\frac{1}{2}})$) and by zone $\mathcal{Y}\cap\{\ell\ge C_2B^{-1}|\lambda_N|\}$; there $\zeta\ge C_1|\lambda_N|^{\frac{1}{2}}$ and therefore its contribution does not exceed $CB\int\ell(x)^{-1}\,dx$ with integral over this zone and it does not exceed $CB\bar{r}^2L$.

So, the truncated semiclassical errors do not exceed $C\max(Z^{\frac{2}{3}},B\bar{r}^2L)$. Meanwhile expression (5.2.5) is no less than $CB^{\frac{1}{2}}\bar{r}^2|\lambda_N|^{\frac{3}{4}}$. Therefore comparing these two expressions as $B\le Z^{\frac{4}{3}}$ and as $Z^{\frac{4}{3}}\le B\le Z^3$ we arrive to (5.4.2).

Consider remaining case $B\le(Z-N)_+^{\frac{4}{3}}$. Semiclassical arguments remain valid while estimate of (5.2.5) from below by $\epsilon_0|\lambda_N|^{\frac{3}{4}}$ also could be proven easily.

(ii) Again, assume first that $(Z-N)_+^{\frac{4}{3}}\le B\le Z^3$. Again, in the calculation of the truncated semiclassical errors we integrate over $\ell\ge C_2B^{-1}|\lambda_N-\nu|$ where

$\zeta \geq C_1|\lambda_N|^{\frac{1}{2}}$ and therefore its contribution does not exceed $CB\int \ell(x)^{-1}\,dx$ with integral over this zone and it does not exceed $CB\bar{r}^2 L_1$[22].

Again, expression (5.3.8) is larger than the expressions afterwards and comparing with the semiclassical error estimate we arrive to (5.4.4).

Consider remaining case $B \leq (Z-N)_+^{\frac{4}{3}}$. Semiclassical arguments remain valid while estimate of (5.2.5) from below by $\epsilon_0|\lambda_N - \nu|\cdot^{-\frac{1}{4}}$ also could be proven easily. $\square$

Corollary 5.4.2. *In the framework of proposition 5.4.1* $|\lambda_N-\nu|\cdot \mathsf{N}([\lambda_N,\nu])$ *does not exceed expression* (4.4.6)*.*

5.4.2 Estimate for D-terms under assumption (5.4.1)

We need to estimate semiclassical error D-term (5.3.11) with $\lambda=\lambda_N$ because for $\lambda=\nu$ we already estimated it, and also we need to estimate a term (5.3.12). We start from the latter one. Recall that under assumption (5.4.1) we take $\nu=0$. The trivial estimate is based on

$$|P'_B(W) - P'_B(W+\lambda)| \leq CW^{\frac{1}{2}}|\lambda| + CBW^{-\frac{1}{4}}|\lambda|^{\frac{3}{4}}, \tag{5.4.9}$$

leading to

$$J \leq C\mathsf{D}(W^{\frac{1}{2}}, W^{\frac{1}{2}})|\lambda|^2 + CB^2|\lambda|^{\frac{3}{2}}\mathsf{D}(W^{-\frac{1}{4}}\theta, W^{-\frac{1}{4}}\theta) \tag{5.4.10}$$

where here and below J is expression (5.3.12), θ is a characteristic function of domain$\{x:\, \gamma(x)\geq \hbar^{\frac{1}{3}}\}$ and we can ignore contribution of $\{x:\,\gamma(x)\leq \hbar^{\frac{1}{3}}\}$. Really, contribution of this zone does not exceed a semiclassical error estimate CRL^2.

Note that even without assumption (5.4.1)

$$\mathsf{D}(W^{\frac{1}{2}}, W^{\frac{1}{2}}) \asymp (B^{-\frac{1}{4}};\, B^{-\frac{8}{5}}Z^{\frac{9}{5}}) \qquad \text{as}\ \ B\leq Z^{\frac{4}{3}},\ Z^{\frac{4}{3}}\leq B\leq Z^3 \tag{5.4.11}$$

respectively and (5.4.2) implies that the first term is much less than R.

Meanwhile under assumption (5.4.1)

$$\mathsf{D}(W^{-\frac{1}{4}}\theta, W^{-\frac{1}{4}}\theta) \asymp B^{-1}\mathsf{D}(\ell^{-1}\theta,\ell^{-1}\theta) \asymp B^{-1}\bar{r}^3\mathsf{D}(\gamma^{-1}\theta,\gamma^{-1}\theta) \tag{5.4.12}$$

---

[22] In (i) this would not lead to any improvement.

where in the right-hand expression D and γ,θ are in the scale $x\mapsto x\bar{r}^{-1}$ and then $\mathsf{D}(\gamma^{-1}\theta,\gamma^{-1}\theta)\asymp L^2$ so the second term in (5.4.10) does not exceed $CB\bar{r}^3|\lambda_N|^{\frac{3}{2}}L^2$ which due to (5.4.2) does not exceed

$$C\max\bigl(Z^{\frac{5}{3}},\, Z^{\frac{3}{5}}B^{\frac{3}{5}}L^4\bigr). \tag{5.4.13}$$

Consider now term (5.3.11) with $\lambda=\lambda_N$. Let us consider zones $\Omega_1:=\{x:\,|\lambda-\nu|\lesssim\zeta\ell^{-1}\}$ and $\Omega_2:=\{x:\,|\lambda-\nu|\gtrsim\zeta\ell^{-1}\}$.

Remark 5.4.3. Contribution to the term in question of each couple of balls contained in $\Omega_1\times\Omega_1$ does not exceed estimate for the same term with $\lambda=\nu$; really, after rescaling $x\mapsto x/\ell$ and $\tau\mapsto\tau/\zeta^2$ we conclude that the difference between energy levels does not exceed local semiclassical parameter $C/(\zeta\ell)$.

Therefore the total contribution of $\Omega_1\times\Omega_1$ to this term does not exceed $C\max\bigl(Z^{\frac{5}{3}},\, Z^{\frac{3}{5}}B^{\frac{3}{5}}L^2\bigr)$.

On the other hand, contribution to the term in question of each couple of balls contained in $\Omega_2\times\Omega_2$ does not exceed its contribution to (5.3.12) and therefore the total contribution of $\Omega_2\times\Omega_2$ to this term does not exceed expression (5.4.13).

So, term (5.3.11) with $\lambda=\lambda_N$ does not exceed (5.4.13).

Therefore we arrive immediately to

Theorem 5.4.4. *Let $M\ge 2$, $B\le Z^3$ and condition* (0.4.1) *be fulfilled. Then under assumption* (5.4.1)

(i) The following estimate holds:

$$E^{\mathsf{TF}}\le\mathcal{E}^{\mathsf{TF}}+\Bigl(\operatorname{Tr}(H_{A,W}-\nu)^-+\int P_B(W^{\mathsf{TF}}+\nu)\,dx\Bigr)+ C\max\bigl(Z^{\frac{5}{3}},\, Z^{\frac{3}{5}}B^{\frac{3}{5}}L^4\bigr); \tag{5.4.14}$$

(ii) As $\mathsf{a}\ge Z^{-1}$ The following estimate holds:

$$E^{\mathsf{TF}}\le\mathcal{E}^{\mathsf{TF}}+\mathsf{Scott}+C\max\bigl(Z^{\frac{5}{3}},\, Z^{\frac{3}{5}}B^{\frac{3}{5}}L^4\bigr)+CZ^{\frac{4}{3}}B^{\frac{1}{3}}+C\mathsf{a}^{-\frac{1}{2}}Z^{\frac{3}{2}}; \tag{5.4.15}$$

as $\mathsf{a}\le Z^{-1}$ one should replace the last term in the right-hand expression by CZ^2 and skip Scott*;*

(iii) As $B \le Z$ *and* $a \ge Z^{-\frac{1}{3}}$

$$E^{\mathsf{TF}} \le \mathcal{E}^{\mathsf{TF}} + \mathsf{Scott} + \mathsf{Dirac} + \mathsf{Schwinger} + CZ^{\frac{5}{3}}\bigl(Z^{-\delta} + (BZ^{-1})^{\delta} + (aZ^{\frac{1}{3}})^{-\delta}\bigr). \tag{5.4.16}$$

Here proof of (iii) is due to the same arguments as in the case $B = 0$. Combining with the estimate from below we also conclude that

Theorem 5.4.5. *(i) In the framework of theorem 5.4.4 the following estimate holds:*

$$\mathsf{D}\bigl(\rho_\Psi - \rho^{\mathsf{TF}},\, \rho_\Psi - \rho^{\mathsf{TF}}\bigr) \le C\max\bigl(Z^{\frac{5}{3}},\, Z^{\frac{3}{5}}B^{\frac{3}{5}}L^4\bigr); \tag{5.4.17}$$

(ii) In the framework of theorem 5.4.4(iii) the following estimate holds:

$$\mathsf{D}\bigl(\rho_\Psi - \rho^{\mathsf{TF}},\, \rho_\Psi - \rho^{\mathsf{TF}}\bigr) \le CZ^{\frac{5}{3}}\bigl(Z^{-\delta} + (BZ^{-1})^{\delta} + (aZ^{\frac{1}{3}})^{-\delta}\bigr). \tag{5.4.18}$$

5.4.3 Estimate for D-terms under assumption (5.4.3)

Let assumption (5.4.3) be fulfilled. Let $W = W_\nu$ and $\ell = \ell_\nu$ be a potential and a scaling function (used to derive semiclassical remainder estimates) for this $\nu < 0$ (and $N < Z$) while W_0 and ℓ_0 be a potential and a scaling function for $\nu = 0$ (and $N = Z$).

Let us start from the trivial arguments. Note that

$$|P'_B(W + \lambda) - P'_B(W + \nu)| \le CW^{\frac{1}{2}}|\lambda - \nu|\theta_1 + CB|\lambda - \nu|^{\frac{1}{2}}\theta_2 \tag{5.4.19}$$

where θ_1 and θ_2 are characteristic functions of $\mathcal{Y}_1 = \{x:\ W(x) + \nu \ge |\nu|\}$ and $\{x:\ 0 < W(x) + \nu \le |\nu|\}$ respectively. Let

$$J_k := \mathsf{D}\bigl([P'_B(W + \lambda_N) - P'_B(W + \nu)]\theta_k, [P'_B(W + \lambda_N) - P'_B(W + \nu)]\theta_k\bigr).$$

Then in virtue of (5.4.19)

$$J_1 \le C\mathsf{D}(W^{\frac{1}{2}}\theta_1, W^{\frac{1}{2}}\theta_1)|\lambda_N - \nu|^2.$$

Note that $\mathsf{D}(W^{\frac{1}{2}}\theta_1, W^{\frac{1}{2}}\theta_1) \asymp \bigl((Z - N)_+^{-\frac{1}{3}},\, B^{-\frac{1}{4}},\, Z^{\frac{9}{5}}B^{-\frac{8}{5}}\bigr)$ in our cases and using (5.4.4) one can prove easily that $J_1 \le CZ^{\frac{5}{3}}$.

However estimate for a contribution of zone $\{x:\, 0 < W(x) \le C_0|\nu|\}$ is much worse:

$$J_2 \le CB^2\mathsf{D}(\theta_2,\theta_2)|\lambda_N - \nu| \le CB^2\bar{r}^3(|\nu|/B^2)^{\frac{1}{2}}L_1^2|\lambda_N-\nu|, \tag{5.4.20}$$

where for $B \le (Z-N)_+^{\frac{1}{3}}$ we should replace $(|\nu|/B^2)^{\frac{1}{2}}L_1^2$ by $\bar{r}^2$. Using (5.4.4) we conclude that for $(Z-N)_+^{\frac{4}{3}} \lesssim B \lesssim Z^3$

$$J_2 \le CB\bar{r}^3|\nu|^{\frac{3}{4}}L_1^2\max(Z^{\frac{2}{3}}, B^{\frac{1}{2}}) \asymp CB(Z-N)_+^{\frac{3}{4}}\bar{r}^{\frac{9}{4}}\max(Z^{\frac{2}{3}}, B^{\frac{1}{2}}L_1)L_1^2,$$

and therefore we arrive to the last two cases below; the first case is proven similarly:

$$J_2 \le C\begin{cases} (Z-N)_+^{-\frac{4}{3}}Z^{\frac{2}{3}}B^2,\\ (Z-N)_+^{\frac{3}{4}}B^{\frac{7}{16}}\max(Z^{\frac{2}{3}}, B^{\frac{1}{2}}L_1)L_1^2,\\ (Z-N)_+^{\frac{3}{4}}Z^{\frac{9}{20}}B^{\frac{3}{5}}L_1^3. \end{cases} \tag{5.4.21}$$

This is not a good estimate. To improve it let us note that

(5.4.22) If $|\lambda_N-\nu| \le C\max(B^{\frac{2}{3}}, (Z-N)_+^{\frac{8}{9}})$ then J_2 does not exceed (4.4.6)

and therefore we can assume that

$$|\lambda_N-\nu| \ge C\max(B^{\frac{2}{3}}, (Z-N)_+^{\frac{8}{9}}). \tag{5.4.23}$$

Proposition 5.4.6. *(i) Let* $(Z-N)_+^{\frac{4}{3}} \le B \le Z^3$ *and*

$$C_0B^{\frac{2}{3}} \le |\lambda_N-\nu| \le C_1B^{\frac{1}{2}}|\nu|^{\frac{1}{4}}, \tag{5.4.24}$$

then the truncated semiclassical error does not exceed

$$M := CZ^{\frac{2}{3}} + CB\bar{r}^2(|\nu|/B^2)^{\frac{1}{4}}L \times (B^{-1}|\lambda_N-\nu|)^{-1}; \tag{5.4.25}$$

(ii) Let $(Z-N)_+^{\frac{4}{3}} \le B \le Z^3$ *and*

$$C_1B^{\frac{1}{2}}|\nu|^{\frac{1}{4}} \le |\lambda_N-\nu| \le C_1B^{\frac{1}{2}}|\nu|^{\frac{1}{4}}L, \tag{5.4.26}$$

then the truncated semiclassical error does not exceed

$$M := CZ^{\frac{2}{3}} + CB\bar{r}^2L; \tag{5.4.27}$$

(iii) Let $B \le (Z-N)_+^{\frac{4}{3}}$ *and*

$$(Z-N)_+^{\frac{8}{9}} \le |\lambda_N - \nu| \le C_1 (Z-N)_+, \tag{5.4.28}$$

then the truncated semiclassical error does not exceed

$$M := CZ^{\frac{2}{3}} + C(Z-N)_+^{\frac{5}{3}} |\lambda_N - \nu|^{-1}; \tag{5.4.29}$$

(iv) Let $B \le (Z-N)_+^{\frac{4}{3}}$ *and*

$$C_0 (Z-N)_+ \le |\lambda_N - \nu| \le C_1 Z^{\frac{2}{3}} (Z-N)_+^{\frac{1}{3}}, \tag{5.4.30}$$

then the truncated semiclassical error does not exceed $M := CZ^{\frac{2}{3}}$.

Proof. Easy proof using arguments of the proof of proposition 5.4.1 is left to the reader. ☐

Proposition 5.4.7. *In the framework of proposition (i)–(iv) term* (5.3.12) *does not exceed* $CM^{\frac{5}{3}}(B|\lambda_N - \nu|^{\frac{1}{2}})^{\frac{1}{3}} +$ (4.4.6) *with* M *defined defined in the corresponding cases in Proposition 5.4.6.*

Proof. Using proposition 5.4.1 one can prove easily that

(5.4.31) The contribution of $\{x : \min_m |x - \mathsf{y}_m| \le \epsilon \bar{r}\}$ to (5.3.12) does not exceed $CZ^{\frac{5}{3}}$.

Now we need to estimate excess of expression (5.3.12) over semiclassical D-term (with $\lambda = \nu$) which has been estimated by (4.4.6). To do so we need to estimate

$$\mathsf{D}\bigl([P_B(W+\nu) - P_B(W+\lambda)]\theta,\, [P_B(W+\nu) - P_B(W+\lambda)]\theta\bigr) \tag{5.4.32}$$

which is the contribution of the domain $\Omega' := \{x : \ell(x) \le CB^{-1}|\lambda - \nu|\}$ where θ is the characteristic function of Ω'. Recall that on the complimentary domain $|P_B(W+\nu) - P_B(W+\lambda)| \le C\ell^{-1}$. Let us consider

$$\mathsf{D}\bigl([P_B(W+\nu) - P_B(W+\lambda)]\theta_0,\, [P_B(W+\nu) - P_B(W+\lambda)]\theta_0\bigr) \tag{5.4.33}$$

and

$$\text{(5.4.34)}\quad \mathsf{D}\bigl([P_B(W+\nu)-P_B(W+\lambda)]\theta_t,\,[P_B(W+\nu)-P_B(W+\lambda)]\theta_{t'}\bigr)$$

where θ_0 is a characteristic function of

$$\Omega'_0 := \{x:\ \ell(x)\le t_0 := (|\lambda_N-\nu|B^{-2})^{\frac14}\}$$

and θ_t is a characteristic function of $\Omega'_t := \{x:\ t\le \ell(x)\le 2t\}$ with $t\ge t'\ge t_0$.

When calculating (5.2.5) the contribution of Ω'_0 is $\asymp B|\lambda_N-\nu|^{\frac12}\,\mathsf{mes}(\Omega'_0)$ and therefore due to proposition 5.4.6

$$\mathsf{mes}(\Omega'_0)\le CM\bigl(B|\lambda_N-\nu|^{\frac12}\bigr)^{-1} \tag{5.4.35}$$

while term (5.4.33) is

$$\asymp B^2|\lambda_N-\nu|\mathsf{D}(\theta_0,\theta_0)\le CB^2|\lambda_N-\nu|\bigl(\mathsf{mes}(\Omega_0)\bigr)^{\frac53}\le CM^{\frac53}\bigl(B|\lambda_N-\nu|^{\frac12}\bigr)^{\frac13}$$

where the middle inequality

$$\mathsf{D}(\chi_G,\chi_G)\le C\bigl(\mathsf{mes}(G)\bigr)^{\frac53} \tag{5.4.36}$$

is well known[23] and the last one is due to (5.4.35); χ_G denotes characteristic function of G.

Similarly, when calculating expression (5.2.5) one can see easily that the contribution of Ω'_t is $\asymp B|\lambda_N-\nu|(B^2t^4)^{-1}\,\mathsf{mes}(\Omega'_t)$ and therefore

$$\mathsf{mes}(\Omega'_t)\le CM|\lambda_N-\nu|^{-1}t^2 \tag{5.4.37}$$

while term (5.4.34) is $\asymp|\lambda_N-\nu|^2t^{-2}t'^{-2}\mathsf{D}(\theta_t,\theta_{t'})$ which does not exceed

$$C|\lambda_N-\nu|^2t^{-2}t'^{-2}\,\mathsf{mes}(\Omega_t)\,\mathsf{mes}(\Omega_{t'})\bigl[\max\bigl(\mathsf{mes}(\Omega_t),\mathsf{mes}(\Omega_{t'})\bigr)\bigr]^{-\frac13} \tag{5.4.38}$$

due to inequality

$$\mathsf{D}(\chi_G,\chi_{G'})\le C\,\mathsf{mes}(G)\,\mathsf{mes}(G')\bigl[\max\bigl(\mathsf{mes}(G),\mathsf{mes}(G')\bigr)\bigr]^{-\frac13} \tag{5.4.39}$$

---

[23] Really, among uniform solids of equal mass and density the ball has the least potential energy; then $C=\frac15(12\pi)^{\frac13}$.

which trivially follows from the obvious inequality $\mathsf{D}(\chi_G, \delta_z) \le C(\mathsf{mes}(G))^{\frac{2}{3}}$, where $\delta_z(x) = \delta(x-z)$.

Due to (5.4.37) expression (5.4.38) does not exceed $CM^{\frac{5}{3}} t^{-\frac{2}{3}}$; recall that $t \ge t'$. Since summation with respect to $t \ge t'$ and then with respect to $t' \ge t_0$ returns $CM^{\frac{5}{3}} t_0^{-\frac{2}{3}}$ we conclude that term (5.4.33) with θ_0 replaced by θ'' (the characteristic function of $\{x : \ell(x) \ge t_0\}$) also does not exceed $CM^{\frac{5}{3}}(B|\lambda_N - \nu|^{\frac{1}{2}})^{\frac{1}{3}}$. $\square$

So, we have now two estimates for an excess of expression (5.3.12) over (4.4.6): one estimate is $CM^{\frac{5}{3}}(B|\lambda_N - \nu|^{\frac{1}{2}})^{\frac{1}{3}}$ with $M = M(|\lambda_N - \nu|)$ derived in proposition 5.4.6 and another one is due to (5.4.20). Let us consider the best of them. Note that the former estimate consists of two terms each due to the corresponding term in the definition of M. The second term in the framework of proposition 5.4.6(i) is

$$C\bigl(B^{\frac{3}{2}}\bar{r}^2|\nu|^{\frac{1}{4}}L|\lambda_N - \nu|^{-1}\bigr)^{\frac{5}{3}}\bigl(B|\lambda_N - \nu|^{\frac{1}{2}}\bigr)^{\frac{1}{3}} \asymp B^{\frac{17}{6}}\bar{r}^{\frac{10}{3}}|\nu|^{\frac{5}{12}}L^{\frac{5}{3}}|\lambda_N - \nu|^{-\frac{3}{2}}$$

and taking minimum of this expression and $CB\bar{r}^3|\nu|^{\frac{1}{2}}L^2|\lambda_N - \nu|$ we see that this minimum does not exceed

$$C\bigl(B^{\frac{17}{6}}\bar{r}^{\frac{10}{3}}|\nu|^{\frac{5}{12}}L^{\frac{5}{3}}\bigr)^{\frac{2}{5}}\bigl(B\bar{r}^3|\nu|^{\frac{1}{2}}L^2\bigr)^{\frac{3}{5}} \asymp CB^{\frac{5}{3}}\bar{r}^{\frac{8}{3}}(Z-N)_+^{\frac{7}{15}}L^{\frac{28}{15}} \asymp \\ C\begin{cases} (Z-N)_+^{\frac{7}{15}}BL^{\frac{28}{15}} & (Z-N)_+^{\frac{4}{3}} \le B \le Z_+^{\frac{4}{3}}, \\ (Z-N)_+^{\frac{7}{15}}Z_+^{\frac{8}{15}}B^{\frac{3}{5}}L^{\frac{28}{15}} & Z_+^{\frac{4}{3}} \le B \le Z^3 \end{cases} \tag{5.4.40}$$

which is achieved as $|\lambda_N - \nu| \asymp B^{\frac{11}{15}}\bar{r}^{\frac{2}{15}}|\nu|^{-\frac{1}{30}}L^{-\frac{2}{15}}$ and one can see easily that it is in the framework of assumption of proposition 5.4.6(i).

Furthermore, the first term is $(Z^{\frac{2}{3}})^{\frac{5}{3}}(B|\lambda_N - \nu|^{\frac{1}{2}})^{\frac{1}{3}}$ and it reaches its maximum at the largest value of $|\lambda_N - \nu|$ (which is $\mathsf{max}(Z^{\frac{2}{3}}, B^{\frac{1}{2}}L)|\nu|^{\frac{1}{4}}$) and we leave to the reader to check that it does not exceed (5.4.40).

In the framework of proposition 5.4.6(iii) one should replace $(\nu/B^2)^{\frac{1}{4}}$ by $\bar{r}$ and L by 1, so $B^{\frac{3}{2}}\bar{r}^2|\nu|^{\frac{1}{4}}L|\lambda_N - \nu|^{-1} \mapsto B^2\bar{r}^3|\lambda_N - \nu|^{-1}$; further, one should preserve $B|\lambda_N - \nu|^{\frac{1}{2}}$ and therefore the second term becomes

$$\bigl(B^2\bar{r}^3|\lambda_N - \nu|^{-1}\bigr)^{\frac{5}{3}}\bigl(B|\lambda_N - \nu|^{\frac{1}{2}}\bigr)^{\frac{1}{3}} \asymp B^{\frac{11}{3}}\bar{r}^5|\lambda_N - \nu|^{-\frac{3}{2}}$$

which should be run versus $\bigl(B|\lambda_N - \nu|^{\frac{1}{2}} + |\nu|^{\frac{1}{2}}|\lambda_N - \nu|\bigr)^2\bar{r}^5$ resulting in

$$CB^{\frac{8}{3}}\bar{r}^5 + CB^{\frac{44}{21}}|\nu|^{\frac{3}{7}}\bar{r}^5 \asymp CB^{\frac{8}{3}}(Z-N)_+^{-\frac{5}{3}} + CB^{\frac{44}{21}}(Z-N)_+^{-\frac{23}{21}}$$

and both of them are less than in (4.4.6). Meanwhile the first term becomes $Z^{\frac{10}{9}}B^{\frac{1}{3}}|\lambda_N-\nu|^{\frac{1}{6}}\le (Z-N)_+^{\frac{1}{18}}Z^{\frac{11}{9}}B^{\frac{1}{3}}$. Combining with (5.4.40) we arrive to

$$C\left\{\begin{aligned} &(Z-N)_+^{\frac{1}{18}}Z^{\frac{11}{9}}B^{\frac{1}{3}} &&B\le (Z-N)_+^{\frac{4}{3}},\\ &(Z-N)_+^{\frac{7}{15}}BL^{\frac{28}{15}} &&(Z-N)_+^{\frac{4}{3}}\le B\le Z_+^{\frac{4}{3}},\\ &(Z-N)_+^{\frac{7}{15}}Z_+^{\frac{8}{15}}B^{\frac{3}{5}}L^{\frac{28}{15}} &&Z_+^{\frac{4}{3}}\le B\le Z^3.\end{aligned}\right. \tag{5.4.41}$$

5.4.4 Summary

Theorem 5.4.8. *Let* $M\ge 2$, $B\le Z^3$. *Then*

(i) The following estimate holds:

$$E^{\mathsf{TF}}\le \mathcal{E}^{\mathsf{TF}}+\Bigl(\operatorname{Tr}(H_{A,W}-\nu)^-+\int P_B(W^{\mathsf{TF}}+\nu)\,dx\Bigr)+CR+ \\ (4.4.6)+(5.4.41); \tag{5.4.42}$$

(ii) The following estimate holds:

$$E^{\mathsf{TF}}\le \mathcal{E}^{\mathsf{TF}}+\mathsf{Scott}+CR+CZ^{\frac{4}{3}}B^{\frac{1}{3}}+(4.4.6)+(5.4.41); \tag{5.4.43}$$

(iii) As $B\le Z$ *and* $a\ge Z^{-\frac{1}{3}}$

$$E^{\mathsf{TF}}\ge \mathcal{E}^{\mathsf{TF}}+\mathsf{Scott}+\mathsf{Dirac}+\mathsf{Schwinger}+ \\ CZ^{\frac{5}{3}}\bigl(Z^{-\delta}+(BZ^{-1})^{\delta}+(aZ^{\frac{1}{3}})^{-\delta}\bigr) \tag{5.4.44}$$

Remark 5.4.9. Unfortunately there are gaps in the proofs of V. Ivrii [Ivr2] in the case of $M\ge 2$ and large $Z-N>0$ and I was unable to fill them.

6 Negatively charged systems (coming)

7 Positively charged systems (coming)

8 Appendices

8.A Electrostatic inequalities

There are two kinds of electrostatic inequalities: those which hold for any fermionic state Ψ and those which hold only for ground-state (or near ground state) Ψ. Inequalities of the first kind do not depend on the quantum Hamiltonian and they are (24.2.1) of [Ivr11] repeated here:

$$\sum_{1\le j<k\le N}\int |x_j-x_k|^{-1}|\Psi(x_1,\ldots,x_N)|^2\,dx_1\cdots dx_N \ge \frac{1}{2}\mathsf{D}(\rho_\Psi,\rho_\Psi)-C\int \rho_\Psi^{\frac{4}{3}}(x)\,dx \tag{8.A.1}$$

and (8.A.5) below.

Inequalities of the second kind are as $B=0$:

$$\sum_{1\le j<k\le N}\int |x_j-x_k|^{-1}|\Psi(x_1,\ldots,x_N)|^2\,dx_1\cdots dx_N \ge \frac{1}{2}\mathsf{D}(\rho_\Psi,\rho_\Psi)-CZ^{\frac{5}{3}}, \tag{8.A.2}$$

and more precise one (8.A.26) below

As $\vec{B}=\mathsf{const}$ there is inequality established in E. Lieb, J. P. Solovej and J. Yngvarsson [LSY2] (p. 122):

Theorem 8.A.1. *Let $\vec{B}=\mathsf{const}$. Then for the ground state Ψ*

$$\int \rho_\Psi^{\frac{4}{3}}\,dx \le CZ^{\frac{5}{6}}N^{\frac{1}{2}}(Z+N)^{\frac{1}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}} \tag{8.A.3}$$

In particular for $c^{-1}N \le Z \le cN$ *the right-hand expression does not exceed*

$$CZ^{\frac{5}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}} \asymp C\begin{cases} Z^{\frac{5}{3}} & \textit{as}\quad B\le Z^{\frac{4}{3}},\\ Z^{\frac{17}{15}}B^{\frac{2}{5}} & \textit{as}\quad B\ge Z^{\frac{4}{3}}.\end{cases} \tag{8.A.4}$$

We want to establish inequality similar to (24.A.2) of [Ivr11] but in magnetic case. We will use for this the following

Theorem 8.A.2. [24] *Fix* $0<\delta\le 1/6$. *Then for any density matrix* G *an any density* $\rho_0(x)\ge 0$ *we have*

$$\begin{aligned}\sum_{1\le j<k\le N}\int |x_j-x_k|^{-1}|\Psi(x_1,\ldots,x_N)|^2\,dx_1\cdots dx_N \ge\\ \mathsf{D}(\rho_0,\rho_\gamma)-\frac{1}{2}\mathsf{D}(\rho_0,\rho_0)-\frac{1}{2}\sum_{\varsigma,\varsigma'}\iint |G(x,\varsigma;y,\varsigma')|^2|x-y|^{-1}\,dxdy\\ -C\|\rho\|_{5/3}^{5/6}\cdot\|\rho\|_1^{1/6+\delta}\upsilon(\gamma,G)^{1/3-\delta}\end{aligned} \tag{8.A.5}$$

where $\rho=\rho_0+\rho_G+\rho_\Psi$, $\upsilon(\gamma,G):=\mathsf{Tr}(\gamma(I-G))$,

$$\gamma=\gamma_\Psi(x,y)=N\int \Psi(x,x_2,\ldots,x_N)\Psi^\dagger(y,x_2,\ldots,x_N)\,dx_2\cdots x_N \tag{8.A.6}$$

is two-point one particle density.

Recall that $\|.\|_p$ denotes $\mathscr{L}^p$-norm.

Connection between (8.A.1) and (8.A.5): as we set $G=0$ we get $\varsigma=\|\rho\|_1$ and the last term in (8.A.5) becomes $\|\rho\|_{5/3}^{5/6}\cdot\|\rho\|_1^{1/2}$; on the other hand, $\|\rho\|_{4/3}^{4/3}\le\|\rho\|_{5/3}^{5/6}\cdot\|\rho\|_1^{1/2}$, so (8.A.5) is slightly deteriorated (8.A.1) with $G=0$ but with "free" ρ_0.

Let us follow G. Graf and J. P. Solovej [GS] further albeit in the case of magnetic field. Let us estimate first $\|\rho\|_{5/3}$.

---

[24] Lemma 6 of G. Graf and J. P. Solovej [GS].

If $\rho=\rho^{\mathsf{TF}}$ direct calculations show that as $N\asymp Z$

$$\int \rho^{\mathsf{TF}}\,dx=\min(Z,N), \tag{8.A.7}$$

$$\int(\rho^{\mathsf{TF}})^{\frac{4}{3}}\,dx\asymp C\rho^{*\,\frac{4}{3}}r^{*\,3}=CZ^{\frac{5}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}}, \tag{8.A.8}$$

$$\int(\rho^{\mathsf{TF}})^{\frac{5}{3}}\,dx\asymp C\rho^{*\,\frac{5}{3}}r^{*\,3}=CZ^{\frac{7}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{4}{5}} \tag{8.A.9}$$

with

$$r^*=\min(Z^{-\frac{1}{3}},B^{-\frac{2}{5}}Z^{\frac{1}{5}})\asymp Z^{-\frac{1}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{-\frac{2}{5}}, \tag{8.A.10}$$

$$\rho^*=\min(N,Z)r^{*,3} \tag{8.A.11}$$

and we have $\|\rho\|_{5/3}^{5/6}\cdot\|\rho\|_1^{1/2}\asymp\|\rho\|_{4/3}^{4/3}$ for $\rho=\rho^{\mathsf{TF}}$.

If $\rho=\rho_\Psi$ we use magnetic Lieb-Thirring inequality (see f. e. L. Erdš [E])

$$\operatorname{Tr}(H_{A,W})^-\ge -C\int P_B(W)\,dx \tag{8.A.12}$$

and therefore

$$\langle \mathsf{H}\Psi,\Psi\rangle\ge \operatorname{Tr}^-(H_{A,W})+\int W\rho_\Psi\,dx \\ -\int V\rho_\Psi\,dx+\frac{1}{2}\mathsf{D}(\rho_\Psi,\rho_\Psi)-C\|\rho_\Psi\|_{4/3}^{4/3} \tag{8.A.13}$$

which due to (8.A.12) is greater than

$$\int\bigl(-CP_B(W)+W\rho_\Psi\bigr)\,dx- \\ \int V\rho_\Psi\,dx+\frac{1}{2}\mathsf{D}(\rho_\Psi,\rho_\Psi)-C\|\rho_\Psi\|_{4/3}^{4/3}\ge \\ 3\epsilon_0\int\tau_B(\rho_\Psi)\,dx-\int V\rho_\Psi\,dx+\frac{1}{2}\mathsf{D}(\rho_\Psi,\rho_\Psi)-C\|\rho_\Psi\|_{4/3}^{4/3}. \tag{8.A.14}$$

as we picked up $W: CP'_B(W)=\rho_\Psi$.

The first two terms in the right-hand expression are estimated from below by

$$2\epsilon_0\int\tau_B(\rho_\Psi)\,dx-C\int P_B(V)\phi\,dx-C\int V\rho_\Psi(1-\phi)\,dx$$

where $\mathsf{supp}(\phi) \subset \{x, \ell(x) \le 2r^*\}$ and $\mathsf{supp}(1-\phi) \subset \{x, \ell(x) \ge r^*\}$.

One can see easily that the absolute value of the second term is $\asymp Z^{\frac{7}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}}$ while the absolute value of the third term does not exceed $CZ\int V(1-\phi)\,dx \asymp CZ^2 r^{*\,-1}$ which does not exceed the same expression $Z^{\frac{7}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}}$. Therefore

$$\langle \mathsf{H}\Psi,\Psi\rangle + C_1 Z^{\frac{7}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}} \ge \\ 2\epsilon_0\int \tau_B(\rho_\Psi)\,dx + \frac{1}{2}\mathsf{D}(\rho_\Psi,\rho_\Psi) - C\|\rho_\Psi\|_{4/3}^{4/3}. \tag{8.A.15}$$

Note that $\|\rho_\Psi\|_{4/3}^{4/3}$ calculated over domain $\{x:\rho_\Psi(x)\ge B^{\frac{4}{3}}\}$ does not exceed $C\|\rho_\Psi\|_{5/3}^{5/6}\cdot\|\rho\|_1^{\frac{1}{2}}$ with norms calculated over the same domain which does not exceed $CT^{\frac{1}{2}}Z^{\frac{1}{2}}$ with $T=\int \tau_B(\rho_\Psi)\,dx$.

Meanwhile $\|\rho_\Psi\|_{4/3}^{4/3}$ calculated over domain $\{x:\rho_\Psi(x)\ge B^{\frac{4}{3}}\}$ does not exceed $C\|\rho\|_3^{\frac{1}{2}}\cdot\|\rho\|_1^{\frac{5}{6}}$ with norms calculated over the same domain which does not exceed $CZ^{\frac{1}{2}}B^{\frac{1}{3}}T^{\frac{1}{6}}$.

Therefore

$$\|\rho_\Psi\|_{4/3}^{4/3} \le CT^{\frac{1}{2}}Z^{\frac{1}{2}} + CZ^{\frac{5}{6}}B^{\frac{1}{3}}T^{\frac{1}{6}} \tag{8.A.16}$$

and therefore (8.A.15) implies that if

$$\langle \mathsf{H}\Psi,\Psi\rangle \le C_1 Z^{\frac{7}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}} \tag{8.A.17}$$

then

$$T=\int \tau_B(\rho_\Psi)\,dx \le C_2 Z^{\frac{7}{3}}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}} \tag{8.A.18}$$

and

$$\|\rho_\Psi\|_{4/3}^{4/3} \le CZ^{5/3}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}}; \tag{8.A.19}$$

taking $G=0$ we arrive to (8.A.3) as $N\asymp Z$.

However on our preparatory step we need to estimate also $\|\rho_\Psi\|_{5/3}^{5/3}$ and due to (8.A.18) we need to consider only norms over $\{x:\rho_\Psi(x)\ge B^{\frac{4}{3}}\}$. Then $\|\rho_\Psi\|_{5/3}^{5/3} \le C\|\rho_\Psi\|_{4/3}^{16/15}\|\rho_\Psi\|_3^{3/5} \le$ and plugging the same estimates (8.A.19), (8.A.19) we conclude that

$$\|\rho_\Psi\|_{5/3}^{5/3} \le CZ^{7/3}\bigl(1+BZ^{-\frac{4}{3}}\bigr)^{\frac{4}{5}}. \tag{8.A.20}$$

Now we assume that $B \le Z^3$, take $G = e(x,y,\nu)$ where $e(x,y,\nu)$ is the Schwartz kernel of spectral projector for potential W approximating W^{TF} and $\nu \le 0$ is a chemical potential. One can prove easily that $\|\rho_G\|_{5/3}^{5/3}$ satisfies the same estimate and we need to estimate $\operatorname{Tr}\bigl(\gamma_\Psi(I - E(\mu))\bigr)$.

Consider

$$\begin{aligned}
N\langle H_{A,W(x_1)}\Psi,\Psi\rangle &- \operatorname{Tr}(HE(\nu)) - \alpha\operatorname{Tr}\bigl(\gamma_\Psi(I-E(\nu))\bigr)\\
&\ge \int_{\beta<0}\beta\, d_\beta \operatorname{Tr}E(\beta) - \int_{\beta\le\nu}(\beta-\nu+\alpha)\, d_\beta\operatorname{Tr}E(\beta)\\
&= -\int_{\nu-\alpha<\beta<\nu}(\beta-\nu+\alpha)d_\beta E(\beta)\\
&= -\alpha E(\nu) + \int_{\nu-\alpha<\beta<\nu} E(\beta)\, d\beta.
\end{aligned}\tag{8.A.21}$$

We can replace $E(\beta)$ by $\int P'(W+\beta)\,dx$ with a resulting error $O(Z\alpha\hbar^\delta)$, $\hbar := BZ^{-3}$. Then the right-hand expression becomes

$$\begin{aligned}
-L(\alpha) := \int\Bigl(-\alpha P'_B(W+\nu) + \int_0^\alpha P'_B(W+\nu-\beta)\,d\beta\Bigr)\,dx =\\
-\int_0^\alpha (\alpha-\beta)\Bigl(\int P''_B(W+\nu-\beta)\,dx\Bigr)\,d\beta.
\end{aligned}\tag{8.A.22}$$

Therefore

$$\begin{aligned}
\alpha\Bigl(\operatorname{Tr}\bigl(\gamma_\Psi(I-E(\mu))\bigr) - CZ\hbar^\delta\Bigr) \le\\
\underbrace{N\langle H_{A,W(x_1)}\Psi,\Psi\rangle - \operatorname{Tr}(HE(\nu))} + L(\alpha).
\end{aligned}\tag{8.A.23}$$

Note that adding to the selected terms $-\frac{1}{2}\mathsf{D}(\rho^{\mathsf{TF}},\rho^{\mathsf{TF}})$ we obtain exactly the snippet occurring in the lower estimate of E_N but in virtue of the upper estimate it should not exceed $Q = CZ^{\frac{5}{3}}\bigl(1 + BZ^{-\frac{4}{3}}\bigr)^{\frac{2}{5}} \le CZ^{\frac{5}{3}} + C\hbar^2 Z^{\frac{7}{3}}$ and therefore plugging $\alpha = Z^{\frac{4}{3}}\hbar^\delta$ we conclude that

$$\operatorname{Tr}\bigl(\gamma_\Psi(I-E(\mu)\bigr) \le Z\hbar^\delta \tag{8.A.24}$$

provided we prove that

$$L(\alpha) \le Q \qquad \text{as}\ \ \alpha = Z^{\frac{4}{3}}\hbar^\delta. \tag{8.A.25}$$

Therefore modulo proof of (8.A.25) we arrive to the estimate (8.A.26) below:

Theorem 8.A.3. *Let $N \asymp Z$ and $B \le Z$. Then for the ground state energy*

$$\sum_{1\le j<k\le N} \int |x_j - x_k|^{-1} |\Psi(x_1,\dots,x_N)|^2\, dx_1\cdots dx_N \ge \frac{1}{2}\mathsf{D}(\rho^{\mathsf{TF}},\rho^{\mathsf{TF}}) + \mathsf{Dirac} - CZ^{\frac{5}{3}-\delta}(1+B^{\delta}) \tag{8.A.26}$$

To prove (8.A.25) we note that $0 \le P''_B(w) \asymp w^{\frac{1}{2}} + Bw^{-\frac{1}{2}}$. One can prove easily then that $L(\alpha) \le C\alpha^{\frac{7}{4}} + CB^{\frac{1}{4}}\alpha^{\frac{3}{2}}$ which obviously implies (8.A.25).

8.B Very strong magnetic field case

Let us consider now the case $Z^2 \lesssim B \lesssim Z^3$.

Proposition 8.B.1. *Consider Schrödinger operator $H_{A,W}$ with a constant magnetic field of intensity B and potential W: $W \le Z|x|^{-1}$. Let $\phi(x) := \phi_r(x)$ be r-admissible function. Then as $Z^2 \lesssim B \lesssim Z^3$ and $r \asymp Z^{-1}$*

$$|e(x,y,0)| \le CZB \qquad \textit{in}\ \ B(0,r) \tag{8.B.1}$$

and

(8.B.2) *All eigenvalues are $\ge -CZ^2$.*

Proof. Without any loss of the generality one can assume that

$$H_{A,W} = D_3^2 + D_2^2 + (D_1 - Bx_2)^2 - W. \tag{8.B.3}$$

Consider $f \in sL^2$; then $\|E(\lambda)f\| \le \|f\|$ and then one can prove easily (8.B.2) and that

$$\|H_{A,0}E(\lambda)f\| \le (CZ^2 + \lambda_+)\|f\|. \tag{8.B.4}$$

Really, $\frac{1}{2}D_3^2 + CZ^2 \ge W$ in the operator sense.

Then (8.B.4) implies that in $B(0,r)\times B(y',r')$ with $r' = B^{-\frac{1}{2}}$

$$|P^{\alpha}E(\lambda)f| \le CZ^{\alpha_3}B^{\frac{1}{2}|\alpha'|} \qquad \forall \alpha : |\alpha| \le 2\ \forall \lambda \le Z^2$$

with $P = (D_1 - Bx_2, D_2, D_3)$ and therefore $\|E(\lambda)f\|_{\mathscr{C}} \le CZ^{\frac{1}{2}}B^{\frac{1}{2}}\|f\|$. Then $\|E(x,.,\lambda)\|_{\mathscr{L}^2_y}\| \le CZ^{\frac{1}{2}}B^{\frac{1}{2}}$.

Repeating the same arguments with respect to y we arrive to (8.B.1).□

Corollary 8.B.2. *In the framework of Proposition 8.B.1 with* $\phi \in \mathscr{L}^\infty(B(0,r))$, $\|\phi\|_{\mathscr{L}^\infty} \le 1$

$$|\int \phi(x) e(x,x,0)\,dx| \le CZ^{-2}B, \tag{8.B.5}$$

$$\mathsf{D}\bigl(\phi(x)e(x,x,0), \phi(x)e(x,x,0)\bigr) \le CZ^{-3}B^2 \tag{8.B.6}$$

and

$$|\int_{-\infty}^{0} \int \phi(x) e(x,x,\tau)\,d\tau dx| \le CB. \tag{8.B.7}$$

8.C Riemann sums and integrals

If $f \in \mathscr{C}^\infty(\mathbb{R}^+)$ and fast decays at $+\infty$ then

$$f(0)h + \sum_{n\ge 1} 2f(2nh)h \sim \int_0^\infty f(t)\,dt + \sum_{m\ge 1} \kappa_m f^{(2m-1)}(0) h^{2m}, \tag{8.C.1}$$

$$\sum_{n\ge 0} 2f((2n+1)h)h \sim \int_0^\infty f(t)\,dt + \sum_{m\ge 1} \kappa'_m f^{(2m-1)}(0) h^{2m} \tag{8.C.2}$$

as $h \to +0$. The proofs of both formulae follow from Taylor decomposition and observation that the odd powers of h should disappear. Taking $f(t) = e^{-tz/h}$ with $\operatorname{Re} z > 0$ we arrive to

$$1 - \frac{\cosh(z)}{\sinh(z)} z \sim \sum_{m\ge 1} \kappa_m z^{2m}, \tag{8.C.3}$$

$$1 - \frac{1}{\sinh(z)} z \sim \sum_{m\ge 1} \kappa'_m z^{2m}. \tag{8.C.4}$$

as $|z| \ll 1$. In particular, $\kappa_1 = \frac{1}{3}$ and $\kappa'_1 = -\frac{1}{6}$.

Bibliography

[A] V. I. Arnold: *Mathematical Methods of Classical Mechanics.* Springer-Verlag (1990).

[Ba] V. Bach: *Error bound for the Hartree-Fock energy of atoms and molecules,* Commun. Math. Phys. 147:527–548 (1992).

[Be] R. Benguria: *Dependence of the Thomas-Fermi energy on the nuclear coordinates,* Commun. Math. Phys., 81:419–428 (1981).

[BeL] R. Benguria and E. H. Lieb: *The positivity of the pressure in Thomas-Fermi theory,* Commun. Math. Phys., 63:193–218 (1978).

[BrL] H. Brezis and E. H. Lieb: *Long range potentials in Thomas-Fermi theory,* Commun. Math. Phys. 65, 231-246 (1979).

[E] L. Erdős, Magnetic Lieb-Thirring inequalities. Commun. Math. Phys., 170:629–668 (1995).

[ES3] L. Erdős, J.P. Solovej, *Ground state energy of large atoms in a self-generated magnetic field.* Commun. Math. Phys. **294**, No. 1, 229-249 (2009) arXiv:0903.1816

[EFS1] L. Erdős, S. Fournais, J.P. Solovej: *Stability and semiclassics in self-generated fields.* arXiv:1105.0506

[EFS2] L. Erdős, S. Fournais, J.P. Solovej: *Second order semiclassics with self-generated magnetic fields.* arXiv:http://arxiv.org/abs/1105.0512

[EFS3] L. Erdős, S. Fournais, J.P. Solovej: *Scott correction for large atoms and molecules in a self-generated magnetic field* arXiv:1105.0521

[FS] C. Fefferman and L.A. Seco: *On the energy of a large atom*, Bull. AMS **23**, 2, 525–530 (1990).

[FSW1] R. L. Frank, H. Siedentop, S. Warzel: *The ground state energy of heavy atoms: relativistic lowering of the leading energy correction.* Commun. Math. Phys. **278** no. 2, 549–566 (2008)

[FSW2] R. L. Frank, H. Siedentop, S. Warzel: *The energy of heavy atoms according to Brown and Ravenhall: the Scott correction.* Doc. Math. **14**, 463–516 (2009).

[FLL] J. Fröhlich, E. H. Lieb, and M. Loss: *Stability of Coulomb systems with magnetic fields. I. The one-electron atom.* Commun. Math. Phys. **104** 251–270 (1986)

[GS] G. M. Graf and J. P. Solovej: *A correlation estimate with applications to quantum systems with Coulomb interactions,* Rev. Math. Phys., 6(5a):977–997 (1994). Reprinted in The state of matter a volume dedicated to E. H. Lieb, Advanced series in mathematical physics, 20, M. Aizenman and H. Araki (Eds.), 142–166, World Scientific 1994.

[H] W. Hughes: *An atomic energy bound that gives Scott's correction*, Adv. Math. **79**, 213–270 (1990).

[Ivr1] V. Ivrii, Asymptotics of the ground state energy of heavy molecules in the strong magnetic field. I. Russian Journal of Mathematical Physics, 4(1):29–74 (1996).

[Ivr2] Asymptotics of the ground state energy of heavy molecules in the strong magnetic field. II. Russian Journal of Mathematical Physics, 5(3):321–354 (1997).

[Ivr3] V. Ivrii, Heavy molecules in the strong magnetic field. Russian Journal of Math. Phys., 4(1):29–74 (1996).

[Ivr4] V. Ivrii, Heavy molecules in the strong magnetic field. Estimates for ionization energy and excessive charge 6(1):56–85 (1999).

[Ivr5] Sharp spectral asymptotics for operators with irregular coefficients. Pushing the limits. II. Comm. Part. Diff. Equats., 28 (1&2):125–156, (2003).

[Ivr6] V. Ivrii Sharp spectral asymptotics for operators with irregular coefficients. III. Schrödinger operator with a strong magnetic field, arXiv:math/0510326 (Apr. 11, 2011), 101pp.

[Ivr7] V. Ivrii Local trace asymptotics in the self-generated magnetic field, arXiv:math/1108.4188 (December 23, 2011), 24pp.

[Ivr8] V. Ivrii Global trace asymptotics in the self-generated magnetic field in the case of Coulomb-like singularities, arXiv:math/1112.2487 (December 23, 2011), 19pp.

[Ivr9] V. Ivrii Asymptotics of the ground state energy for atoms and molecules in the self-generated magnetic field, arXiv:math/1112.5538 (December 23, 2011), 11pp.

[Ivr10] V. Ivrii Asymptotics of the ground state energy of heavy molecules and related topics, arXiv:math/0510326 (October 03, 2012), 70pp.

[Ivr11] V. Ivrii *Microlocal Analysis and Sharp Spectral Asymptotics*, in progress: available online at http://www.math.toronto.edu/ivrii/futurebook.pdf

[IS] Ivrii, V. and Sigal, I. M Asymptotics of the ground state energies of large Coulomb systems. Ann. of Math., 138:243–335 (1993).

[L] E. H. Lieb: *Thomas-Fermi and related theories of atoms and molecules*, Rev. Mod. Phys. **65**. No. 4, 603-641 (1981)

[L2] E. H. Lieb: *Variational principle for many-fermion systems*, Phys. Rev. Lett. **46**, 457–459 (1981) and **47** 69(E) (1981)

[L2] *The stability of matter: from atoms to stars* (Selecta). Springer-Verlag (1991).

[LLS] E. H. Lieb, M. Loss and J. P. Solovej: *Stability of Matter in Magnetic Fields*, Phys. Rev. Lett. **75**, 985–989 (1995)

[LO] E. H. Lieb and S. Oxford: *Improved Lower Bound on the Indirect Coulomb Energy*, Int. J. Quant. Chem. **19**, 427–439, (1981)

[LS] E. H. Lieb and B. Simon: *The Thomas-Fermi theory of atoms, molecules and solids*, Adv. Math. **23**, 22-116 (1977)

[LSY1] E. H. Lieb, J. P. Solovej and J. Yngvarsson: *Asymptotics of heavy atoms in high magnetic fields: I. Lowest Landau band regions,* Comm. Pure Appl. Math. 47:513–591 (1994).

[LSY2] E. H. Lieb, J. P. Solovej and J. Yngvarsson:. *Asymptotics of heavy atoms in high magnetic fields: II. Semiclassical regions,* Comm. Math. Phys., 161: 77–124 (1994).

[RS] M. B. Ruskai, M. B. and J. P. Solovej: *Asymptotic neutrality of polyatomic molecules.* In Schrödinger Operators, Springer Lecture Notes in Physics 403, E. Balslev (Ed.), 153–174, Springer Verlag (1992).

[SW1] H. Siedentop and R. Weikard: *On the leading energy correction for the statistical model of an atom: interacting case*, Commun. Math. Phys. **112**, 471–490 (1987)

[SW2] H. Siedentop and R. Weikard: *On the leading correction of the Thomas-Fermi model: lower bound*, Invent. Math. **97**, 159–193 (1990)

[SW3] H. Siedentop and R. Weikard: *A new phase space localization technique with application to the sum of negative eigenvalues of Schrödinger operators*, Ann. Sci. École Norm. Sup. (4), **24**, no. 2, 215–225 (1991).

[Sob1] A. V. Sobolev: *Quasi-classical asymptotics of local Riesz means for the Schrödinger operator in a moderate magnetic field.* Ann. Inst. H. Poincaré, **62** no. 4, 325-360, (1995)

[Sob] A. V. Sobolev: *Discrete spectrum asymptotics for the Schrödinger operator with a singular potential and a magnetic field*, Rev. Math. Phys **8** (1996) no. 6, 861–903.

[SS] J. P. Solovej, W. Spitzer: *A new coherent states approach to semiclassics which gives Scott's correction.* Comm. Math. Phys. **241** (2003), no. 2-3, 383–420.

[SSS] J. P. Solovej, T.Ø. Sørensen, W. Spitzer: *Relativistic Scott correction for atoms and molecules.* Comm. Pure Appl. Math. Vol. LXIII. 39-118 (2010).

[Zh] G. Zhislin, *Discussion of the spectrum of Schrodinger operator for systems of many particles.* Tr. Mosk. Mat. Obs., 9, 81–128 (1960).